

## HW 6

### 3.2.9

**a**

We can rewrite  $S$  as  $S = \{(x_1, \dots, x_{p-1}, (x_1 \dots x_{p-1})^{-1}) | x_i \in G\}$ . Since there are no restrictions on the  $x_i$  and there are  $p-1$  choices of the  $x_i$ , the size of this set is  $|G|^{p-1}$ .

**b**

We prove this for the cyclic permutation  $x_k \dots x_{k-1}, k \in [1, p)$  by induction on  $k$ , where the base case  $k = 1$  holds by definition of  $S$ . For the inductive step, we are given  $x_k \dots x_{k-1} = 1$ . Multiplying by  $x_k^{-1}$  on the left, we have  $x_{k+1} \dots x_{k-1} = x_k^{-1}$ . Multiplying by  $x_k$  on the right, we have  $x_{k+1} \dots x_k = 1$ .

**c**

Notation: let  $S_p$  (the symmetric group on  $[1, p]$ ) act on  $S$  by permuting the indices, that is for  $\tau \in S_p$  we have  $\tau \cdot (x_1, \dots, x_p) = (x_{\tau(1)}, \dots, x_{\tau(p)})$ .

I will assume that a cyclic permutation of  $(x_1, \dots, x_p)$  is  $\sigma^j \cdot (x_1, \dots, x_p) = (x_{\sigma^j(1)}, \dots, x_{\sigma^j(p)})$  where  $\sigma^j \in S_p$  is some power of the  $p$ -cycle  $\sigma = (1, \dots, p)$ .

Reflexive: this holds because  $\sigma^0$  is the identity.

Symmetric: this holds because  $\sigma^{-1} = \sigma^{p-1}$ .

Transitive: this holds because  $\sigma^a \sigma^b = \sigma^{a+b}$ .

**d**

$\Leftarrow$  : clearly  $(x, \dots, x) \in S$ . Every cyclic permutation is also of the form  $(x, \dots, x)$ . Hence the equivalence class has exactly 1 element.

$\Rightarrow$  : let  $E = \{(x_1, \dots, x_k)\}$  be the equivalence class. For any  $k \in [1, p)$  we have  $(x_1, \dots, x_k) = (x_k, \dots, x_{k-1})$  since both the LHS and RHS belong to  $E$ , hence  $x_1 = x_k$ . Hence  $x_1 = x_2 = \dots = x_p$ .

**e**

Let  $E$  be the equivalence class, and fix some  $X = (x_1, \dots, x_p) \in E$ . Let  $j$  be the smallest positive integer such that  $\sigma^j \cdot X = X$ . We have  $j \leq p$  since  $\sigma^j = e$ . If  $j = p$  then all  $p$  cyclic permutations are distinct and  $|E| = p$ . Otherwise,  $j$  and  $p$  are coprime so the sequence  $\sigma, \sigma^j, \sigma^{2j}, \dots$  contains every

power of  $\sigma$ . Every cyclic permutation of  $X$  is  $\sigma^k \cdot X$ , and there exists  $t$  such that  $\sigma^k = \sigma^{jt}$ , and  $\sigma^k \cdot X = \sigma^{jt} \cdot X = \sigma^j \cdot \sigma^j \dots \cdot X = X$ .

Since  $S$  is a disjoint union of its equivalence classes, we have  $|S| = |G|^{p-1} = \sum_E |E| = \sum_{E, |E|=1} |E| + \sum_{E, |E|=p} |E| = k + pd$  where the summation is over equivalence classes.

**f**

Consider the equation  $|G|^{p-1} = k + pd$  modulo  $p$ . The LHS is 0 since  $p$  divides  $|G|$  and the RHS is  $k$ . Hence  $p|k$ . Since  $p \geq 2$  and  $k \geq 1$  (since we have at least one equivalence class of size 1) we have  $k > 1$ , hence there are at least two equivalence classes of size 1, hence at least one of the form  $\{x, \dots x\}$  where  $x \neq 1$ . By the definition of  $G$ ,  $x$  satisfies  $x^p = 1$ .