

## HW 12

### 5.4.14

We show that  $G = D \times U$  as an internal direct product.  $D$  and  $U$  are both contained in  $G$ , and every element of  $d$  commutes with every element of  $u$ . Given  $g \in G$  whose diagonal elements are  $g_0$ , we have  $g_0 > 0$  since otherwise, the determinant (which is the product of the diagonal elements for an upper triangular matrix) is 0, which contradicts  $G \subseteq GL_n(F)$ . We can write  $g = du$  where  $d = g_0 I \in D, u \in U$ .

Claim: this decomposition is unique. Suppose  $g = d_1 u_1 = d_2 u_2$ . Let  $d_i = k_i I$  for some scalars  $k_i$ . Then  $k_1 u_1 = k_2 u_2$ . The upper left element of the LHS is  $k_1$  and that of the RHS is  $k_2$ , hence  $k_1 = k_2$ , hence  $d_1 = d_2$  and  $u_1 = u_2$ . taking determinants,  $|d_1| = |d_2|$  hence they have equal diagonal elements, hence  $d_1 = d_2$ , hence  $u_1 = u_2$ .

The map  $\phi : H \times K \rightarrow G$  given by  $(h, k) \mapsto hk$  is a homomorphism. It maps  $(e, e)$  to  $e$ , it maps  $(h, k)^{-1} = (h^{-1}, k^{-1})$  to  $(hk)^{-1} = h^{-1}k^{-1}$ , and it maps  $(h, k)(h', k') = (hh', kk')$  to  $hh'kk' = (hk)(h'k')$ . There is a map  $G \rightarrow H \times K$  given by the unique decomposition. This is an inverse map since the composed map maps  $hk \rightarrow (h, k) \rightarrow hk$ , hence  $\phi$  is an isomorphism.

### 5.5.5a

By definition,  $G = H \rtimes K$  where  $H = Z_2 \times Z_2$  and  $K = \text{Aut}(H)$ . Let  $Z_2 = \{e, x, y, xy\}$ . Any automorphism of  $H$  fixes  $e$ , hence  $K$  is a subgroup of  $S_{\{x, y, xy\}}$ . In fact, since  $Z_2$  is generated by  $x, y$  with  $x^2 = y^2 = (xy)^2 = e$  which is symmetric in  $x, y, xy$ , any permutation of  $x, y, xy$  is an automorphism of  $H$  (e.g. the permutation  $(x \ y)$  gives the homomorphism  $e \mapsto e, x \mapsto y, y \mapsto x, xy \mapsto xy$ ). Hence  $K = S_{\{x, y, xy\}}$ .

$$|G| = |H||K| = 4 \times 6 = 24.$$

### 5.5.5b

Notation: given  $v \in H, \sigma \in K$  let the action of  $K$  on  $v$  that defines the semidirect product be written  $v^\sigma$ , and let elements of  $K$  be written as  $v\sigma$  where  $v \in H, \sigma \in K$ . Then the multiplication rule for  $G$  can be written as  $v\sigma_1 w\sigma_2 = vw^{\sigma_1} \sigma_1 \sigma_2$ .

Consider  $H$  as a subgroup of  $K$ . There are 4 left-cosets which are  $C = \{K, xK, yK, xyK\}$ .  $G$  acts on  $C$  by left multiplication; denote this action by  $g \cdot wK$  where  $g \in G, w \in H$ . This action has a permutation representation  $\pi : G \rightarrow S_C$ . Write  $g = v\sigma$ . Then  $g \cdot wK = v\sigma \cdot wK = vw^{\sigma} \sigma K = vw^{\sigma} K$ .

Notation: when writing elements of  $\text{im}\pi$  as permutations, we shall replace  $vK$  with  $v$ .

Considering the elements  $g$  for which  $v = e$ , we see that  $g \cdot wK = w^\sigma K$ . Hence  $\text{im}\pi$  contains  $S_{\{x, y, xy\}}$ .

Considering the elements of  $g$  for which  $\sigma$  is the identity automorphism, we see that  $g \cdot wK = vwK$ . Hence  $\text{im}\pi$  contains  $(e \ x)(y \ xy)$  as the image of  $x$ ,  $(e \ y)(x \ xy)$  as the image of  $y$ , and  $(e \ xy)(x \ y)$  as the

image of  $xy$  (these are the permutations that correspond to the left regular representation of the action of  $H$  on itself by left multiplication)

We know  $\text{im}\pi$  contains  $(x\ y)$ . Conjugating this by  $(e\ x)(y\ xy)$  we see that it contains  $(e\ xy)$ . Conjugating this by elements of  $S_{\{x,y,xy\}}$  we see that it contains  $(e\ x)$  and  $(e\ y)$ . Hence  $\text{im}\pi$  contains all the transpositions of  $S_C$ , hence  $\text{im}\pi = S_C$ , which means there is a surjective homomorphism  $G \rightarrow S_4$ ; since  $|G| = |S_4|$ , this is an isomorphism.

### 5.5.16

I will use  $C_k$  for the cyclic group of order  $k$ .

Considering  $C_8$  as the additive group of  $\mathbb{Z}/8\mathbb{Z}$ , every automorphism of  $C_8$  is multiplication by a unit of  $\mathbb{Z}/8\mathbb{Z}$ , i.e. the automorphisms are precisely multiplication by 1, 3, 5, or 7. Every homomorphism  $C_2 \rightarrow C_8$  is determined by the image of the nonidentity element of  $C_2$ , and an automorphism of  $C_8$  is a valid target if it is self-inverse. It is easy to check that each of those four automorphisms is self-inverse (since  $-1, 3, -3$  all square to 1 modulo 8).

Multiplication by one: this is the semidirect product  $C_2 \rtimes C_8$  where the action of  $C_2$  on  $C_8$  is trivial, hence it is the direct product  $C_2 \times C_8$ .

Multiplication by 7 = -1: this is the semidirect product  $C_2 \rtimes C_8$  where the action of  $C_2$  on  $C_8$  is inversion, hence it is the dihedral group.

Multiplication by 3: use exponential notation for the action. The multiplication rule is  $\sigma_1\tau_1\sigma_2\tau_2 = \sigma_1\sigma_2\tau_1^{\sigma_2}\tau_2$ . Let  $\sigma_1 = e, \tau_2 = e$  and let  $\tau, \sigma$  be generators of  $C_8$  and  $C_2$ . The multiplication rule simplifies to  $\tau\sigma = \sigma^\tau\tau = \sigma^3\tau$ . Hence  $\tau\sigma^3 = \sigma^3\tau\sigma^2 = \sigma^6\tau\sigma = \sigma^9\tau = \sigma\tau$ , hence this group has the presentation  $\langle \sigma, \tau \mid \sigma^8 = \tau^2 = e, \sigma\tau = \tau\sigma^3 \rangle$  which is the same presentation as 2.5.11.

Multiplication by 5: let  $u, v$  be generators of  $C_2$  and  $C_8$ ; similar to the above, we have  $uv = v^5u$ . Hence  $uv^5 = v^5uv^4 = v^{10}uv^3 = \dots = v^{25}u = vu$  which is the same presentation in 2.5.14.

### 5.5.18

Let  $H$  be a group; by Cayley's theorem it is a subgroup of  $S_H$ , namely the subgroup of permutations  $P = \{\pi_h : h' \mapsto hh' \mid h \in H\}$ . Take  $G = N_{S_H}(H)$ ; then  $H$  is a normal subgroup of  $G$  by construction. Let  $\sigma$  be an automorphism of  $H$ ; in particular  $\sigma$  is a permutation of  $H$  so we can treat  $\sigma$  as an element of  $S_H$ . The conjugation  $\pi_h^\sigma$  is the map  $\sigma \circ \pi_h \circ \sigma^{-1}$  which maps  $h'$  to  $\sigma(h\sigma^{-1}(h')) = \sigma(h)h'$  hence it is the map  $\pi_{\sigma(h)}$ . Hence conjugation by  $\sigma$  is an automorphism of  $P$  (this also shows  $\sigma \in G$ ).

### 5.5.22a

We apply the recognition theorem (theorem 12):

$U \trianglelefteq G$ : when multiplying upper-triangular matrices, the diagonal entries are multiplied with each other.

Hence,  $\begin{pmatrix} a & \cdots & \cdots \\ 0 & b & \cdots \\ & & \ddots \end{pmatrix}^{-1} = \begin{pmatrix} a^{-1} & \cdots & \cdots \\ 0 & b^{-1} & \cdots \\ & & \ddots \end{pmatrix}$  and conjugation by an upper triangular matrix leaves the diagonal values unchanged.

$U \cap D = \{I\}$ : let  $M$  be in the intersection.  $M$  is a diagonal matrix, and all of its diagonal elements are 1, hence it is the identity matrix.

$UD = G$ : note that diagonal matrices in  $GL_n(F)$  are invertible, with

$$\begin{pmatrix} a & 0 & \cdots \\ 0 & b & \cdots \\ & & \ddots \end{pmatrix}^{-1} = \begin{pmatrix} a^{-1} & 0 & \cdots \\ 0 & b^{-1} & \cdots \\ & & \ddots \end{pmatrix}$$

where every diagonal element is nonzero, since otherwise the determinant would be 0. Given an arbitrary matrix  $\mathcal{G} = \begin{pmatrix} a & \cdots & \cdots \\ 0 & b & \cdots \\ & & \ddots \end{pmatrix}$  let  $\mathcal{D} = \begin{pmatrix} a & 0 & \cdots \\ 0 & b & \cdots \\ & & \ddots \end{pmatrix}$ ; we have  $\mathcal{G}\mathcal{D}^{-1}$  is an upper triangular matrix with diagonal elements 1; call this matrix  $\mathcal{U}$ . Then  $\mathcal{G} = \mathcal{U}\mathcal{D}$ .

### 5.5.22b

Elements of  $U$  are of the form  $\mathcal{U} = \begin{pmatrix} 1 & x \\ 0 & 1 \end{pmatrix}$  where we can take  $x \in F$  to be the image of the isomorphism.

Similarly elements of  $D$  are of the form  $\mathcal{D} = \begin{pmatrix} a & 0 \\ 0 & b \end{pmatrix}$  where  $a, b \in F^\times$ .

The action is given by conjugation of  $\mathcal{U}$  by  $\mathcal{D}$ ; we have  $\mathcal{D}\mathcal{U}\mathcal{D}^{-1} = \begin{pmatrix} 1 & a^{-1}xb \\ 0 & 1 \end{pmatrix}$ . Hence the action is given by  $(a, b) \circ x = a^{-1}bx$ .