



Chapter 10: Treatment of active tuberculosis in special populations

Ryan Cooper, Stan Houston, Christine Hughes & James C. Johnston

To cite this article: Ryan Cooper, Stan Houston, Christine Hughes & James C. Johnston (2022) Chapter 10: Treatment of active tuberculosis in special populations, Canadian Journal of Respiratory, Critical Care, and Sleep Medicine, 6:sup1, 149-166, DOI: [10.1080/24745332.2022.2039500](https://doi.org/10.1080/24745332.2022.2039500)

To link to this article: <https://doi.org/10.1080/24745332.2022.2039500>



Published online: 25 Mar 2022.



Submit your article to this journal [↗](#)



Article views: 6083



View related articles [↗](#)



View Crossmark data [↗](#)



Citing articles: 1 View citing articles [↗](#)

Chapter 10: Treatment of active tuberculosis in special populations

Ryan Cooper^a, Stan Houston^{a,b}, Christine Hughes^c and James C. Johnston^{d,e}

^aDepartment of Medicine, Division of Infectious Diseases, University of Alberta, Edmonton, Alberta, Canada; ^bSchool of Public Health, University of Alberta, Edmonton, Alberta, Canada; ^cFaculty of Pharmacy & Pharmaceutical Sciences, University of Alberta, Edmonton, Alberta, Canada; ^dDivision of Respiratory Medicine, University of British Columbia, Vancouver, British Columbia, Canada; ^eTB Services, British Columbia Centre for Disease Control, Vancouver, British Columbia, Canada

KEY POINTS

- The management of tuberculosis (TB) in people with medical co-morbidities, substance-use disorders, advanced age or pregnancy is challenging; consultation with a TB expert is recommended.
- Adverse events and treatment interruption are more common in these special populations; close monitoring and additional support is often needed.
- Given alterations in TB-drug pharmacokinetics and the potential for severe drug-drug interactions in patients with medical co-morbidities, we recommend consultation with an experienced pharmacist.
- TB disease and its treatment have important implications for the management of co-morbid conditions; close collaboration with medical specialists and allied health professionals is often required.
- The prolonged course of TB treatment is a useful opportunity to screen for co-morbidity and to facilitate referral and linkage into care.

1. Introduction

Medical co-morbidities, substance use-disorders, advanced age and pregnancy can complicate management of TB.^{1,2} Alterations in immunologic control of TB infection may increase the risk of reactivation and lead to more severe forms of TB disease.¹ Co-morbidity may also lead to slower TB treatment response and higher rates of TB relapse.¹ As a result, extension of TB treatment duration is sometimes required to optimize TB treatment outcomes.

These special populations may also differ in absorption, metabolism and clearance of TB drugs. Thus, special attention to dosing of TB medications is needed and, in some groups, therapeutic drug monitoring is recommended.³ Increased susceptibility to adverse events in some groups may mandate closer monitoring and support, including for drug-drug interactions and overlapping toxicities.

Finally, there is often a lack of comparative clinical data directly relevant to these special populations, as these groups

are often excluded from clinical trials due to concerns over harm.

Given these challenges and limitations in the clinical data, consultation with a TB specialist and an experienced pharmacist is strongly recommended. Furthermore, it is important to initiate close collaboration with the patient's medical and allied health care team throughout the TB treatment course.

Here we review additional considerations in TB management of special populations. For each population we will review the impact on epidemiology, clinical presentation, treatment outcomes and management.

2. People with human immunodeficiency virus

Human immunodeficiency virus (HIV) infection increases the risk of TB disease nearly 100-fold.^{4,5} Among people with HIV and *M. tuberculosis* coinfection, the annual risk of active TB may be as high as 10 per 100 person years.^{5,6} Antiretroviral therapy (ART) reduces the incidence of active TB substantially, although the incidence remains higher than in people without HIV infection, even after normal CD4+ lymphocyte counts are attained.^{7,8}

The predominant immunologic effect of HIV is to reduce cell-mediated immune function. By reducing the number of T-helper cells, macrophage activation and granuloma formation is impaired, compromising the immunologic containment of latent and new TB infections.⁹

HIV also alters the clinical and radiologic features of TB, which are partly determined by the host response.¹⁰ Extra-pulmonary and disseminated forms of TB are more common in people with HIV infection, especially in those with CD4+ lymphocyte counts below 50 × 10⁶/L, while cavitary lung disease and sputum smear-positive disease is less common.¹¹ This atypical presentation can contribute to diagnostic delay.

HIV infection also affects TB treatment outcomes. Treatment failure with acquired rifampin mono-resistance has been observed with intermittent treatment regimens in

people with HIV, particularly among people with CD4 counts $<100 \times 10^6/L$.^{12,13} TB recurrence is also more common among people with HIV.¹⁴ When molecular techniques have been used to distinguish between relapse and reinfection in communities with high levels of ongoing transmission, however, the rates of relapse with the original strain have been similar.^{15,16} Mortality is higher among people with both HIV and TB and correlates with the degree of immune suppression.¹⁷ With appropriate anti-TB therapy and timely initiation of ART, however, the difference in outcomes can be attenuated.¹⁶

There are several special considerations in the management of TB and HIV co-infection, including TB drug malabsorption, potential for profound drug-drug interactions and the avoidance of immune reconstitution inflammatory syndrome.

Recommendation

- **We strongly recommend that, due to the profound impact of human immunodeficiency virus (HIV) on patient survival and TB treatment outcomes, all patients with TB be screened for HIV infection (*good evidence*).**

Good practice statement

- **Treatment of TB in people with human immunodeficiency virus (HIV) should be guided by a physician with expertise in the management of both diseases or in close collaboration with a physician expert in HIV care. Consultation with an expert pharmacist is also advised.**

2.1. TB therapy in people with HIV

The anti-TB regimen choice and duration is the same for people with HIV as for those without HIV, except for the selective use of rifabutin in place of rifampin when necessary to achieve compatibility with the ART regimen (see [Table 1](#)).^{16,18–24}

Daily administration of TB drugs throughout the treatment course is recommended, as intermittent therapy is associated with worse treatment outcomes, including treatment failure, relapse and acquired drug resistance.^{16,25} Treatment duration is not extended on the basis of HIV co-infection alone. However, in the uncommon scenario where a patient declines to take ART, TB treatment should be extended to 9 months.^{16,21}

Several studies have found that a substantial proportion of people with both HIV and TB infection have low serum concentrations of anti-tuberculous agents.^{26–31} This is thought to be due to a combination of factors, including drug interactions with ART and decreased absorption related to gastrointestinal dysfunction associated with HIV infection.³² Low serum drug concentrations have been linked to slower response to TB treatment^{21,29,33,34} and acquired rifamycin resistance.³⁵ Thus, we frequently monitor serum drug concentrations in individuals with HIV co-infection to optimize dosing.

HIV-infected individuals are already at increased risk of neuropathy related to HIV and ART. Therefore, in those taking isoniazid (INH), vitamin B6 supplementation is routinely used to prevent additional neurologic toxicity from INH-associated neuropathy.

Recommendations

- **We strongly recommend a rifamycin (rifampin or rifabutin)-containing regimen for treatment of TB, despite the potential for drug-interactions with antiretroviral therapy (*good evidence*).**
- **We strongly recommend that TB treatment in human immunodeficiency virus co-infected individuals be administered daily throughout (*good evidence*).**
- **We conditionally recommend that, if a patient is not taking antiretroviral therapy, TB treatment be extended to 9 months (*poor evidence*).**
- **We conditionally recommend that, where available, serum TB drug concentrations be measured in people with human immunodeficiency virus co-infection and used to optimize TB drug dosing (*poor evidence*).**

2.2. Antiretroviral therapy in people with HIV and TB

ART is strongly recommended for all people with HIV who also have TB.²³ The optimal timing of ART initiation in people receiving TB treatment, balancing the risk of progressive HIV and TB disease with the risk of immune reconstitution inflammatory syndrome (IRIS), has been evaluated in nine randomized-controlled trials.^{36–38} Two systematic reviews of these trials concluded that early initiation of ART, within 2 weeks of TB therapy initiation, reduces overall mortality and the incidence of additional acquired immunodeficiency syndrome (AIDS)-defining illnesses. The benefit of early initiation of ART was most apparent in patients with CD4 counts of $<50 \times 10^6/L$.

As an important caveat, early initiation of ART in cases of central nervous system (CNS) TB may be hazardous, probably because of the unique risks of IRIS reactions in the closed space of the cranium. One randomized study of TB meningitis compared immediate ART with a 2-month delay in ART initiation and found higher rates of severe adverse effects in the immediate ART group. However, the mortality rate, which was more than 50%, did not differ between arms.³⁹ Thus, in those with CNS-TB, the optimal time to initiate ART is not well-established. Delaying therapy at least 2 weeks after initiation of TB treatment appears to provide the best balance between avoiding cerebral complications of IRIS and improved TB treatment outcomes (see [Chapter 7: Extra-pulmonary Tuberculosis](#)).

In patients already receiving effective combination ART at the time of the TB diagnosis, ART should be continued. However, a change in ART regimen may be required to accommodate potential drug interactions with rifamycins. Note that when an ART dose adjustment is made to address an interaction with a rifamycin, the increased dose should

be maintained for 2 weeks after stopping the rifamycin to allow for the liver enzyme induction effect to wear off.

Recommendations

- We strongly recommend initiating antiretroviral therapy within 2 weeks of starting TB therapy, provided there is no documented central nervous system TB (*good evidence*).
- We conditionally recommend delaying antiretroviral therapy (ART) initiation for at least 2 weeks after initiation of anti-TB treatment for those with central nervous system TB, although here the optimal time to initiate ART is less well-established and expert consultation is advised (*poor evidence*).

2.3. Specific antiretroviral regimens in patients taking Rifamycin-containing TB treatment

Rifamycins are the only anti-tuberculous agents to exert clinically important interactions with antiretroviral drugs (see Table 1). Rifabutin is associated with weaker enzyme induction and thus has less potential for serious drug-drug interactions than either rifapentine or rifampin. However, there is less published clinical experience with rifabutin in the treatment of people with both HIV and TB, and rifampin is usually preferred in this population.^{23,38}

Since rifamycin-based TB treatment is strongly recommended in people with HIV and TB (see recommendation in section 2.1), these drug-drug interactions substantially limit the number of compatible ART regimens available. Current recommendations from the World Health Organization (WHO) for people newly diagnosed with HIV and on treatment for TB is dose-adjusted dolutegravir plus 2 nucleoside analogues. An acceptable alternative is efavirenz plus 2 nucleoside analogues (see Table 1 for dosing for both regimens).³⁸

The management of drug-drug interactions between ART and the rifamycin class is an area of active research and recommendations change frequently. Consultation with an experienced pharmacist and a regularly updated clinical drug-interaction resource is prudent (see section 13 on drug-drug interactions).

It is crucial that, while on TB treatment, ongoing monitoring of antiretroviral efficacy is performed. ART efficacy can be compromised in people with HIV and TB co-infection because of drug-drug interactions, reduced adherence related to increased pill burden or overlapping toxicity. Thus, we strongly advise close monitoring of plasma HIV ribonucleic acid (RNA) in people on TB treatment. Monthly testing is recommended until plasma viral load is no longer detectable and then, at minimum, quarterly testing while on TB treatment.

In patients with a suboptimal virologic response to ART in whom an interaction or decreased absorption is a possible explanation, measurement of serum antiretroviral concentrations should be considered, although clinical evidence to support this strategy is lacking. Adherence to ART should also be optimized and antiviral resistance excluded.

Good practice statement

- Close monitoring of plasma human immunodeficiency virus (HIV) RNA in people on treatment for both TB and HIV is suggested with monthly measurements until plasma HIV RNA is no longer detected, followed by quarterly testing during the course of TB treatment.

2.4. Additional considerations in selecting a compatible antiretroviral regimen for people on TB treatment

There are no significant interactions between the nucleoside analogue class and the rifamycin class and these can be used together without dose adjustment. Tenofovir alafenamide serum concentrations are reduced with rifampin but intracellular concentrations of the active form of the drug appear adequate.⁴⁰

All the other antiretroviral drug classes, including protease inhibitors (PI), non-nucleoside reverse transcriptase inhibitors (NNRTI), integrase inhibitors (INI) and CCR5 receptor blockers, demonstrate major interactions with the rifamycins and caution is required.

Dolutegravir and raltegravir may be used with rifampin, provided doses are increased to account for the enhanced metabolism (see Table 1).^{41–44} It should be noted, however, that twice-daily dosing of some components of an ART regimen adds significant complexity and pill-burden and that this can challenge consistent adherence to ART. In a recent randomized controlled trial involving people with HIV and TB receiving rifampin-based treatment, twice-daily raltegravir was less effective at controlling HIV than an efavirenz-based, once-daily regimen.⁴¹ The difference was largely attributed to reduced adherence to the more complicated ART regimen in the raltegravir arm.⁴¹ Although a single, small phase-one study suggests that dolutegravir can be given once daily with rifabutin, there is less published clinical experience with this regimen.⁴⁵ Other members of the integrase-inhibitor class — bictegravir and elvitegravir — cannot be used with the rifamycins.²³

Extensive experience and a controlled trial has shown that the NNRTI efavirenz at standard dosing of 600 mg/day remains effective when used with rifampin, despite variable reduction in efavirenz serum concentrations.⁴⁶ An increase in dose of efavirenz to 800 mg has been used in patients with larger body mass or suboptimal viral suppression. Rilpivirine and doravirine are not compatible with rifampin but with dosage adjustment can be co-administered with rifabutin.²³

No PI dosing regimen has been found to be safe and effective in combination with rifampin. Rifabutin can be substituted for rifampin in TB treatment to permit the use of PIs but is associated with higher rates of hematologic and ocular toxicity.^{20,47} This is because rifabutin concentrations are increased by concomitant therapy with PIs. There is greater experience with ritonavir than cobicistat “boosting” of PIs.

Table 1. Summary of compatible rifamycin-based anti-tuberculous therapy and antiretroviral regimens (last updated September 2021).

Rifamycin	Anchor drug in antiretroviral regimen	Nucleoside analogue component of anti-retroviral regimen
Rifampin 600 mg daily	Efavirenz 600 mg daily ^{38,46} Dolutegravir 50 mg twice daily ^{41,43} Raltegravir 800 mg twice daily ⁴²	No clinically important interactions expected ^{23,38}
Rifabutin 150 mg daily	Ritonavir "boosted" protease inhibitor ^{23,48}	No clinically important interactions expected ^{23,38}
Rifabutin 300 mg daily	Dolutegravir 50 mg daily ⁴⁵ Raltegravir 400 mg twice daily ^{42,49} Rilpivirine 50 mg daily ²³ Doravirine 100 mg twice daily ²³	No clinically important interactions expected ^{23,38}

Note: Expert consultation is strongly advised when selecting antiretroviral therapy (ART) regimens in people on tuberculosis treatment.

2.5. Immune-Reconstitution Inflammatory Syndrome

IRIS is a frequent early complication of ART in people with HIV and TB. There are two types of IRIS. The first occurs during TB therapy, after ART initiation, and is known as paradoxical IRIS. The second occurs following ART initiation in patients with unrecognized TB, and is known as unmasking IRIS.^{47,50,51}

IRIS has been reported with a frequency ranging from 8 to 43%.⁵² IRIS usually presents as fever and disease progression at involved sites, for example as enlarging lymph nodes, worsening pulmonary infiltrates on chest radiograph or exacerbation of inflammatory changes at other sites. Mortality attributed to IRIS appears to be uncommon except in cases with CNS involvement. Most affected patients have low initial CD4 cell counts, typically $100 \times 10^6/L$ or less. Onset has been described between 2 and 40 days after ART initiation.

Diagnosis of IRIS requires exclusion of other possible causes, including treatment failure due to drug resistance or development of a different opportunistic infection.⁵³

Treatment of IRIS is not always required, as the condition is self-limited. However, if the symptoms are severe enough to warrant therapy, corticosteroids such as prednisone at doses in the range of 1 mg/kg of body weight, given over four weeks, have been shown effective in a randomized trial.⁵⁴ Most people can be managed successfully without interruption of ART or TB treatment and in any case, such interruption will not hasten resolution of IRIS.

A single randomized-controlled study has shown that a short course of prednisone given at the same time as ART initiation can reduce the frequency of symptomatic IRIS without increasing the risk of other opportunistic illness.⁵⁵ People who should be considered for this intervention are those at high-risk of paradoxical IRIS, defined by CD4 count less than $100 \times 10^6/L$. It is important to ensure that patients are responding to TB therapy prior to using preemptive prednisone and that rifampin resistance is excluded. The presence of Kaposi's sarcoma or active hepatitis B infection are additional contraindications.⁵⁵

Recommendation

- We strongly recommend, in people with HIV responding to TB therapy and who are about to initiate antiretroviral therapy, and whose CD4 count is less than 100×10^6 , prednisone 40 mg/d for 14 days followed by 20 mg/d for 14 days to reduce the risk of symptomatic

immune reconstitution inflammatory syndrome, unless rifampin-resistance, Kaposi's sarcoma, and active hepatitis B infection are present as contraindications (good evidence).

Diagnostic considerations in people with both HIV and TB are discussed in Chapter 3: Diagnosis of TB disease and Drug-Resistant TB and Chapter 4: Diagnosis of TB Infection. Regimens used for treatment of latent TB infection in people with HIV are discussed in the Chapter 6: TB Preventive Treatment in Adults.

3. People who have undergone solid organ transplant

Recipients of solid-organ transplantation have a substantially elevated risk of TB that can range between 4 and 30 times that of the general population.⁵⁶ This is a result of the potent impact of anti-rejection medications on host T-cell function and cell-mediated immune response.⁵⁷ Most commonly, TB in this population represents reactivation of latent infection in the recipient. However, transmission of unrecognized infection in the allograft (donor-derived infection) and acquisition of new infection post-transplantation can also occur.

TB in transplant recipients is frequently atypical in its clinical presentation.^{58–60} Disseminated disease is present in up to 15–30% at TB diagnosis and cavitary disease is less frequent in this population.^{58,61} TB often occurs within the first year following organ transplantation, reflecting both reactivation of previous infection and the period of the most intense immune suppression. Donor-derived TB (ie, TB arising from the transplanted organ) typically presents within three months of organ transplantation.

TB treatment is more difficult and outcomes are worse in solid-organ transplant recipients compared to other populations.^{58,62} Overlapping drug toxicity and drug-drug interactions lead to adverse events in nearly one-third of transplant recipients receiving TB therapy.^{58,63} Graft dysfunction and organ rejection occur more frequently in transplant recipients with TB compared to those without TB; this is related in part to drug-drug interactions reducing the serum concentrations of immune-suppression therapies. Mortality from TB is higher in transplant recipients, with rates up to 20% in modern cohort studies.^{58,63,64}

Drug-induced liver injury occurs frequently in transplant recipients, with rates especially high in liver recipients. INH

can still be included as part of a first-line regimen, provided liver enzymes and liver function are closely monitored. Pyrazinamide (PZA) should be avoided in liver transplant recipients.^{59,63,65}

Despite potential for drug-interactions, most experts recommend rifamycin-based for treatment of TB in transplant recipients.^{60,65–67} However, interactions between TB and immunosuppressive agents necessitates close collaboration with transplant physicians and pharmacists.^{60,66}

Rifabutin, which exerts less potent drug-interactions than rifampin, is the preferred rifamycin for treatment of active TB in this population.^{68,69} The efficacy of rifabutin against TB has been demonstrated in clinical trials (albeit in non-transplant recipients) and is considered comparable to rifampin^{20,70–73} (see [Chapter 5: Treatment of Tuberculosis Disease](#)). Importantly, rifabutin use appears to lessen the risk of graft rejection in transplant recipients.⁷⁰ Regular therapeutic drug monitoring of anti-rejection medications is recommended while on a rifamycin-containing TB treatment.

Dose reduction of immune suppression in transplant recipients is not needed to achieve cure of TB when using rifamycin-based TB treatment regimens. Furthermore, aggressive dose reduction of anti-rejection medications could potentially lead to an inflammatory immune-reconstitution syndrome that in some cases can be severe or life-threatening.⁷⁴

Mold-active azoles (eg, voriconazole, posaconazole) are used as prophylaxis against fungal infection in some transplant recipients. Serum concentrations of these agents are substantially reduced by rifamycins. There is also a bi-directional interaction with rifabutin, which risks potential rifabutin toxicity. In some centers, azole prophylaxis is avoided.

Therapeutic drug monitoring of anti-TB drugs is also commonly performed in recipients of solid-organ transplants, due to the potential for altered pharmacokinetics in these complicated populations.

TB drugs for solid-organ transplant recipients should be administered daily throughout the treatment course and these patients should be provided with directly observed therapy (DOT) and/or close supportive care. Treatment extension to nine months is endorsed for this population but direct evidence to support this is lacking.^{67,75} Many experts recommend longer treatment courses in transplant recipients, reasoning that, as in untreated HIV infection, where there is also persistent cell-mediated immune dysfunction, treatment extension has been demonstrated in controlled trials to reduce relapse risk.¹⁶

Recommendations

- **We strongly recommend rifamycin-based therapy for optimal treatment of TB in solid-organ transplant recipients (good evidence).**
- **We conditionally recommend rifabutin over rifampin to reduce the risk of potentially severe drug interactions with anti-rejection medications (poor evidence).**

- **We conditionally recommend, given the potentially severe risk of drug-drug interactions, regular therapeutic drug monitoring of anti-rejection medications (poor evidence).**
- **We conditionally recommend treatment extension to nine months for solid-organ transplant recipients (poor evidence).**

4. People prescribed TNF-alpha inhibitors

TNF-alpha inhibitors (TNFi), including infliximab, adalimumab, etanercept, golimumab and certolizumab, are associated with an elevated risk of reactivation of TB disease. Multiple registry analyses estimate the risk of TB to be more than twice that of patients with inflammatory disease not taking TNFi and up to 20 times higher than the general population.^{76–79}

TB reactivation rates appear higher when TNFi therapy is given along with methotrexate or azathioprine.⁸⁰ Furthermore, within the TNFi class, infliximab and adalimumab appear to convey a higher risk of TB reactivation than etanercept.^{81,82} The introduction of systematic screening for TB infection prior to initiation of treatment with TNFi appears to have reduced, but not eliminated, TB in people receiving these agents.^{79,83,84}

TNF-a is a cytokine essential for the activation of macrophages and the formation and maintenance of granulomas.⁸⁵ The inhibition of TNF-a function leads to reduced immunologic containment of TB infection and can alter the clinical presentation of disease.

Indeed, people with TNFi-associated TB are more likely to have disseminated, meningeal and extra-pulmonary disease and less likely to have cavitary chest disease or sputum smear positivity.^{86,87} This atypical presentation may delay TB recognition.^{87,88} TB in patients treated with TNFi usually occurs within a median of 3 months after starting TNFi therapy.^{87,89}

Sometimes, use of TNFi is delayed in patients who are under evaluation for possible TB before the diagnosis is confirmed, out of concern that the immune suppression may hasten disease progression or lead to dissemination before effective anti-TB therapy can be established.^{90–92} However, it should be noted that abrupt TNFi withdrawal may be associated with exacerbation of TB disease despite anti-TB treatment, somewhat analogous to IRIS seen in HIV-coinfected patients.^{93,94} Vigilance for this effect is prudent and re-introduction of TNFi has been reportedly effective in its management.^{86,92,95,96} Small cohort studies and clinical experience suggest that TNFi can be safely administered once patient is established on effective TB therapy and provided that drug-resistance is not suspected.^{95,97–100}

Despite lack of experimental data, many experts recommend longer treatment courses in patients receiving TNFi, reasoning that, as for people with untreated HIV infection, and thus persistent immune dysfunction, treatment extension may reduce relapse risk.¹⁶ Relapse in people re-started on

TNFi therapy and completing standard TB treatment durations has been reported.^{21,95}

Recommendations

- **We conditionally recommend that, where TNF-alpha inhibitors are stopped following the development of TB disease, it can be re-introduced once the patient is established on effective TB therapy (poor evidence).**
- **We conditionally recommend extending TB treatment to nine months in patients receiving TNF-alpha inhibitors (poor evidence).**

5. People with diabetes

Systematic reviews have estimated a three-fold increase in active TB in people with diabetes^{101,102} and risk appears especially high in those with poor glycemic control, insulin dependence and higher HbA1c.^{103–105}

The clinical manifestations of TB are altered by the presence of diabetes. Observational studies suggest people with diabetes are more likely to have cavitary and sputum smear-positive disease but less likely to have extra-pulmonary disease.^{106–109}

Hyperglycemia and poorly controlled diabetes have also been correlated with worse TB treatment outcomes. Multiple retrospective studies observe that people with both TB and diabetes experience delayed sputum culture conversion and higher rates of treatment failure, relapse, mortality and acquisition of rifampin resistance.^{107,110–114}

The prevalence of diabetes in people with newly diagnosed active TB can range up to 25% when routine testing is employed (see [Chapter 1: Epidemiology of Tuberculosis in Canada](#), Figure 11). This has been demonstrated in both high TB-incidence and low TB-incidence countries.^{115–119} In one Canadian study, diabetes was present in 19.7% of people with active TB.¹²⁰ Thus, screening for diabetes in patients with active TB is recommended and measurement of glycosylated hemoglobin A1C percentage is commonly used.^{121–123}

Good practice statement

- **At the time of diagnosis of TB disease, routine screening for diabetes through measurement of glycosylated hemoglobin A1C percentage is suggested and those diagnosed with diabetes should be linked to a diabetes care provider.**

Optimization of glycemic control is associated with improved TB treatment outcomes.^{109,111,124–126} People diagnosed with diabetes during their TB care should receive a referral to a diabetes care provider for long-term management.

Peak serum TB drug concentrations are frequently low in people with diabetes and this may contribute to the poorer TB treatment outcomes seen in this population.^{3,28,29,33,127} It is not entirely clear how diabetes effects the pharmacokinetics of TB drugs, but increased body weight and gastroparesis with delayed absorption have been suggested as potential factors.^{127–129} At least one observational study has linked the routine measurement of serum

TB drug concentrations in diabetic patients with faster microbiological response to TB treatment.¹³⁰

Longer treatment duration may be required in people with diabetes. In one large country-wide registry of people under treatment for pulmonary TB in Taiwan, higher relapse rates seen in people with diabetes was mitigated when TB treatment was extended to 9 months total.¹⁰⁷ However, the impact of treatment extension was small; of 12,688 people with diabetes and TB, relapse rates were 2.23% in those who received 6 months of treatment and 2.00% in those receiving 9 months (aHR 0.75 (95% CI, 0.59–0.97).

Recommendations

- **We conditionally recommend that for people with poorly controlled diabetes or evidence of gastroparesis, serum TB drug concentrations be used to optimize drug dosing (poor evidence).**
- **We conditionally recommend treatment extension to 9 months for those with diabetes and cavitary pulmonary or disseminated TB disease (poor evidence).**

6. People with chronic kidney disease

End-stage kidney disease requiring dialysis is a well-established risk factor for TB disease, with incidence rates 7 times higher than the general population.^{131–133} Evidence is emerging to suggest that TB risk is also increased in people with earlier stages of chronic kidney disease (CKD), with risk rising as estimated glomerular filtration rate drops below 50 ml/min.^{133–135}

Susceptibility to TB in CKD appears multifactorial.^{136,137} People with CKD often have low 25-hydroxy vitamin D levels and higher rates of protein malnutrition. As uremic waste products accumulate in later stages of renal disease, a broad cellular immune dysfunction develops. The risk of TB may be compounded by associated co-morbid conditions such as diabetes and use of immune suppressive drugs.

The clinical presentation of TB in people with CKD is often insidious and atypical.¹³⁶ Systemic symptoms, such as fever, anorexia and weight loss, may mimic uremia and can result in a delay of diagnosis. People with CKD frequently have extra-pulmonary TB, rather than the more recognizable pulmonary disease.¹³⁸ Delayed recognition may contribute to the higher mortality rates seen in these populations.¹³⁶

A special consideration in this population is to ensure renal-adjusted dosing for some TB drugs.¹³⁶ Rifampin and INH are primarily metabolized and excreted through the liver with little urinary clearance. Thus, dose adjustment of these agents is not required for patients with renal insufficiency.¹³⁸ Although PZA is also metabolized primarily through the liver, some of its metabolites are eliminated renally and so its dose should be adjusted in people with advanced CKD ((Stage 4/5, GFR < 30 ml/min).^{138,139} Ethambutol (EMB) is mostly excreted unchanged by the kidneys and people with advanced CKD have substantially reduced clearance of the drug.¹⁴⁰ Ethambutol-induced ocular toxicity is largely dose-related and so dose adjustment and regular visual acuity testing is necessary in those with advanced CKD.

Because therapeutic efficacy of both PZA and EMB appears dependent on peak concentrations, the dosing interval for both drugs should be extended in people with advanced kidney disease, rather than decreasing the dose administered.

There is insufficient evidence to guide dosing of people with moderate kidney disease (GFR 30-60mL/min). In this range, people should be monitored carefully for toxicity, and therapeutic drug monitoring might be necessary to guide appropriate dosing.

The mechanism of drug removal by peritoneal dialysis is not the same as by hemodialysis and so TB drug-dosing recommendations for people receiving hemodialysis may not necessarily apply. There is comparatively little clinical experience in TB drug-dose adjustment in people receiving peritoneal dialysis.^{141,142} Thus, people receiving peritoneal dialysis should be monitored carefully for drug toxicity and therapeutic drug monitoring may be necessary.

Treatment extension solely based on CKD is not recommended, as neither slower response to treatment nor higher relapse rates have been reported in these patients.

Recommendation

- **We conditionally recommend, in people with advanced chronic kidney disease (Stage 4/5, GFR < 30mL/min), that dose intervals should be modified to three times weekly for pyrazinamide (25-30mg/kg) and ethambutol (15-20mg/kg) (poor evidence).**

Good practice statement

- **Pyrazinamide and ethambutol (and other first-line TB drugs) should be dosed after dialysis session in people on hemodialysis.**

7. People with liver disease

TB treatment in patients with underlying liver cirrhosis is challenging because with limited hepatic functional reserve, they are at particular risk of liver decompensation following drug-induced hepatotoxicity. For patients with liver cirrhosis of any stage, PZA and INH are best avoided¹⁴³ and establishing a hepatic-sparing TB regimen in consultation with a TB expert is recommended.

Although rifampin is associated with drug-induced liver injury, the risk is significantly lower than with either INH or PZA.¹⁴⁴ Furthermore, rifampin is considered crucial to achieving relapse-free (or long-term) TB cure. Thus, rifampin is often used in patients with compensated liver cirrhosis (Child-Pugh A) although is usually avoided in those with overt liver decompensation (Child-Pugh B or C).¹⁴³ Fluoroquinolones, especially levofloxacin, are associated with low rates of hepatotoxicity and are sometimes used in people with decompensated liver cirrhosis.^{145,146}

Serological screening for viral hepatitis infection should be a part of routine testing at the time of TB treatment initiation for all people with TB, regardless of whether liver disease is apparent on initial testing.^{147,148} Viral hepatitis

and TB share epidemiologic associations: both hepatitis B (HBV) and C (HCV) are more prevalent in people from Asian and African regions.^{149,150} In Canada, HCV infection is prevalent in people who use drugs, those who are unstably housed and those born before 1965. Viral hepatitis is also a risk factor for drug-induced liver injury during TB treatment.¹⁵¹⁻¹⁵⁵

Observational data has shown that antiviral treatment of active hepatitis B during TB therapy reduces the incidence of subsequent drug-induced liver injury and hospitalization. TB patients found to be seropositive for hepatitis B surface antigen at the time of TB diagnosis should be promptly referred for hepatitis B treatment.¹⁵⁶

Currently recommended antiviral treatment regimens for chronic hepatitis C infection are considered incompatible with the rifamycin class because of significant drug-interactions.¹⁵⁷ Antiviral therapy for HCV is thus usually deferred until completion of rifamycin-based TB treatment. Collaboration with a hepatitis specialist is recommended.

Recommendation

- **We conditionally recommend routine serological screening for viral hepatitis at the time of TB treatment initiation (poor evidence).**

Good practice statement

- **When initiating active TB therapy in people with liver cirrhosis, consultation with a TB expert is advised. A hepatic-sparing regimen, which might exclude pyrazinamide and isoniazid, may be required.**

8. Women who are pregnant and breastfeeding

8.1. TB and pregnancy

Population studies demonstrate an elevated risk of TB in pregnant and postpartum women, with incidence rates nearly twice that of women who are not pregnant.^{2,158}

Pregnancy suppresses T-helper cell-mediated immune function, increasing susceptibility to TB infection and progressive disease.^{159,160} However, immune suppression seen during pregnancy can also mask the symptoms of progressive disease, making recognition more difficult.¹⁶¹ Furthermore, insidious symptoms of active TB may be attributed to pregnancy itself. Hesitancy to perform chest radiography may further delay diagnosis.¹⁶² After delivery, T-helper cell suppression is immediately reversed and, in some cases, the symptoms of TB disease are exacerbated as a result.¹⁵⁹

TB in pregnancy is associated with significant morbidity for both woman and their infants.^{163,164} Pregnant women with TB have higher rates of miscarriage, cesarean sections, anemia, pre-term labor and mortality.^{165,166} Infants born to mothers with active TB are more likely to be premature or low birthweight.¹⁶⁵ Thus, referral to an obstetrician for expert prenatal care is recommended for all pregnant women with TB.

Initiation of treatment for active TB in pregnancy should never be deferred, as the benefits of TB treatment greatly outweigh risks to mother and fetus. TB is not by itself an indication for termination of pregnancy. Treatment of TB in pregnant women is largely the same as in nonpregnant women.^{162,167} Treatment failure and relapse are not more common in pregnancy and treatment extension is not necessary.^{168,169} Dose adjustments are not required with advancing gestation as clinically significant changes to pharmacokinetics of the TB drugs have not been demonstrated.¹⁷⁰ Adverse effects from TB drugs, including drug-induced hepatitis, may be more common in pregnant women and careful monitoring is required.¹⁷¹

The first-line anti-TB drugs are all categorized as “Category C” by the US Food and Drug Administration. This classification reflects the lack of controlled studies in pregnant women and possible harm to fetus in animal reproductive studies. However, INH, rifampin and EMB have long track records of safety in pregnancy and are considered acceptable for first-line treatment.^{167,172,173}

Because PZA is not absolutely necessary to cure TB, and because there is a lack of formal studies on the fetal safety of this drug, its inclusion in treatment regimens for pregnant woman is usually decided on a case-by-case basis. To date, no reports of PZA teratogenicity in humans have emerged despite a long history of use²¹ and the WHO continues to recommend its use in pregnant women.¹⁷⁴ If PZA is not used, TB treatment is extended to nine months.

Fluoroquinolone use in pregnancy has not been associated with adverse pregnancy outcomes.¹⁷⁵ However, larger studies that include pregnant women exposed to longer treatment durations are needed to better establish safety; at the present time, these drugs should be used only if there are no safer alternatives.

There is considerably less experience with second-line TB agents in pregnancy¹⁶⁷ (see [Chapter 8: Drug-resistant Tuberculosis](#)). Consultation with an expert in multidrug-resistant TB is recommended.

Recommendations

- **We strongly recommend using isoniazid, rifampin and ethambutol as initial treatment in pregnant women, as all 3 are considered safe in pregnancy (good evidence).**
- **We conditionally recommend adding pyrazinamide to the regimen in pregnant women with extensive disease, smear-positive pulmonary disease, disseminated TB or intolerance of any of the other first-line drugs (poor evidence).**

8.2. TB and breastfeeding

First-line TB drugs achieve only very minimal concentrations in breast milk and toxicity to infants has not been reported. TB transmission via breast milk has not been reported in the chemotherapeutic era.¹⁷⁶

Good practice statements

- **Breastfeeding is to be encouraged in women taking first-line TB therapy.**

- **Pyridoxine supplementation for the breastfeeding infant is not necessary unless the infant is also taking isoniazid.**

See [Chapter 9: Pediatric Tuberculosis](#) for management of the neonate and peripartum periods.

9. People over 75 years of age with TB

In Canadian-born non-Indigenous and foreign-born Canadians, TB incidence rates are highest in those older than 75 years of age.¹⁷⁷ The higher rate of TB in older adults appears to be driven by a combination of higher prevalence of latent infection, increasing frequency of medical co-morbidities and, possibly, waning immunity.¹⁷⁸

TB disease can be more difficult to recognize in older adults.¹⁷⁹ Chest radiographic patterns are often atypical, with older adults less likely to demonstrate cavitation or upper-lobe predominance of infiltrates. Older adults are also more likely to have smear-negative disease.¹⁸⁰

Age over 75 years appears to be associated with poorer TB treatment outcomes, including higher mortality rates and lower completion rates.^{180,181} Age over 75 years is also associated with more adverse events, including gastrointestinal upset, rash, drug interactions and drug-induced liver injury.^{144,182–185} A systematic review demonstrated that the odds of drug-induced liver injury in active TB treatment was 30% higher in older people than in younger people.¹⁸⁵ In older patients with active TB, PZA is the most common cause of adverse events. However, these risks should be balanced with the benefit of PZA in people with high bacillary burden.

Recommendation

- **We conditionally recommend the routine use of pyrazinamide be avoided in older adults, particularly in those over 75 years of age (poor evidence).**

10. People with alcohol-use disorder

Excessive alcohol use is a well-established risk for TB. A systematic review found that people who drink more than 40g of alcohol per day or who have a diagnosed alcohol-use disorder are at a 3-fold higher risk of TB disease compared to people with lower alcohol use.^{186–188}

Alcohol likely exerts a direct toxic effect on the cellular immune system, increasing susceptibility to TB.¹⁸⁸ Furthermore, people with alcohol-use disorder experience higher rates of social marginalization, homelessness, micro- and macronutrient deficiency and incarceration, all of which are associated with elevated rates of TB infection and reactivation.¹⁸⁸

Alcohol-use disorder has been associated with delay in diagnosis of TB and a higher likelihood of sputum-smear positivity at diagnosis.^{189,190}

People with heavy alcohol use have nearly double the risk of treatment failure and have elevated TB-related mortality.^{191–193} These adverse outcomes are only partly explained by loss to follow-up; high rates of adverse events, intermittent nonadherence and severity of initial disease also contribute. In people

with alcohol-use disorder, community-based, rather than clinic-based, DOT is preferred.^{191,194}

Good practice statement

- **People with an alcohol-use disorder should receive supportive care, including community-based direct observed therapy, to ensure optimal adherence, and should be linked to alcohol counseling and support services while undergoing TB therapy.**

11. People who inject drugs

Although precise estimates are not available, people who inject drugs appear to have a higher risk of TB infection and disease than the general population.^{195–198} This susceptibility is likely mediated by several factors, including co-morbid HIV infection, tobacco use and undernutrition, as well as increased risk of exposure related to homelessness and incarceration. The presentation of TB is not evidently altered by injection drug use itself, absent co-morbid, immune-altering conditions.

Injection drug use has been associated with reduced adherence to, and lower rates of, TB treatment completion.¹⁹⁹ However, enhanced adherence supports such as provision of monetary or material treatment incentives, peer support, integration with opiate agonist therapy and DOT can improve adherence to TB treatment and monitoring.^{195,200,201} Several studies have highlighted that, with adequate adherence support, high treatment completion rates and good outcomes can be achieved in people who inject drugs.^{195,202,203}

Co-morbid liver disease is common in people who inject drugs and screening for viral hepatitis and close treatment monitoring is prudent.¹⁹⁵

Rifampin substantially reduces serum concentrations of methadone and buprenorphine, and this can precipitate opiate withdrawal syndrome.²⁰⁴ Furthermore, once rifampin is stopped, hepatic enzyme induction will wear off, usually over a period of about 2 weeks, and opiate serum concentrations can then increase, risking opiate toxicity. Rifabutin exerts less of an effect on the metabolism of these drugs and is not usually associated with withdrawal symptoms.^{205,206} Hydromorphone does not interact with the rifamycin drug class and dose adjustment is not necessary.

Given these clinically important drug-drug interactions, dosage adjustment of opioid agonist therapy and close monitoring is required throughout TB therapy. It is important that TB treatment providers alert opioid agonist providers prior to initiating rifamycin therapies and that an opiate monitoring strategy is in place. A notification to the opiate prescriber when the rifamycin is soon to stop is also strongly advised.

For patients who may not tolerate any change to opiate-agonist therapy, rifabutin can be substituted for rifampin in the TB treatment regimen.

Good practice statements

- **People with drug-use disorders should receive supportive care, including community-based directly observed**

therapy, to ensure optimal adherence, and should be linked to drug-use counseling and support services while undergoing TB therapy.

- **In patients receiving treatment for TB with rifamycin-based regimens who are also on opioid agonist therapy, it is important to adjust opioid agonist therapy in close collaboration with provider at both initiation and completion of the TB treatment course.**

12. People who smoke tobacco

People who smoke tobacco are at increased risk of TB infection and progression to active TB, likely due to biologic impacts on innate immune responses and social factors related to exposure.^{207,208} Recent systematic reviews estimate the risk of TB disease to be twice as high in people who smoke compared to nonsmokers.^{207,209}

Smoking may also affect the clinical presentation of TB. Large patient registries from Spain and Hong Kong have demonstrated that people who smoke are more likely to have pulmonary disease, lung cavitation and sputum-smear positivity and are more likely to require hospitalization for TB treatment than are nonsmokers with TB.^{210,211}

Smoking is also associated with worse TB treatment outcomes, including a higher risk of recurrence and increased mortality.^{211–213} One cohort study found those who smoke more than 10 cigarettes per day are twice as likely to relapse as those who do not currently smoke.²¹⁴ Additionally, TB is associated with an increased incidence of airway disease after treatment completion, which may compound the adverse effects of smoke exposure and existing airway disease.

TB patients are engaged into medical care for several months and have frequent visits with nursing staff, pharmacists and physicians. This represents a good opportunity to offer help in smoking cessation. A systematic review demonstrated that smoking cessation interventions appear effective in people receiving treatment for TB.²¹⁵ No controlled trials have examined the impact of smoking cessation interventions affect TB treatment outcomes.²¹⁶

Good practice statements

- **People with TB who smoke tobacco should be offered tobacco cessation interventions during TB therapy.**
- **People with pulmonary TB who smoke tobacco should be offered pulmonary function testing at the end of treatment.**

13. Drug-drug interactions

A complete medication review with assessment for potential drug interactions and need for dose adjustments of concomitant medications is recommended for all patients at initiation of TB treatment. Any adjustments of doses of these medications should be reconsidered within 2 weeks of stopping TB treatment.^{217–219}

There are several freely available online drug interaction resources that we suggest can be helpful to guide this assessment:

- Medscape (<https://reference.medscape.com/drug-interactionchecker>)
- University of Liverpool HIV Drug Interactions (<https://www.hiv-druginteractions.org/checker>)
- HIV/HCV Drug Therapy Guide from Toronto (<https://hivclinic.ca/wp-content/plugins/php/app.php>)

Rifampin is a well-documented potent inducer of hepatic and intestinal cytochrome P450 (CYP) enzymes, as well as the P-glycoprotein (P-gp) transport system. The onset of induction effects is gradual, with maximal effects on metabolizing enzymes and drug transporters by about two weeks. Induction can last for up to 4 weeks after stopping rifampin.²²⁰

Rifabutin is a less potent inducer of CYP P450 than rifampin. However, unlike rifampin, rifabutin is also a substrate of CYP3A4 and, therefore, can participate in bi-directional interactions, meaning that the metabolism of rifabutin can be altered by other drugs.^{68,221}

Rifapentine, when dosed weekly, exerts less induction of CYP P450 than does rifampin. However, when rifapentine is dosed daily, it appears to have an even greater inductive effect than standard-dosed rifampin.²²² Rifapentine is not a substrate for CYP3A.

Isoniazid is primarily metabolized via N-acetylation and is an inhibitor of several CYP450 isoenzymes. Isoniazid may

inhibit the metabolism of concomitant agents, including some antiepileptics or benzodiazepines.^{223,224}

The absorption of quinolones is significantly affected when administered at the same time as antacids and minerals containing multivalent cations, such as aluminum, magnesium or iron. Separating administration times by at least 2 hours is recommended.²¹

EMB and PZA have a low risk of drug-drug interactions.

Good practice statement

- **Drug-drug interactions and altered pharmacokinetics of TB drugs are frequently encountered in people with HIV co-infection, organ transplantation, medical co-morbidities, liver disease, renal dysfunction, and advanced age. Consultation with an experienced pharmacist is recommended when treating TB in these populations.**

Disclosure statement

The CTS TB Standards editors and authors declared potential conflicts of interest at the time of appointment and these were updated throughout the process in accordance with the CTS Conflict of Interest Disclosure Policy. Individual member conflict of interest statements are posted on the CTS website.

Table 2. Common drug-drug interactions with Rifamycins (Last updated Dec 2021).

Drugs/Drug Classes Impacted	Potential Mitigation Strategy
Anticoagulants Warfarin Apixaban, dabigatran, edoxaban, rivaroxaban	Continue warfarin with frequent INR monitoring while on rifampin and for 4 weeks after stopping rifampin. Alternatively, could change to LMWH injection. ^{225,226} Avoid use of direct-acting oral anticoagulants with rifampin. Rifabutin may be used with dabigatran. ^{227–231}
Anticonvulsants carbamazepine, phenytoin, lamotrigine	Consult with neurologist to discuss the possibility of using alternative anti-epileptic agent. Levetiracetam is a preferred anticonvulsant when taking rifamycins as there are no clinically relevant interactions. Alternatively, therapeutic drug monitoring of these anti-epileptic agents with subsequent dose adjustment is required. ^{232–235}
Antidiabetics agents	More frequent blood glucose monitoring recommended. Potential for decrease in drug concentration of most oral antidiabetic agents in patients on rifamycins and potential for increased blood glucose levels. Dose adjustment of antidiabetic agents may be required. ^{236–243} No significant interaction with metformin or insulin. ²⁴³
Antifungals Azoles (itraconazole, fluconazole, posaconazole, voriconazole, isavuconazole)	Subtherapeutic azole concentrations may occur when used with any of the rifamycins. ^{244–248}
Antihypertensives	Increased BP monitoring recommended with most antihypertensives. Rifampin has a strong interaction with calcium channel blockers and clinicians may be required to increase dose of the calcium channel blocker or change/add an alternate antihypertensive agent (consult provider). Rifabutin has a less potent interaction as compared to rifampin. ^{249–252}
Antiretroviral agents	See Text and Table 1.
Corticosteroids dexamethasone, methylprednisolone, prednisone	Monitor clinical response. May require increase in dose of corticosteroids. ^{253–255}
Immunosuppressive agents cyclosporine, tacrolimus	Rifabutin use is preferred over rifampin to minimize impact on concentrations of calcineurin inhibitors and to reduce the risk of allograft rejection. Monitor calcineurin inhibitor concentrations and adjust dose as required. ^{256–260}
Levothyroxine	Continue both agents but monitor TSH monthly. May require increase in dose of levothyroxine with rifamycins. Monitor TSH after stopping rifamycins. ^{261–263}
Opiate agonist therapy methadone, buprenorphine/naloxone (Suboxone)	Methadone and buprenorphine serum concentrations decrease substantially with rifampin and precipitation of withdrawal symptoms is frequent. May need preemptive OAT regimen dosage adjustment in addition to close monitoring. Discuss OAT with provider. ^{204,264,265} Rifabutin has much less impact on methadone and buprenorphine serum concentrations and is not associated with withdrawal symptoms. ^{205,206}
Oral hormonal contraceptive ethinyl estradiol, norethindrone, etc	Add a barrier method of contraception when taking a rifamycin with oral hormonal contraceptives. ²⁶⁶

Abbreviations: INR, international normalized ratio; LMWH, Low molecular weight heparin; BP, blood pressure; TSH, thyroid-stimulating hormone; OAT, opiate agonist therapy.

Funding

The 8th edition Canadian Tuberculosis Standards are jointly funded by the Canadian Thoracic Society (CTS) and the Public Health Agency of Canada, edited by the CTS and published by the CTS in collaboration with AMMI Canada. However, it is important to note that the clinical recommendations in the Standards are those of the CTS. The CTS TB Standards editors and authors are accountable to the CTS CRGC and the CTS Board of Directors. The CTS TB Standards editors and authors are functionally and editorially independent from any funding sources and did not receive any direct funding from external sources.

The CTS receives unrestricted grants which are combined into a central operating account to facilitate the knowledge translation activities of the CTS Assemblies and its guideline and standards panels. No corporate funders played any role in the collection, review, analysis or interpretation of the scientific literature or in any decisions regarding the recommendations presented in this document.

References

1. Creswell J, Raviglione M, Ottmani S, et al. Tuberculosis and noncommunicable diseases: neglected links and missed opportunities. *Eur Respir J*. 2011;37(5):1269–1282. doi:10.1183/09031936.00084310.
2. Mathad JS, Gupta A. Tuberculosis in Pregnant and Postpartum Women: Epidemiology, Management, and Research Gaps. *Clin Infect Dis*. 2012;55(11):1532–49. doi:10.1093/cid/cis732.
3. Perumal R, Naidoo K, Naidoo A, et al. A systematic review and meta-analysis of first-line tuberculosis drug concentrations and treatment outcomes. *Int J Tuberc Lung Dis*. 2020;24(1):48–64. doi:10.5588/ijtld.19.0025.
4. Wood R, Maartens G, Lombard CJ. Risk Factors for Developing Tuberculosis in HIV-1-infected adults from communities with a low or very high incidence of tuberculosis. *J Acquir Immune Defic Syndr*. 2000;23(1):75–80. doi:10.1097/00126334-200001010-00010.
5. Selwyn PA, Hartel D, Lewis VA, et al. A Prospective Study of the Risk of Tuberculosis among Intravenous Drug Users with Human Immunodeficiency Virus Infection. *N Engl J Med*. 1989;320(9):545–550. doi:10.1056/nejm198903023200901.
6. Pape JW, Jean SS, Ho JL, Hafner A, Johnson WD. Effect of isoniazid prophylaxis on incidence of active tuberculosis and progression of HIV infection. *The Lancet*. 1993;342(8866):268–272. doi:10.1016/0140-6736(93)91817-6.
7. Suthar AB, Lawn SD, del Amo J, et al. Antiretroviral therapy for prevention of tuberculosis in adults with HIV: a systematic review and meta-analysis. *PLoS Med*. 2012;9(7):e1001270. doi:10.1371/journal.pmed.1001270.
8. Gupta A, Wood R, Kaplan R, Bekker LG, Lawn SD. Tuberculosis incidence rates during 8 years of follow-up of an antiretroviral treatment cohort in south africa: Comparison with rates in the community. *PLoS One*. 2012;7(3):e34156–8. doi:10.1371/journal.pone.0034156.
9. Houben RMGJ, Crampin AC, Ndhlovu R, et al. Human immunodeficiency virus associated tuberculosis more often due to recent infection than reactivation of latent infection. *International Journal of Tuberculosis and Lung Disease*. 2011;15(1):24–31. doi:10.5588/ijtld.11.0115.
10. Oni T, Burke R, Tsekela R, et al. High prevalence of subclinical tuberculosis in HIV-1-infected persons without advanced immunodeficiency: Implications for TB screening. *Thorax*. 2011;66(8):669–673. doi:10.1136/thx.2011.160168.
11. Aderaye G, Bruchfeld J, Assefa G, et al. The relationship between disease pattern and disease burden by chest radiography, M. tuberculosis load, and HIV status in patients with pulmonary tuberculosis in Addis Ababa. *Infection*. 2004;32(6):333–338. doi:10.1007/s15010-004-3089-x.
12. Li J, Munsiff SS, Driver CR, Sackoff J. Relapse and acquired rifampin resistance in HIV-infected patients with tuberculosis treated with rifampin- or rifabutin-based regimens in New York City, 1997–2000. *Clinical Infectious Diseases*. 2005;41(1):83–91. doi:10.1086/430377.
13. Nettles RE, Mazo D, Alwood K, et al. Risk factors for relapse and acquired rifampin resistance after directly observed tuberculosis treatment: A comparison by HIV serostatus and rifampin use. *Clin Infect Dis*. 2004;38(5):731–736. doi:10.1086/381675.
14. Comstock GW, Golub JE, Panjabi R. Recurrent tuberculosis and its risk factors: Adequately treated patients are still at high risk. *International Journal of Tuberculosis and Lung Disease*. 2007;11(8):828–837.
15. Sonnenberg P, Murray J, Glynn JR, Shearer S, Kambashi B, Godfrey-Faussett P. HIV-1 and recurrence, relapse, and reinfection of tuberculosis after cure: A cohort study in South African mineworkers. *Lancet*. 2001;358(9294):1687–1693. doi:10.1016/S0140-6736(01)06712-5.
16. Ahmad Khan F, Minion J, Al-Motairi A, Benedetti A, Harries AD, Menzies D. An updated systematic review and meta-analysis on the treatment of active tuberculosis in patients with hiv infection. *Clin Infect Dis*. 2012;55(8):1154–1163. doi:10.1093/cid/cis630.
17. Haar CH, Cobelens FGJ, Kalisvaart NA, Van Gerven PJHJ, Van Der Have JJ. HIV-related mortality among tuberculosis patients in the Netherlands, 1993–2001. *International Journal of Tuberculosis and Lung Disease*. 2007;11(9):1038–1041.
18. El-Sadr WM, Perlman DC, Matts JP, et al. Evaluation of an intensive intermittent-induction regimen and duration of short-course treatment for human immunodeficiency virus-related pulmonary tuberculosis. Terry Beinr Community Programs for Clinical Research on AIDS (CPCRA) and the AIDS Clinical Trials Group (ACTG). *Clin Infect Dis*. 1998;26(5):1148–1158. doi:10.1086/520275.
19. Perriens JH, St. Louis ME, Mukadi YB, et al. Pulmonary Tuberculosis in HIV-Infected Patients in Zaire — A Controlled Trial of Treatment for Either 6 or 12 Months. *N Engl J Med*. 1995;332(12):779–785. doi:10.1056/NEJM199503233321204.
20. Davies G, Cerri S, Richeldi L. Rifabutin for treating pulmonary tuberculosis. *The Cochrane Database of Systematic Reviews*. 2005; (1):22–24. doi:10.1002/14651858.cd005159.
21. Nahid P, Dorman SE, Alipanah N, et al. Official American Thoracic Society/Centers for Disease Control and Prevention/ Infectious Diseases Society of America Clinical Practice Guidelines: Treatment of Drug-Susceptible Tuberculosis. *Clin Infect Dis*. 2016;63(7):e147–e195. doi:10.1093/cid/ciw376.
22. Bracchi M, van Halsema C, Post F, et al. British HIV Association guidelines for the management of tuberculosis in adults living with HIV 2019. *HIV Med*. 2019;20:s2–s83. doi:10.1111/hiv.12748.
23. Panel on Guidelines for the Prevention and Treatment of Opportunistic Infections in Adults and Adolescents with HIV. Guidelines for the Prevention and Treatment of Opportunistic Infections in HIV-infected Adults and Adolescents: Recommendations from the Centers for Disease Control and Prevention, the National Institutes of Health, and the HIV Medicine Association of the Infectious Diseases Society of America. 2021. <https://clinicalinfo.hiv.gov/en/guidelines/adult-and-adolescent-opportunistic-infection/whats-new-guidelines>. Accessed November 1, 2021.
24. Mfinanga SG, Kirenga BJ, Chanda DM, et al. Early versus delayed initiation of highly active antiretroviral therapy for HIV-positive adults with newly diagnosed pulmonary tuberculosis (TB-HAART): A prospective, international, randomised, placebo-controlled trial. *Lancet Infect Dis*. 2014;14(7):563–571. doi:10.1016/S1473-3099(14)70733-9.
25. Gopalan N, Santhanakrishnan RK, Palaniappan AN, et al. Daily vs intermittent antituberculosis therapy for pulmonary tuberculosis in patients with HIV: A Randomized Clinical Trial. *JAMA*

- Intern Med.* 2018;178(4):485–493. doi:10.1001/jamainternmed.2018.0141.
26. Tappero JW, Bradford WZ, Agerton TB, et al. Serum concentrations of antimycobacterial drugs in patients with pulmonary tuberculosis in Botswana. *Clinical Infectious Diseases.* 2005;41(4):461–469. doi:10.1086/431984.
 27. Chideya S, Winston CA, Peloquin CA, et al. Isoniazid, rifampin, ethambutol, and pyrazinamide pharmacokinetics and treatment outcomes among a predominantly HIV-infected cohort of adults with tuberculosis from Botswana. *Clin Infect Dis.* 2009;48(12):1685–1694. doi:10.1086/599040.
 28. Alsultan A, Peloquin CA. Therapeutic drug monitoring in the treatment of tuberculosis: An update. *Drugs.* 2014;74(8):839–854. doi:10.1007/s40265-014-0222-8.
 29. Babalik A, Babalik A, Mannix S, Francis D, Menzies D. Therapeutic drug monitoring in the treatment of active tuberculosis. *Can Respir J.* 2011;18(4):225–229. doi:10.1155/2011/307150.
 30. Daskapan A, Idrus LR, Postma MJ, et al. A Systematic Review on the Effect of HIV Infection on the Pharmacokinetics of First-Line Tuberculosis Drugs. *Clin Pharmacokinet.* 2019;58(6):747–766. doi:10.1007/s40262-018-0716-8.
 31. Kwara A, Enimil A, Gillani FS, et al. Pharmacokinetics of first-line antituberculosis drugs using WHO revised dosage in children with tuberculosis with and without HIV coinfection. *J Pediatric Infect Dis Soc.* 2016;5(4):356–365. doi:10.1093/jpids/piv035.
 32. Mårtson AG, Burch G, Ghimire S, Alffenaar JWC, Peloquin CA. Therapeutic drug monitoring in patients with tuberculosis and concurrent medical problems. *Expert Opin Drug Metab Toxicol.* 2021;17(1):23–39. doi:10.1080/17425255.2021.1836158.
 33. Sileshi T, Tadesse E, Makonnen E, Aklilu E. The impact of first-line anti-tubercular Drugs' Pharmacokinetics on Treatment Outcome: A Systematic Review. *Clin Pharmacol.* 2021;13:1–12. doi:10.2147/CPAA.S289714.
 34. Pasipanodya JG, McIlleron H, Burger A, Wash PA, Smith P, Gumbo T. Serum drug concentrations predictive of pulmonary tuberculosis outcomes. *J Infect Dis.* 2013;208(9):1464–1473. doi:10.1093/infdis/jit352.
 35. Weiner M, Benator D, Burman W, Tuberculosis Trials Consortium, et al. Association between acquired rifamycin resistance and the pharmacokinetics of rifabutin and isoniazid among patients with HIV and tuberculosis. *Clin Infect Dis.* 2005;40(10):1481–1491. doi:10.1086/429321.
 36. Uthman OA, Okwundu C, Gbenga K, et al. Optimal timing of antiretroviral therapy initiation for HIV-infected adults with newly diagnosed pulmonary tuberculosis: A systematic review and meta-analysis. *Ann Intern Med.* 2015;163(1):32–39. doi:10.7326/M14-2979.
 37. Burke RM, Rickman HM, Singh V, et al. What is the optimum time to start antiretroviral therapy in people with HIV and tuberculosis coinfection? A systematic review and meta-analysis. *J Int AIDS Soc.* 2021;24(7):e25772. doi:10.1002/jia2.25772.
 38. World Health Organization. *Consolidated Guidelines on HIV Prevention, Testing, Treatment, Service Delivery and Monitoring: recommendations for a Public Health Approach.* 2021. Geneva, Switzerland. <https://www.who.int/publications/i/item/9789240031593>. Accessed November 1, 2021.
 39. Török ME, Yen NTB, Chau TTH, et al. Timing of initiation of antiretroviral therapy in human immunodeficiency virus (HIV)-associated tuberculous meningitis. *Clin Infect Dis.* 2011;52(11):1374–1383. doi:10.1093/cid/cir230.
 40. Cerrone M, Alfarisi O, Neary M, et al. Rifampicin effect on intracellular and plasma pharmacokinetics of tenofovir alafenamide. *J Antimicrob Chemother.* 2019;74(6):1670–1678. doi:10.1093/jac/dkz068.
 41. De Castro N, Marcy O, Chazallon C, et al. Standard dose raltegravir or efavirenz-based antiretroviral treatment for patients co-infected with HIV and tuberculosis (ANRS 12 300 Replate TB 2): an open-label, non-inferiority, randomised, phase 3 trial. *The Lancet Infectious Diseases.* 2021;21(6):813–822. doi:10.1016/S1473-3099(20)30869-0.
 42. Wenning LA, Hanley WD, Brainard DM, et al. Effect of rifampin, a potent inducer of drug-metabolizing enzymes, on the pharmacokinetics of raltegravir. *Antimicrob Agents Chemother.* 2009;53(7):2852–2856. doi:10.1128/AAC.01468-08.
 43. Dooley KE, Kaplan R, Mwelase N, International Study of Patients with HIV on Rifampicin ING study group, et al. Dolutegravir-based Antiretroviral Therapy for Patients Coinfected with Tuberculosis and Human Immunodeficiency Virus: A Multicenter, Noncomparative, Open-label, Randomized Trial. *Clinical Infectious Diseases.* 2019;70(4):549–556. doi:10.1093/cid/ciz256.
 44. Wang X, Cerrone M, Ferretti F, et al. Pharmacokinetics of dolutegravir 100 mg once daily with rifampicin. *Int J Antimicrob Agents.* 2019;54(2):202–206. doi:10.1016/j.ijantimicag.2019.04.009.
 45. Dooley KE, Sayre P, Borland J, et al. Safety, tolerability, and pharmacokinetics of the HIV integrase inhibitor dolutegravir given twice daily with rifampin or once daily with rifabutin: Results of a phase 1 study among healthy subjects. *J Acquir Immune Defic Syndr.* 2013;62(1):21–27. doi:10.1097/QAI.0b013e318276cda9.
 46. Manosuthi W, Kiertiburanakul S, Sungkanuparph S, et al. Efavirenz 600 mg/day versus efavirenz 800 mg/day in HIV-infected patients with tuberculosis receiving rifampicin: 48 Weeks results. *AIDS.* 2006;20(1):131–132. doi:10.1097/01.aids.0000196181.18916.9b.
 47. Meintjes G, Brust JCM, Nuttall J, Maartens G. Management of active tuberculosis in adults with HIV. *Lancet HIV.* 2019;6(7):e463–e474. doi:10.1016/S2352-3018(19)30154-7.
 48. Cohen K, Meintjes G. Management of individuals requiring antiretroviral therapy and TB treatment. *Curr Opin HIV Aids.* 2010;5(1):61–69. doi:10.1097/COH.0b013e3283339309.
 49. Brainard DM, Kassahun K, Wenning LA, et al. Lack of a clinically meaningful pharmacokinetic effect of rifabutin on raltegravir: In vitro/in vivo correlation. *J Clin Pharmacol.* 2011;51(6):943–950. doi:10.1177/0091270010375959.
 50. Meintjes G, Lawn SD, Scano F, International Network for the Study of HIV-associated IRIS, et al. Tuberculosis-associated immune reconstitution inflammatory syndrome: case definitions for use in resource-limited settings. *Lancet Infect Dis.* 2008;8(8):516–523. doi:10.1016/S1473-3099(08)70184-1.
 51. Marais S, Meintjes G, Pepper DJ, et al. Frequency, severity, and prediction of tuberculous meningitis immune reconstitution inflammatory syndrome. *Clin Infect Dis.* 2013;56(3):450–460. doi:10.1093/cid/cis899.
 52. Manabe YC, Breen R, Perti T, Girardi E, Sterling TR. Unmasked tuberculosis and tuberculosis immune reconstitution inflammatory disease: A disease spectrum after initiation of antiretroviral therapy. *J Infect Dis.* 2009;199(3):437–444. doi:10.1086/595985.
 53. Meintjes G, Rangaka MX, Maartens G, et al. Novel relationship between tuberculosis immune reconstitution inflammatory syndrome and antitubercular drug resistance. *Clin Infect Dis.* 2009;48(5):667–676. doi:10.1086/596764.
 54. Meintjes G, Wilkinson RJ, Morroni C, et al. Randomized placebo-controlled trial of prednisone for paradoxical tuberculosis-associated immune reconstitution inflammatory syndrome. *AIDS.* 2010;24(15):2381–2390. doi:10.1097/QAD.0b013e32833dfc68.
 55. Meintjes G, Stek C, Blumenthal L, PredART Trial Team, et al. Prednisone for the Prevention of Paradoxical Tuberculosis-Associated IRIS. *N Engl J Med.* 2018;379(20):1915–1925. doi:10.1056/nejmoa1800762.
 56. Subramanian AK. Mycobacterium tuberculosis in Transplantation. *Emerging Transplant Infections: Clinical Challenges and Implications.* 2020;1–16. doi:10.1007/978-3-030-01751-4_15-1.
 57. Torre-Cisneros J, Doblas A, Aguado JM, Spanish Network for Research in Infectious Diseases, et al. Tuberculosis after solid-organ transplant: incidence, risk factors, and clinical characteristics in the resitra (spanish network of infection in

- transplantation) cohort. *Clin Infect Dis*. 2009;48(12):1657–1665. doi:10.1086/599035.
58. Benito N, García-Vázquez E, Horcajada JP, et al. Clinical features and outcomes of tuberculosis in transplant recipients as compared with the general population: A retrospective matched cohort study. *Clinical Microbiology and Infection*. 2015;21(7):651–658. doi:10.1016/j.cmi.2015.03.010.
 59. Singh N, Paterson DL. Mycobacterium tuberculosis infection in solid-organ transplant recipients: Impact and implications for management. *Clin Infect Dis*. 1998;27(5):1266–1277. doi:10.1086/514993.
 60. Subramanian AK, Theodoropoulos NM. Mycobacterium tuberculosis infections in solid organ transplantation: Guidelines from the infectious diseases community of practice of the American Society of Transplantation. *Clin Transplant*. 2019;33(9):e13513. doi:10.1111/ctr.13513.
 61. Giacomelli IL, Neto RS, Marchiori E, Pereira M, Hochhegger B. Chest x-ray and chest ct findings in patients diagnosed with pulmonary tuberculosis following solid organ transplantation: A systematic review. *J Bras Pneumol*. 2018;44(2):161–166. doi:10.1590/s1806-37562017000000459.
 62. Muñoz P, Rodríguez C, Bouza E. Mycobacterium tuberculosis infection in recipients of solid organ transplants. *Clin Infect Dis*. 2005;40(4):581–587. doi:10.1086/427692.
 63. Abad CLR, Razonable RR. Mycobacterium tuberculosis after solid organ transplantation: A review of more than 2000 cases. *Clin Transplant*. 2018;32(6):e13259. doi:10.1111/ctr.13259.
 64. Sun HY, Munoz P, Torre-Cisneros J, et al. Tuberculosis in solid-organ transplant recipients: Disease characteristics and outcomes in the current era. *Prog Transplant*. 2014;24(1):37–43. doi:10.7182/pit2014398.
 65. Aguado J, Torre-Cisneros J, Fortun J, et al. Tuberculosis in solid-organ transplant recipients: Consensus statement of the Group for the Study of Infection in transplant recipients (GESITRA) of the Spanish society of infectious diseases and clinical microbiology. *Clin Infect Dis*. 2009;48(9):1276–1284. doi:10.1086/597590.
 66. Meije Y, Piersimoni C, Torre-Cisneros J, Dilektasli AG, Aguado JM. Mycobacterial infections in solid organ transplant recipients. *Clinical Microbiology and Infection*. 2014;20:89–101. doi:10.1111/1469-0691.12641.
 67. Pennington KM, Kennedy CC, Chandra S, et al. Management and diagnosis of tuberculosis in solid organ transplant candidates and recipients: Expert survey and updated review. *J Clin Tuberc Other Mycobact Dis*. 2018;11:37–46. doi:10.1016/j.jc-tube.2018.04.001.
 68. Baciewicz AM, Chrisman CR, Finch CK, Self TH. Update on rifampin, rifabutin, and rifapentine drug interactions. *Curr Med Res Opin*. 2013;29(1):1–12. doi:10.1185/03007995.2012.747952.
 69. Hickey MD, Quan DJ, Chin-Hong PV, Roberts JP. Use of rifabutin for the treatment of a latent tuberculosis infection in a patient after solid organ transplantation. *Liver Transpl*. 2013;19(4):457–461. doi:10.1002/lt.23622.
 70. López-Montes A, Gallego E, López E, et al. Treatment of tuberculosis with rifabutin in a renal transplant recipient. *American Journal of Kidney Diseases*. 2004;44(4):e59–63. doi:10.1053/j.ajkd.2004.06.015.
 71. Grassi C, Peona V. Use of rifabutin in the treatment of pulmonary tuberculosis. *Clinical Infectious Diseases*. 1996;22(Supplement_1):S50–S54. doi:10.1093/clinids/22.supplement_1.s50.
 72. McGregor MM, Oliario P, Wolmarans L, et al. Efficacy and safety of rifabutin in the treatment of patients with newly diagnosed pulmonary tuberculosis. *Am J Respir Crit Care Med*. 1996;154(5):1462–1467. doi:10.1164/ajrccm.154.5.8912765.
 73. Crabol Y, Catherinot E, Veziris N, Jullien V, Lortholary O. Rifabutin: Where do we stand in 2016? *J Antimicrob Chemother*. 2016;71(7):1759–1771. doi:10.1093/jac/dkw024.
 74. Sun HY, Munoz P, Torre-Cisneros J, et al. Mycobacterium tuberculosis-associated immune reconstitution syndrome in solid-organ transplant recipients. *Transplantation*. 2013;95(9):1173–1181. doi:10.1097/TP.0b013e31828719c8.
 75. Park YS, Choi JY, Cho CH, et al. Clinical outcomes of tuberculosis in renal transplant recipients. *Yonsei Med J*. 2004;45(5):865–872. doi:10.3349/ymj.2004.45.5.865.
 76. Zhang Z, Fan W, Yang G, et al. Risk of tuberculosis in patients treated with TNF- α antagonists: A systematic review and meta-analysis of randomised controlled trials. *BMJ Open*. 2017;7(3):e012567–8. doi:10.1136/bmjopen-2016-012567.
 77. Sartori NS, de Andrade NPB, da Silva Chakr RM. Incidence of tuberculosis in patients receiving anti-TNF therapy for rheumatic diseases: a systematic review. *Clin Rheumatol*. 2020;39(5):1439–1447. doi:10.1007/s10067-019-04866-x.
 78. Dobler CC. Biological agents and tuberculosis: Risk estimates and screening strategies. *Int J Rheum Dis*. 2015;18(3):264–267. doi:10.1111/1756-185X.12672.
 79. Carmona L, Gómez-Reino JJ, Rodríguez-Valverde V, BIOBADASER Group, et al. Effectiveness of recommendations to prevent reactivation of latent tuberculosis infection in patients treated with tumor necrosis factor antagonists. *Arthritis Rheum*. 2005;52(6):1766–1772. doi:10.1002/art.21043.
 80. Lorenzetti R, Zullo A, Ridola L, et al. Higher risk of tuberculosis reactivation when anti-TNF is combined with immunosuppressive agents: A systematic review of randomized controlled trials. *Ann Med*. 2014;46(7):547–554. doi:10.3109/07853890.2014.941919.
 81. Dixon WG, Hyrich KL, Watson KD, BSR Biologics Register, et al. Drug-specific risk of tuberculosis in patients with rheumatoid arthritis treated with anti-TNF therapy: Results from the British Society for Rheumatology Biologics Register (BSRBR). *Ann Rheum Dis*. 2010;69(3):522–528. doi:10.1136/ard.2009.118935.
 82. Singh JA, Wells GA, Christensen R, Cochrane Musculoskeletal Group, et al. Adverse effects of biologics: a network meta-analysis and Cochrane overview. *The Cochrane Database of Systematic Reviews*. 2011;81(2):CD008794. doi:10.1002/14651858.CD008794.pub2.
 83. Gómez-Reino JJ, Carmona L, Rodríguez Valverde V, Mola EM, Montero MD, BIOBADASER Group. Treatment of rheumatoid arthritis with tumor necrosis factor inhibitors may predispose to significant increase in tuberculosis risk: A multicenter active-surveillance report. *Arthritis Rheum*. 2003;48(8):2122–2127. doi:10.1002/art.11137.
 84. Ai JW, Zhang S, Ruan QL, et al. The risk of tuberculosis in patients with rheumatoid arthritis treated with tumor necrosis factor- α antagonist: A metaanalysis of both randomized controlled trials and registry/cohort studies. *J Rheumatol*. 2015;42(12):2229–2237. doi:10.3899/jrheum.150057.
 85. Solovic I, Sester M, Gomez-Reino JJ, Series "update on tuberculosis" edited by C. Lange, M. Raviglione, W.W. Yew and G.B, et al. Migliori number 2 in this series: The risk of tuberculosis related to tumour necrosis factor antagonist therapies: A TBNET consensus statement. *Eur Respir J*. 2010;36(5):1185–1206. doi:10.1183/09031936.00028510.
 86. Arend SM, Leyten EMS, Franken WPJ, Huisman EM, Van Dissel JT. A patient with de novo tuberculosis during anti-tumor necrosis factor-alpha therapy illustrating diagnostic pitfalls and paradoxical response to treatment. *Clin Infect Dis*. 2007;45(11):1470–1475. doi:10.1086/522993.
 87. Wallis RS, Broder MS, Wong JY, Hanson ME, Beenhouwer DO. Granulomatous infectious diseases associated with tumor necrosis factor antagonists. *Clinical Infectious Diseases*. 2004;38(9):1261–1265. doi:10.1086/383317.
 88. Mohan AK, Coté TR, Block JA, Manadan AM, Siegel JN, Braun MM. Tuberculosis following the use of etanercept, a tumor necrosis factor inhibitor. *Clin Infect Dis*. 2004;39(3):295–299. doi:10.1086/421494.
 89. Godfrey MS, Friedman LN. Tuberculosis and Biologic Therapies: Anti-Tumor Necrosis Factor- α and Beyond. *Clin Chest Med*. 2019;40(4):721–739. doi:10.1016/j.ccm.2019.07.003.

90. Centers for Disease Control and Prevention. Tuberculosis associated with blocking agents against tumor necrosis factor- α - California, 2002-2003. *MMWR Recommendations and Reports: Morbidity and Mortality Weekly Report Recommendations and Reports/Centers for Disease Control*. 2004;53(30):683-685. doi:10.1001/jama.292.14.1676.
91. Cantini F, Prignano F, Goletti D. Restarting biologics and management of patients with flares of inflammatory rheumatic disorders or psoriasis during active tuberculosis treatment. *J Rheumatol Suppl*. 2014;91(0):78-82. doi:10.3899/jrheum.140106.
92. Wallis RS, van Vuuren C, Potgieter S. Adalimumab Treatment of life-threatening tuberculosis. *Clin Infect Dis*. 2009;48(10):1429-1432. doi:10.1086/598504.
93. Matsumoto T, Tanaka T, Kawase I. Infliximab for Rheumatoid Arthritis in a Patient with Tuberculosis. *N Engl J Med*. 2006;355(7):740-741. doi:10.1056/nejmc053468.
94. Garcia Vidal C, Rodríguez Fernández S, Martínez Lacasa J, et al. Paradoxical response to antituberculous therapy in infliximab-treated patients with disseminated tuberculosis. *Clin Infect Dis*. 2005;40(5):756-759. doi:10.1086/427941.
95. Dobler CC. Biologic Agents and Tuberculosis. *Microbiol Spectr*. 2016;4(6). doi:10.1128/microbiolspec.TNMI7-0026-2016.
96. Hsu DC, Faldetta KF, Pei L, et al. A Paradoxical Treatment for a Paradoxical Condition: Infliximab Use in Three Cases of Mycobacterial IRIS. *Clin Infect Dis*. 2016;62(2):258-261. doi:10.1093/cid/civ841.
97. Cho SK, Kim D, Won S, et al. Safety of resuming biologic DMARDs in patients who develop tuberculosis after anti-TNF treatment. *Semin Arthritis Rheum*. 2017;47(1):102-107. doi:10.1016/j.semarthrit.2017.01.004.
98. Kim HW, Kwon SR, Jung KH, Korean Society of Spondyloarthritis Research, et al. Safety of resuming tumor necrosis factor inhibitors in ankylosing spondylitis patients concomitant with the treatment of active tuberculosis: A retrospective nationwide registry of the Korean Society of spondyloarthritis research. *PLoS One*. 2016;11(4):e0153816-14. doi:10.1371/journal.pone.0153816.
99. Kim YJ, Kim YG, Shim TS, et al. Safety of resuming tumour necrosis factor inhibitors in patients who developed tuberculosis as a complication of previous TNF inhibitors. *Rheumatology (Oxford)*. 2014;53(8):1477-1481. doi:10.1093/rheumatology/keu041.
100. Ozguler Y, Hatemi G, Ugurlu S, et al. Re-initiation of biologics after the development of tuberculosis under anti-TNF therapy. *Rheumatol Int*. 2016;36(12):1719-1725. doi:10.1007/s00296-016-3575-3.
101. Jeon CY, Murray MB. Diabetes mellitus increases the risk of active tuberculosis: A systematic review of 13 observational studies. *PLoS Med*. 2008;5(7):e152-1101. doi:10.1371/journal.pmed.0050152.
102. Al-Rifai RH, Pearson F, Critchley JA, Abu-Raddad LJ. Association between diabetes mellitus and active tuberculosis: A systematic review and meta-analysis. *PLoS One*. 2017;12(11):e0187967-26. doi:10.1371/journal.pone.0187967.
103. Dooley KE, Chaisson RE. Tuberculosis and diabetes mellitus: convergence of two epidemics. *Lancet Infect Dis*. 2009;9(12):737-746. doi:10.1016/S1473-3099(09)70282-8.
104. Viney K, Mills T, Harley D. Tuberculosis and diabetes mellitus: A dose-response relationship between the odds of tuberculosis and HbA1c. *Int J Tuberc Lung Dis*. 2019;23(10):1055-1059. doi:10.5588/ijtld.18.0657.
105. Golub JE, Mok Y, Hong S, Jung KJ, Jee SH, Samet JM. Diabetes mellitus and tuberculosis in Korean adults: Impact on tuberculosis incidence, recurrence and mortality. *Int J Tuberc Lung Dis*. 2019;23(4):507-513. doi:10.5588/ijtld.18.0103.
106. Leung CC, Yew WW, Mok TYW, et al. Effects of diabetes mellitus on the clinical presentation and treatment response in tuberculosis. *Respirology*. 2017;22(6):1225-1232. doi:10.1111/resp.13017.
107. Wang CS, Yang CJ, Chen HC, et al. Impact of type 2 diabetes on manifestations and treatment outcome of pulmonary tuberculosis. *Epidemiol Infect*. 2009;137(2):203-210. doi:10.1017/S0950268808000782.
108. Morris JT, Seaworth BJ, McAllister CK. Pulmonary tuberculosis in diabetics. *Chest*. 1992;102(2):539-541. doi:10.1378/chest.102.2.539.
109. Park SW, Shin JW, Kim JY, et al. The effect of diabetic control status on the clinical features of pulmonary tuberculosis. *Eur J Clin Microbiol Infect Dis*. 2012;31(7):1305-1310. doi:10.1007/s10096-011-1443-3.
110. Baker MA, Harries AD, Jeon CY, et al. The impact of diabetes on tuberculosis treatment outcomes: A systematic review. *BMC Med*. 2011;9(1):81. doi:10.1186/1741-7015-9-81.
111. Degner NR, Wang JY, Golub JE, Karakousis PC. Metformin Use Reverses the Increased Mortality Associated with Diabetes Mellitus during Tuberculosis Treatment. *Clin Infect Dis*. 2018;66(2):198-205. doi:10.1093/cid/cix819.
112. Huangfu P, Pearson F, Ugarte-Gil C, Critchley J. Diabetes and poor tuberculosis treatment outcomes: Issues and implications in data interpretation and analysis. *Int J Tuberc Lung Dis*. 2017;21(12):1214-1219. doi:10.5588/ijtld.17.0211.
113. Martinez L, Zhu L, Castellanos ME, et al. Glycemic Control and the Prevalence of Tuberculosis Infection: A Population-based Observational Study. *Clin Infect Dis*. 2017;65(12):2060-2068. doi:10.1093/cid/cix632.
114. Mave V, Gaikwad S, Barthwal M, et al. Diabetes Mellitus and Tuberculosis Treatment Outcomes in Pune, India. *Open Forum Infect Dis*. 2021;8(4):ofab097. doi:10.1093/ofid/ofab097.
115. Armstrong LR, Steve Kammerer J, Haddad MB. Diabetes mellitus among adults with tuberculosis in the USA, 2010-2017. *BMJ Open Diabetes Research and Care*. 2020;8(1):1-6. doi:10.1136/bmjdr-2020-001275.
116. Ugarte-Gil C, Alisjahbana B, Riza AL. Diabetes mellitus among pulmonary tuberculosis patients from four TB - endemic countries: the TANDEM study. *Clin Infect Dis*. 2020;70(5):780-788. doi:10.1093/cid/ciz284.
117. White LV, Edwards T, Lee N, et al. Patterns and predictors of co-morbidities in Tuberculosis: A cross-sectional study in the Philippines. *Sci Rep*. 2020;10(1):1-12. doi:10.1038/s41598-020-60942-2.
118. Workneh MH, Bjune GA, Yimer SA. Prevalence and associated factors of tuberculosis and diabetes mellitus comorbidity: A systematic review. *PLoS One*. 2017;12(4):e0175925-25. doi:10.1371/journal.pone.0175925.
119. Segafredo G, Kapur A, Robbiati C, et al. Integrating TB and non-communicable diseases services: Pilot experience of screening for diabetes and hypertension in patients with Tuberculosis in Luanda, Angola. *PLoS One*. 2019;14(7):e0218052-11. doi:10.1371/journal.pone.0218052.
120. Barss L, Sabur N, Orlikow E, et al. Diabetes Mellitus in adult patients with active tuberculosis in Canada: Presentation and treatment outcomes. *Canadian Journal of Respiratory, Critical Care, and Sleep Medicine*. 2019;3(2):84-90. doi:10.1080/24745332.2018.1507615.
121. Grint D, Alisjahbana B, Ugarte-Gil C, TANDEM consortium, et al. Accuracy of diabetes screening methods used for people with tuberculosis, Indonesia, Peru, Romania, South Africa. *Bull World Health Organ*. 2018;96(11):738-749. doi:10.2471/BLT.17.206227.
122. Lin Y, Harries AD, Kumar AMV, et al. Management of Diabetes Mellitus-Tuberculosis. *International Union against Tuberculosis and Lung Disease*. 2019. https://theunion.org/sites/default/files/2020-11/TheUnion_DMTB_Guide.pdf. Accessed November 1, 2021.
123. Huangfu P, Ugarte-Gil C, Golub J, Pearson F, Critchley J. The effects of diabetes on tuberculosis treatment outcomes: an updated systematic review and meta-analysis. *Int J Tuberc Lung Dis*. 2019;23(7):783-796. doi:10.5588/ijtld.18.0433.
124. Kornfeld H, Singhal A. Enlisting the Host to Fight TB. *Chest*. 2018;153(6):1292-1293. doi:10.1016/j.chest.2018.01.011.
125. Shewade HD, Jeyashree K, Mahajan P, et al. Effect of glycemic control and type of diabetes treatment on unsuccessful TB treatment outcomes among people with TB-Diabetes: A systematic review. *PLoS One*. 2017;12(10):e0186697-17. doi:10.1371/journal.pone.0186697.

126. Yu X, Li L, Xia L, et al. Impact of metformin on the risk and treatment outcomes of tuberculosis in diabetics: A systematic review. *BMC Infect Dis*. 2019;19(1):859. doi:10.1186/s12879-019-4548-4.
127. Mtabho CM, Semvua HH, Van Den Boogaard J, et al. Effect of diabetes mellitus on TB drug concentrations in Tanzanian patients. *J Antimicrob Chemother*. 2019;74(12):3537–3545. doi:10.1093/jac/dkz368.
128. Alfarisi O, Mave V, Gaikwad S, et al. Effect of diabetes mellitus on the pharmacokinetics and pharmacodynamics of tuberculosis treatment. *Antimicrob Agents Chemother*. 2018;62(11):1–14. doi:10.1128/AAC.01383-18.
129. Dekkers BGJ, Akkerman OW, Alffenaar JWC. Role of therapeutic drug monitoring in treatment optimization in tuberculosis and diabetes mellitus comorbidity. *Antimicrob Agents Chemother*. 2019;63(2):1–7. doi:10.1128/AAC.02074-18.
130. Alkabab Y, Keller S, Dodge D, Houpt E, Staley D, Heysell S. Early interventions for diabetes related tuberculosis associate with hastened sputum microbiological clearance in Virginia, USA. *BMC Infect Dis*. 2017;17(1):1–8. doi:10.1186/s12879-017-2226-y.
131. Pradhan RP, Katz LA, Nidus BD, Matalon R, Eisinger RP. Tuberculosis in dialyzed patients. *JAMA*. 1974;229(7):798–800.
132. Dobler CC, McDonald SP, Marks GB. Risk of tuberculosis in dialysis patients: A nationwide cohort study. *PLoS One*. 2011;6(12):e29563–6. doi:10.1371/journal.pone.0029563.
133. Park S, Lee S, Kim Y, et al. Association of CKD with incident tuberculosis. *CJASN*. 2019;14(7):1002–1010. doi:10.2215/CJN.14471218.
134. Cho PJY, Wu CY, Johnston J, Wu MY, Shu CC, Lin HH. Progression of chronic kidney disease and the risk of tuberculosis: An observational cohort study. *Int J Tuberc Lung Dis*. 2019;23(5):555–562. doi:10.5588/ijtld.18.0225.
135. Yan M, Puyat JH, Shulha HP, Clark EG, Levin A, Johnston JC. Risk of tuberculosis associated with chronic kidney disease: a population-based analysis. *Nephrology Dialysis Transplantation*. 2021;37(1):197–116. doi:10.1093/ndt/gfab222.
136. Romanowski K, Clark EG, Levin A, Cook VJ, Johnston JC. Tuberculosis and chronic kidney disease: an emerging global syndemic. *Kidney Int*. 2016;90(1):34–40. doi:10.1016/j.kint.2016.01.034.
137. Dalrymple LS, Katz R, Kestenbaum B, et al. The risk of infection-related hospitalization with decreased kidney function. *American Journal of Kidney Diseases: The Official Journal of the National Kidney Foundation*. 2012;59(3):356–363. doi:10.1053/j.ajkd.2011.07.012.
138. Milburn H, Ashman N, Davies P, British Thoracic Society Standards of Care C, Joint Tuberculosis C, et al. Guidelines for the prevention and management of Mycobacterium tuberculosis infection and disease in adult patients with chronic kidney disease. *Thorax*. 2010;65(6):557–570. doi:10.1136/thx.2009.133173.
139. Stamatakis G, Montes C, Trouvin JH, et al. Pyrazinamide and pyrazinoic acid pharmacokinetics in patients with chronic renal failure. *Clin Nephrol*. 1988;30(4):230–234.
140. Varughese A, Brater DC, Benet LZ, Lee CSC. Ethambutol kinetics in patients with impaired renal function. *American Review of Respiratory Disease*. 1986;134(1):34–38. doi:10.1016/0278-2316(86)90005-8.
141. Ahn C, Oh KH, Kim K, et al. Effect of peritoneal dialysis on plasma and peritoneal fluid concentrations of isoniazid, pyrazinamide, and rifampin. *Perit Dial Int*. 2003;23(4):362–367. doi:10.1177/089686080302300409.
142. Si M, Li H, Chen Y, Peng H. Ethambutol and isoniazid induced severe neurotoxicity in a patient undergoing continuous ambulatory peritoneal dialysis. *BMJ Case Reports*. 2018;2018:bcr-2017-223187–5. doi:10.1136/bcr-2017-223187.
143. Saukkonen JJ, Cohn DL, Jasmer RM, ATS (American Thoracic Society) Hepatotoxicity of Antituberculosis Therapy Subcommittee, et al. An official ATS statement: Hepatotoxicity of antituberculosis therapy. *Am J Respir Crit Care Med*. 2006;174(8):935–952. doi:10.1164/rccm.200510-1666ST.
144. Yee D, Valiquette C, Pelletier M, Parisien I, Rocher I, Menzies D. Incidence of Serious Side Effects from First-Line Antituberculosis Drugs among Patients Treated for Active Tuberculosis. *Am J Respir Crit Care Med*. 2003;167(11):1472–1477. doi:10.1164/rccm.200206-626OC.
145. Alshammari TM, Larrat EP, Morrill HJ, Caffrey AR, Quilliam BJ, LaPlante KL. Risk of hepatotoxicity associated with fluoroquinolones: A national case-control safety study. *Am J Health Syst Pharm*. 2014;71(1):37–43. doi:10.2146/ajhp130165.
146. Paterson JM, Mamdani MM, Manno M, Juurlink DN, Canadian Drug Safety and Effectiveness Research Network Fluoroquinolone therapy and idiosyncratic acute liver injury: A population-based study. *CMAJ*. 2012;184(14):1565–1570. doi:10.1503/cmaj.111823.
147. Shah H, Bilodeau M, Burak KW, Canadian Association for the Study of the Liver, et al. The management of chronic hepatitis C: 2018 guideline update from the Canadian Association for the Study of the Liver. *CMAJ*. 2018;190(22):E677–E687. doi:10.1503/cmaj.170453.
148. Coffin CS, Fung SK, Alvarez F, et al. Management of Hepatitis B Virus Infection: 2018 Guidelines from the Canadian Association for the Study of the Liver and Association of Medical Microbiology and Infectious Disease Canada. *CanLivJ*. 2018;1(4):156–217. doi:10.3138/canlivj.2018-0008.
149. Chitnis AS, Cheung R, Gish RG, Wong RJ. Epidemiology and Prevention of Tuberculosis and Chronic Hepatitis B Virus Infection in the United States. *J Immigrant Minority Health*. 2021;23(6):1213–1267. doi:10.1007/s10903-021-01231-6.
150. Nooredinwand HA, Connell DW, Asgheddi M, et al. Viral hepatitis prevalence in patients with active and latent tuberculosis. *World J Gastroenterol*. 2015;21(29):8920–8980. doi:10.3748/wjg.v21.i29.8920.
151. Chua APG, Lim LKY, Gan SH, Chee CBE, Wang YT. The role of chronic viral hepatitis on tuberculosis treatment interruption. *Int j Tuberc Lung Dis*. 2018;22(12):1486–1494. doi:10.5588/ijtld.18.0195.
152. Chen L, Bao D, Gu L, et al. Co-infection with hepatitis B virus among tuberculosis patients is associated with poor outcomes during anti-tuberculosis treatment. *BMC Infect Dis*. 2018;18(1):1–10. doi:10.1186/s12879-018-3192-8.
153. Kim WS, Lee SS, Lee CM, et al. Hepatitis C and not Hepatitis B virus is a risk factor for anti-tuberculosis drug induced liver injury. *BMC Infect Dis*. 2016;16(1):1–7. doi:10.1186/s12879-016-1344-2.
154. Zheng J, Guo MH, Peng HW, Cai XL, Wu YL, Peng XE. The role of hepatitis B infection in anti-tuberculosis drug-induced liver injury: A meta-analysis of cohort studies. *Epidemiol Infect*. 2020;148:E290. doi:10.1017/S0950268820002861.
155. Ungo JR, Jones D, Ashkin D, et al. Antituberculosis drug-induced hepatotoxicity. The role of hepatitis C virus and the human immunodeficiency virus. *Am J Respir Crit Care Med*. 1998;157(6):1871–1876. doi:10.1164/ajrccm.157.6.9711039.
156. Lui GCY, Wong NS, Wong RYK, et al. Antiviral Therapy for Hepatitis B Prevents Liver Injury in Patients with Tuberculosis and Hepatitis B Coinfection. *Clin Infect Dis*. 2020;70(4):660–666. doi:10.1093/cid/ciz241.
157. Kempker RR, Alghamdi WA, Al-Shaer MH, Burch G, Peloquin CA. A Pharmacology Perspective of Simultaneous Tuberculosis and Hepatitis C Treatment. *Antimicrob Agents Chemother*. 2019;63(12):1–14. doi:10.1128/AAC.01215-19.
158. Zenner D, Kruijshaar ME, Andrews N, Abubakar I. Risk of tuberculosis in pregnancy: a national, primary care-based cohort and self-controlled case series study. *Am J Respir Crit Care Med*. 2012;185(7):779–784. doi:10.1164/rccm.201106-1083OC.
159. Singh N, Perfect JR. Immune reconstitution syndrome and exacerbation of infections after pregnancy. *Clin Infect Dis*. 2007;45(9):1192–1199. doi:10.1086/522182.
160. Jonsson J, Kuhlmann-Berenzon S, Berggren I, Bruchfeld J. Increased risk of active tuberculosis during pregnancy and postpartum: a register-based cohort study in Sweden. *Eur Respir J*. 2020;55(3):1901886. doi:10.1183/13993003.01886-2019.

161. Llewelyn M, Cropley I, Wilkinson RJ, Davidson RN. Tuberculosis diagnosed during pregnancy: a prospective study from London. *Thorax*. 2000;55(2):129–132. doi:10.1136/thorax.55.2.129.
162. Miele K, Bamrah Morris S, Tepper NK. Tuberculosis in Pregnancy. *Obstet Gynecol*. 2020;135(6):1444–1453. doi:10.1097/AOG.0000000000003890.
163. Jana N, Vasishta K, Jindal SK, Khunnu B, Ghosh K. Perinatal outcome in pregnancies complicated by pulmonary tuberculosis. *Int J Gynaecol Obstet*. Feb. 1994;44(2):119–124. doi:10.1016/0020-7292(94)90064-7.
164. Figueroa-Damian R, Arredondo-Garcia JL. Neonatal outcome of children born to women with tuberculosis. *Arch Med Res*. 2001;32(1):66–69. doi:10.1016/s0188-4409(00)00266-6.
165. Sobhy S, Babiker Z, Zamora J, Khan KS, Kunst H. Maternal and perinatal mortality and morbidity associated with tuberculosis during pregnancy and the postpartum period: a systematic review and meta-analysis. *BJOG*. 2017;124(5):727–733. doi:10.1111/1471-0528.14408.
166. S KJ, B A, H GK, G SM. Tuberculosis in pregnant women and neonates: A meta-review of current evidence. *Paediatr Respir Rev*. 2020;36:27–32. doi:10.1016/j.prrv.2020.02.001.
167. Gupta A, Hughes MD, Garcia-Prats AJ, McIntire K, Hesselning AC. Inclusion of key populations in clinical trials of new anti-tuberculosis treatments: Current barriers and recommendations for pregnant and lactating women, children, and HIV-infected persons. *PLoS Med*. 2019;16(8):e1002882. doi:10.1371/journal.pmed.1002882.
168. Snider D. Pregnancy and Tuberculosis. *Chest*. 1984;86(3 Suppl):10S–13S. doi:10.1378/chest.86.3.10S.
169. Water BJVD, Brooks MB, Huang C-c, et al. Tuberculosis clinical presentation and treatment outcomes in pregnancy: a prospective cohort study. *BMC Infect Dis*. 2020;20(1):686. doi: 10.1186/s12879-020-05416-6.
170. Abdelwahab MT, Leisegang R, Dooley KE, et al. Population Pharmacokinetics of Isoniazid, Pyrazinamide, and Ethambutol in Pregnant South African Women with Tuberculosis and HIV. *Antimicrob Agents Chemother*. 2020;64(3):e01978–19. doi: 10.1128/AAC.01978-19.
171. Gupta A, Montepiedra G, Aaron L, IMPAACT P1078 TB APPRISE Study Team, et al. Isoniazid Preventive Therapy in HIV-Infected Pregnant and Postpartum Women. *N Engl J Med*. 2019;381(14):1333–1346. doi:10.1056/nejmoa1813060.
172. Czeizel AE, Rockenbauer M, Olsen J, Sørensen HT. A population-based case-control study of the safety of oral anti-tuberculosis drug treatment during pregnancy. *Int J Tuberc Lung Dis*. 2001;5(6):564–568.
173. Bothamley G. Drug treatment for tuberculosis during pregnancy: safety considerations. *Drug Saf*. 2001;24(7):553–565. doi:10.2165/00002018-200124070-00006.
174. Treatment of tuberculosis: guidelines, World Health Organization Previous editions had different title: Treatment 2010.
175. Yefet E, Schwartz N, Chazan B, Salim R, Romano S, Nachum Z. The safety of quinolones and fluoroquinolones in pregnancy: a meta-analysis. *BJOG: Int J Obstet Gy*. 2018; 125(9):1069–1076. doi:10.1111/1471-0528.15119.
176. Pediatrics AAO. *Red Book: 2018 Report of the Committee on Infectious Diseases*. 31st ed. Itasca, Illinois: American Academy of Pediatrics; 2018.
177. Gallant V, Duvvuri V, McGuire M. Tuberculosis in Canada - Summary 2015. *Can Commun Dis Rep*. Mar 2 2017;43(3-4):77–82. doi:10.14745/ccdr.v43i34a04.
178. Hochberg NS, Horsburgh CR.Jr. Prevention of tuberculosis in older adults in the United States: obstacles and opportunities. *Clin Infect Dis*. 2013;56(9):1240–1247. Maydoi:10.1093/cid/cit027.
179. Symes MJ, Probyn B, Daneshvar C, Telisinghe L. Diagnosing Pulmonary Tuberculosis in the Elderly. *Curr Geri Rep*. 2020;9(2):30–39. doi:10.1007/s13670-020-00319-5.
180. Gardner Toren K, Spitters C, Pecha M, Bhattarai S, Horne DJ, Narita M. Tuberculosis in Older Adults: Seattle and King County, Washington. *Clin Infect Dis*. 2020;70(6):1202–1207. doi:10.1093/cid/ciz306.
181. Hase I, Toren KG, Hirano H, et al. Pulmonary Tuberculosis in Older Adults: Increased Mortality Related to Tuberculosis Within Two Months of Treatment Initiation. *Drugs and Aging*. 2021;38(9):807–815. doi:10.1007/s40266-021-00880-4.
182. Kwon BS, Kim Y, Lee SH, et al. The high incidence of severe adverse events due to pyrazinamide in elderly patients with tuberculosis. *PLoS One*. 2020;15(7):e0236109–10. doi:10.1371/journal.pone.0236109.
183. Marra F, Marra CA, Bruchet N, et al. Adverse drug reactions associated with first-line anti-tuberculosis drug regimens. *Int J Tuberc Lung Dis*. 2007;11(8):868–875.
184. Tweed CD, Wills GH, Crook AM, et al. Liver toxicity associated with tuberculosis chemotherapy in the REMoxTB study. *BMC Med*. 2018;16(1):46. doi:10.1186/s12916-018-1033-7.
185. Hosford JD, von Fricken ME, Lauzardo M, et al. Hepatotoxicity from antituberculous therapy in the elderly: a systematic review. *Tuberculosis (Edinb)*. 2015;95(2):112–122. doi:10.1016/j.tube.2014.10.006.
186. Lönnroth K, Williams BG, Stadlin S, Jaramillo E, Dye C. Alcohol use as a risk factor for tuberculosis - A systematic review. *BMC Public Health*. 2008;8(1):289. doi:10.1186/1471-2458-8-289.
187. Imtiaz S, Shield KD, Roerecke M, Samokhvalov AV, Lönnroth K, Rehm J. Alcohol consumption as a risk factor for tuberculosis: Meta-analyses and burden of disease. *Eur Respir J*. 2017;50(1):1700216. doi:10.1183/13993003.00216-2017.
188. Rehm J, Samokhvalov AV, Neuman MG, et al. The association between alcohol use, alcohol use disorders and tuberculosis (TB). A systematic review. *BMC Public Health*. 2009;9(1):412–450. doi:10.1186/1471-2458-9-450.
189. Storla DG, Yimer S, Bjune GA. A systematic review of delay in the diagnosis and treatment of tuberculosis. *BMC Public Health*. 2008;8(1):15–19. doi:10.1186/1471-2458-8-15.
190. Volkman T, Moonan PK, Miramontes R, Oeltmann JE. Tuberculosis and excess alcohol use in the United States, 1997–2012. *Int J Tuberc Lung Dis*. 2015;19(1):111–119. doi:10.5588/ijtld.14.0516.
191. Ragan EJ, Kleinman MB, Sweigart B, et al. The impact of alcohol use on tuberculosis treatment outcomes: a systematic review and meta-analysis. *Int J Tuberc Lung Dis*. 2020;24(1):73–82. doi:10.5588/ijtld.19.0080.
192. Samuels JP, Sood A, Campbell JR, Ahmad Khan F, Johnston JC. Comorbidities and treatment outcomes in multidrug resistant tuberculosis: A systematic review and meta-analysis. *Sci Rep*. 2018;8(1):1–13. doi:10.1038/s41598-018-23344-z.
193. Torres NMC, Rodríguez JJQ, Andrade PSP, Arriaga MB, Netto EM. Factors predictive of the success of tuberculosis treatment: A systematic review with meta-analysis. *PLoS One*. 2019;14(12):e0226507–24. doi:10.1371/journal.pone.0226507.
194. Burman WJ, Cohn DL, Rietmeijer CA, Judson FN, Sbarbaro JA, Reves RR. Noncompliance with directly observed therapy for tuberculosis: Epidemiology and effect on the outcome of treatment. *Chest*. 1997;111(5):1168–1173. doi:10.1378/chest.111.5.1168.
195. Deiss RG, Rodwell TC, Garfein RS. Tuberculosis and illicit drug use: Review and update. *Clin Infect Dis*. 2009;48(1):72–82. doi:10.1086/594126.
196. Minja LT, Hella J, Mbawambo J, et al. High burden of tuberculosis infection and disease among people receiving medication-assisted treatment for substance use disorder in Tanzania. *PLoS One*. 2021;16(4):e0250038. doi:10.1371/journal.pone.0250038.
197. Getahun H, Gunneberg C, Sculier D, Verster A, Raviglione M. Tuberculosis and HIV in people who inject drugs: Evidence for action for tuberculosis, HIV, prison and harm reduction services. *Curr Opin HIV Aids*. 2012;7(4):345–353. doi:10.1097/COH.0b013e328354bd44.
198. Duarte R, Lönnroth K, Carvalho C, et al. Tuberculosis, social determinants and co-morbidities (including HIV). *Pulmonology*. 2018;24(2):115–119. doi:10.1016/j.rppnen.2017.11.003.

199. Pablos-Méndez A, Knirsch CA, Barr RG, Lerner BH, Frieden TR. Nonadherence in tuberculosis treatment: Predictors and consequences in New York City. *American Journal of Medicine*. 1997;102(2):164–170. doi:10.1016/S0002-9343(96)00402-0.
200. Chaisson RE, Barnes GL, Hackman J, et al. A randomized, controlled trial of interventions to improve adherence to isoniazid therapy to prevent tuberculosis in injection drug users. *American Journal of Medicine*. 2001;110(8):610–615. doi:10.1016/S0002-9343(01)00695-7.
201. Batki SL, Gruber VA, Bradley JM, Bradley M, Delucchi K. A controlled trial of methadone treatment combined with directly observed isoniazid for tuberculosis prevention in injection drug users. *Drug Alcohol Depend*. 2002;66(3):283–293. doi:10.1016/S0376-8716(01)00208-3.
202. Smirnoff M, Goldberg R, Indyk L, Adler JJ. Directly observed therapy in an inner city hospital. *International Journal of Tuberculosis and Lung Disease*. 1998;2(2):134–139.
203. Schluger N, Ciotoli C, Cohen D, Johnson H, Rom WN. Comprehensive tuberculosis control for patients at high risk for noncompliance. *Am J Respir Crit Care Med*. 1995;151(5):1486–1490. doi:10.1164/ajrccm.151.5.7735604.
204. Raistrick D, Hay A, Wolff K. Methadone maintenance and tuberculosis treatment. *BMJ*. 1996;313(7062):925–926. doi:10.1136/bmj.313.7062.925.
205. McCance-Katz EF, Moody DE, Prathikanti S, Friedland G, Rainey PM. Rifampin, but not rifabutin, may produce opiate withdrawal in buprenorphine-maintained patients. *Drug Alcohol Depend*. 2011;118(2-3):326–334. doi:10.1016/j.drugalcdep.2011.04.013.
206. Brown LS, Sawyer RC, Li R, Cobb MN, Colborn DC, Narang PK. Lack of a pharmacologic interaction between rifabutin and methadone in HIV-infected former injecting drug users. *Drug Alcohol Depend*. 1996;43(1-2):71–77. doi:10.1016/S0376-8716(97)84352-9.
207. Bates MN, Khalakdina A, Pai M, Chang L, Lessa F, Smith KR. Risk of tuberculosis from exposure to tobacco smoke: A systematic review and meta-analysis. *Arch Intern Med*. 2007;167(4):335–342. doi:10.1001/archinte.167.4.335.
208. Van Zyl Smit RN, Pai M, Yew WW, et al. Global lung health: The colliding epidemics of tuberculosis, tobacco smoking, HIV and COPD. *Eur Respir J*. 2010;35(1):27–33. doi:10.1183/09031936.00072909.
209. Obore N, Kawuki J, Guan J, Papabathini SS, Wang L. Association between indoor air pollution, tobacco smoke and tuberculosis: an updated systematic review and meta-analysis. *Public Health*. 2020;187:24–35. doi:10.1016/j.puhe.2020.07.031.
210. Altet-Gómez MN, Alcaide J, Godoy P, Romero MA, Hernández Del Rey I. Clinical and epidemiological aspects of smoking and tuberculosis: A study of 13 038 cases. *International Journal of Tuberculosis and Lung Disease*. 2005;9(4):430–436.
211. Leung CC, Yew WW, Chan CK, et al. Smoking adversely affects treatment response, outcome and relapse in tuberculosis. *Eur Respir J*. 2015;45(3):738–745. doi:10.1183/09031936.00114214.
212. Burusie A, Enquesilassie F, Addissie A, Dessalegn B, Lamaro T. Effect of smoking on tuberculosis treatment outcomes: A systematic review and meta-analysis. *PLoS One*. 2020;15(9):e0239333–20. doi:10.1371/journal.pone.0239333.
213. Gajalakshmi V, Peto R, Kanaka TS, Jha P. Smoking and mortality from tuberculosis and other diseases in India: Retrospective study of 43 000 adult male deaths and 35 000 controls. *Lancet*. 2003;362(9383):507–515. doi:10.1016/S0140-6736(03)14109-8.
214. Yen YF, Yen MY, Lin YS, et al. Smoking increases risk of recurrence after successful anti-tuberculosis treatment: A population-based study. *Int J Tuberc Lung Dis*. 2014;18(4):492–498. doi:10.5588/ijtld.13.0694.
215. Whitehouse E, Lai J, Golub JE, Farley JE. A systematic review of the effectiveness of smoking cessation interventions among patients with tuberculosis. *Public Health Action*. 2018;8(2):37–49. doi:10.5588/pha.18.0006.
216. Jeyashree K, Kathirvel S, Shewade HD, Kaur H, Goel S, Cochrane Tobacco Addiction Group. Smoking cessation interventions for pulmonary tuberculosis treatment outcomes. *The Cochrane Database of Systematic Reviews*. 2016;CD011125. doi:10.1002/14651858.CD011125.pub2.
217. Alfenaar JWC, Gumbo T, Dooley KE, et al. Integrating pharmacokinetics and pharmacodynamics in operational research to end tuberculosis. *Clin Infect Dis*. 2020;70(8):1774–1780. doi:10.1093/cid/ciz942.
218. Hall RG, Leff RD, Gumbo T. Treatment of active pulmonary tuberculosis in adults: Current standards and recent advances. Insights from the Society of Infectious Diseases Pharmacists. *Pharmacotherapy*. 2009;29(12):1468–1481. doi:10.1592/phco.29.12.1468.
219. Abrogoua DP, Kamenan BAT, Ahui BJM, Doffou E. Pharmaceutical interventions in the management of tuberculosis in a pneumophthiology department, ivory coast. *Ther Clin Risk Manag*. 2016;12:1749–1756. doi:10.2147/TCRM.S118442.
220. Reitman ML, Chu X, Cai X, et al. Rifampin's acute inhibitory and chronic inductive drug interactions: Experimental and model-based approaches to drug-drug interaction trial design. *Clin Pharmacol Ther*. 2011;89(2):234–242. doi:10.1038/clpt.2010.271.
221. Sekaggya-Wiltshire C, Dooley KE. Pharmacokinetic and pharmacodynamic considerations of rifamycin antibiotics for the treatment of tuberculosis. *Expert Opin Drug Metab Toxicol*. 2019;15(8):615–618. doi:10.1080/17425255.2019.1648432.
222. Dooley KE, Bliven-Sizemore EE, Weiner M, et al. Safety and pharmacokinetics of escalating daily doses of the antituberculosis drug rifapentine in healthy volunteers. *Clin Pharmacol Ther*. 2012;91(5):881–888. doi:10.1038/clpt.2011.323.
223. Sahasrabudhe V, Zhu T, Vaz A, Tse S. Drug Metabolism and Drug Interactions: Potential Application to Antituberculosis Drugs. *J Infect Dis*. 2015;211(suppl 3):S107–S114. doi:10.1093/infdis/jiv009.
224. McFeely SJ, Yu J, Zhao P, et al. Drug-Drug Interactions of Infectious Disease Treatments in Low-Income Countries: A Neglected Topic? *Clin Pharmacol Ther*. 2019;105(6):1378–1385. doi:10.1002/cpt.1397.
225. Yang CS, Boswell R, Bungard TJ. A case series of the rifampin-warfarin drug interaction: focus on practical warfarin management. *Eur J Clin Pharmacol*. 2021;77(3):341–348. doi:10.1007/s00228-020-03057-x.
226. Martins MAP, Reis AMM, Sales MF, et al. Rifampicin-warfarin interaction leading to macroscopic hematuria: A case report and review of the literature. *BMC Pharmacol Toxicol*. 2013;14(1):10–27. doi:10.1186/2050-6511-14-27.
227. Wiggins BS, Dixon DL, Neyens RR, Page RL, Gluckman TJ. Select Drug-Drug Interactions With Direct Oral Anticoagulants: JACC Review Topic of the Week. *J Am Coll Cardiol*. 2020;75(11):1341–1350. doi:10.1016/j.jacc.2019.12.068.
228. Lutz JD, Kirby BJ, Wang L, et al. Cytochrome P450 3A Induction Predicts P-glycoprotein Induction; Part 2: Prediction of Decreased Substrate Exposure After Rifabutin or Carbamazepine. *Clin Pharmacol Ther*. 2018;104(6):1191–1198. doi:10.1002/cpt.1072.
229. Chang SH, Chou IJ, Yeh YH, et al. Association between use of Non-Vitamin K Oral Anticoagulants With and Without Concurrent Medications and Risk of Major Bleeding in Nonvalvular Atrial Fibrillation. *JAMA - Journal of the American Medical Association*. 2017;318(13):1250–1259. doi:10.1001/jama.2017.13883.
230. Mendell J, Chen S, He L, Desai M, Parasramupria DA. The effect of rifampin on the pharmacokinetics of edoxaban in healthy adults. *Clin Drug Investig*. 2015;35(7):447–453. doi:10.1007/s40261-015-0298-2.
231. Vakkalagadda B, Frost C, Byon W, et al. Effect of rifampin on the pharmacokinetics of apixaban, an oral direct inhibitor of factor xa. *Am J Cardiovasc Drugs*. 2016;16(2):119–127. doi:10.1007/s40256-015-0157-9.
232. Kay L, Kampmann J, Svendsen T, et al. Influence of rifampicin and isoniazid on the kinetics of phenytoin. *Br J Clin Pharmacol*. 1985;20(4):323–326. doi:10.1111/j.1365-2125.1985.tb05071.x.
233. Ebert U, Thong NQ, Oertel R, Kirch W. Effects of rifampicin and cimetidine on pharmacokinetics and pharmacodynamics

- of lamotrigine in healthy subjects. *European Journal of Clinical Pharmacology*. 2000;56(4):299–304. doi:10.1007/s002280000146.
234. Wimpelmann J, Høvik H, Riedel B, Slørdal L. The interaction between rifampicin and lamotrigine: A case report. *Br J Clin Pharmacol*. 2019;85(8):1859–1860. doi:10.1111/bcp.13973.
 235. Egelund EF, Mohamed MEF, Fennelly KP, Peloquin CA. Concomitant use of carbamazepine and rifampin in a patient with mycobacterium avium complex and seizure disorder. *J Pharm Technol*. 2014;30(3):93–96. doi:10.1177/8755122514523934.
 236. Niemi M, Backman JT, Neuvonen M, Neuvonen PJ. Effect of rifampicin on the pharmacokinetics and pharmacodynamics of nateglinide in healthy subjects. *Br J Clin Pharmacol*. 2003;56(4):427–432. doi:10.1046/j.1365-2125.2003.01884.x.
 237. Niemi M, Backman JT, Neuvonen M, Neuvonen PJ, Kivistö KT. Effects of rifampin on the pharmacokinetics and pharmacodynamics of glyburide and glipizide. *Clinical Pharmacology and Therapeutics*. 2001;69(6):400–406. doi:10.1067/mcp.2001.115822.
 238. Niemi M, Backman JT, Neuvonen PJ. Effects of trimethoprim and rifampin on the pharmacokinetics of the cytochrome P450 2C8 substrate rosiglitazone. *Clinical Pharmacology and Therapeutics*. 2004;76(3):239–249. doi:10.1016/j.clpt.2004.05.001.
 239. Niemi M, Backman JT, Fromm MF, Neuvonen PJ, Kivistö KT. Pharmacokinetic interactions with rifampicin : clinical relevance. *Clin Pharmacokinet*. 2003;42(9):819–850. doi:10.2165/00003088-200342090-00003.
 240. Park JY, Kim KA, Kang MH, Kim SL, Shin JG. Effect of rifampin on the pharmacokinetics of rosiglitazone in healthy subjects. *Clin Pharmacol Ther*. 2004;75(3):157–162. doi:10.1016/j.clpt.2003.10.003.
 241. Sahi J, Black CB, Hamilton GA, et al. Comparative effects of thiazolidinediones on in vitro P450 enzyme induction and inhibition. *Drug Metab Dispos*. 2003;31(4):439–446. doi:10.1124/dmd.31.4.439.
 242. Jaakkola T, Backman JT, Neuvonen M, Laitila J, Neuvonen PJ. Effect of rifampicin on the pharmacokinetics of pioglitazone. *Br J Clin Pharmacol*. Jan 2006;61(1):70–78. doi:10.1111/j.1365-2125.2005.02515.x.
 243. Riza AL, Pearson F, Ugarte-Gil C, et al. Clinical management of concurrent diabetes and tuberculosis and the implications for patient services. *Lancet Diabetes Endocrinol*. 2014;2(9):740–753. Sep doi:10.1016/S2213-8587(14)70110-X.
 244. Tucker RM, Denning DW, Hanson LH, et al. Interaction of azoles with rifampin, phenytoin, and carbamazepine: in vitro and clinical observations. *Clin Infect Dis*. 1992;14(1):165–174. doi:10.1093/clinids/14.1.165.
 245. Hohmann C, Kang EM, Jancel T. Rifampin and posaconazole coadministration leads to decreased serum posaconazole concentrations. *Clin Infect Dis*. 2010;50(6):939–940. doi:10.1086/650740.
 246. Geist MJ, Egerer G, Burhenne J, Riedel KD, Mikus G. Induction of voriconazole metabolism by rifampin in a patient with acute myeloid leukemia: importance of interdisciplinary communication to prevent treatment errors with complex medications. *Antimicrob Agents Chemother*. 2007;51(9):3455–3456. doi:10.1128/AAC.00579-07.
 247. Townsend R, Dietz A, Hale C, et al. Pharmacokinetic Evaluation of CYP3A4-Mediated Drug-Drug Interactions of Isavuconazole With Rifampin, Ketoconazole, Midazolam, and Ethinyl Estradiol/Norethindrone in Healthy Adults. *Clin Pharmacol Drug Dev*. 2017;6(1):44–53. doi:10.1002/cpdd.285.
 248. Ayudhya DPN, Thanompuangserree N, Tansuphaswadikul S. Effect of rifampicin on the pharmacokinetics of fluconazole in patients with AIDS. *Clin Pharmacokinet*. 2004;43(11):725–732. doi:10.2165/00003088-200443110-00003.
 249. Kirch W, Rose I, Klingmann I, Pabst J, Ohnhaus EE. Interaction of bisoprolol with cimetidine and rifampicin. *Eur J Clin Pharmacol*. 1986;31(1):59–62. doi:10.1007/BF00870987.
 250. Tada Y, Tsuda Y, Otsuka T, et al. Case report: Nifedipine-rifampicin interaction attenuates the effect on blood pressure in a patient with essential hypertension. *Am J Med Sci*. 1992;303(1):25–27. doi:10.1097/00000441-199201000-00006.
 251. Williamson KM, Patterson JH, McQueen RH, Adams KE, Pieper JA. Effects of erythromycin or rifampin on losartan pharmacokinetics in healthy volunteers. *Clin Pharmacol Ther*. 1998;63(3):316–323. doi:10.1016/S0009-9236(98)90163-1.
 252. Liu W, Yan T, Chen K, Yang L, Benet LZ, Zhai S. Predicting Interactions between Rifampin and Antihypertensive Drugs Using the Biopharmaceutics Drug Disposition Classification System. *Pharmacotherapy*. 2020;40(4):274–290. doi:10.1002/phar.2380.
 253. McAllister WA, Thompson PJ, Al-Habet SM, Rogers HJ. Rifampicin reduces effectiveness and bioavailability of prednisolone. *Br Med J (Clin Res Ed)*. 1983;286(6369):923–925. doi:10.1136/bmj.286.6369.923.
 254. Powell-Jackson PR, Gray BJ, Heaton RW, Costello JE, Williams R, English J. Adverse effect of rifampicin administration on steroid-dependent asthma. *Am Rev Respir Dis*. 1983;128(2):307–310. doi:10.1164/arrd.1983.128.2.307.
 255. Lee KH, Shin JG, Chong WS, et al. Time course of the changes in prednisolone pharmacokinetics after co-administration or discontinuation of rifampin. *Eur J Clin Pharmacol*. 1993;45(3):287–289. doi:10.1007/BF00315399.
 256. Bhaloo S, Prasad GV. Severe reduction in tacrolimus levels with rifampin despite multiple cytochrome P450 inhibitors: a case report. *Transplant Proc*. 2003;35(7):2449–2451. doi:10.1016/j.transproceed.2003.08.019.
 257. van Buren D, Wideman CA, Ried M, et al. The antagonistic effect of rifampin upon cyclosporine bioavailability. *Transplant Proc*. 1984;16(6):1642–1645.
 258. Hebert ME, Fisher RM, Marsh CL, Dressler D, Bekersky I. Effects of rifampin on tacrolimus pharmacokinetics in healthy volunteers. *J Clin Pharmacol*. 1999;39(1):91–96. doi:10.1177/00912709922007499.
 259. Wang Y-C, Salvador NG, Lin C-C, et al. Comparative analysis of the drug-drug interaction between immunosuppressants, safety and efficacy of rifabutin from rifampicin-based Anti-TB treatment in living donor liver transplant recipients with active tuberculosis. *Biomedical Journal*. 2020;1–9. doi:10.1016/j.bj.2020.08.010.
 260. Strayhorn VA, Baciewicz AM, Self TH. Update on rifampin drug interactions, III. *Arch Intern Med*. Nov 24 1997;157(21):2453–2458.
 261. Nolan SR, Self TH, Norwood JM. Interaction between rifampin and levothyroxine. *South Med J*. 1999;92(5):529–531. doi:10.1097/00007611-199905000-00018.
 262. Kim HI, Kim TH, Kim H, et al. Effect of Rifampin on thyroid function test in patients on levothyroxine medication. *PLoS One*. 2017;12(1):e0169775–11. doi:10.1371/journal.pone.0169775.
 263. Takasu N, Kinjou Y, Kouki T, Takara M, Ohshiro Y, Komiya I. Rifampin-induced hypothyroidism. *J Endocrinol Invest*. 2006;29(7):645–649. doi:10.1007/BF03344165.
 264. Kreek MJ, Garfield JW, Gutjahr CL, Giusti LM. Rifampin-induced methadone withdrawal. *N Engl J Med*. 1976;294(20):1104–1106. doi:10.1056/NEJM197605132942008.
 265. Badhan RKS, Gittins R, Al Zabit D. The optimization of methadone dosing whilst treating with rifampicin: A pharmacokinetic modeling study. *Drug Alcohol Depend*. 2019;200:168–180. doi:10.1016/j.drugalcdep.2019.03.013.
 266. Simmons KB, Haddad LB, Nanda K, Curtis KM. Drug interactions between rifamycin antibiotics and hormonal contraception: a systematic review. *BJOG*. 2018;125(7):804–811. doi:10.1111/1471-0528.15027.