



Chapter 7: Extra-pulmonary tuberculosis

Leila Barss, William J. A. Connors & Dina Fisher

To cite this article: Leila Barss, William J. A. Connors & Dina Fisher (2022) Chapter 7: Extra-pulmonary tuberculosis, Canadian Journal of Respiratory, Critical Care, and Sleep Medicine, 6:sup1, 87-108, DOI: [10.1080/24745332.2022.2036073](https://doi.org/10.1080/24745332.2022.2036073)

To link to this article: <https://doi.org/10.1080/24745332.2022.2036073>



Published online: 25 Mar 2022.



Submit your article to this journal [↗](#)



Article views: 8579



View related articles [↗](#)



View Crossmark data [↗](#)



Citing articles: 1 View citing articles [↗](#)

Chapter 7: Extra-pulmonary tuberculosis

Leila Barss^a, William J. A. Connors^b and Dina Fisher^a

^aDivision of Respiratory Medicine, Peter Lougheed Centre, Calgary, Alberta, Canada; ^bDivision of Infectious Diseases, Department of Medicine, University of British Columbia, British Columbia Centre for Disease Control, Vancouver, British Columbia, Canada

KEY POINTS

Diagnosis

- Samples (fluid and tissue) for extra-pulmonary tuberculosis (TB) should be sent for acid-fast bacilli smear, mycobacterial culture and nucleic acid amplification test. Tissue biopsy should be sent in sterile saline for these mycobacterial tests as well as in formalin for histopathologic assessment. Drug susceptibility testing should be requested for positive culture samples.
- If the specimen is insufficient for all testing, mycobacterial culture should be prioritized given it has the highest diagnostic yield and allows for gold-standard phenotypic drug testing.
- Every person with presumed extra-pulmonary TB should also be assessed for pulmonary TB to assess infectiousness and potentially assist with diagnosis.

intravenously [IV]) along with standard dose isoniazid, pyrazinamide, and ethambutol, during intensive phase of treatment for drug susceptible TB meningitis (TBM).

4. We conditionally recommend against using fluoroquinolone for TBM unless there is a concern regarding drug resistance.
5. We conditionally recommend initial adjunctive corticosteroid treatment in all human immunodeficiency virus (HIV)-negative patients with TB pericarditis. Routine use of adjunctive corticosteroids in people with HIV NOT on antiretroviral treatment is NOT recommended. Lack of data means we are unable to provide a specific recommendation on use of adjunctive corticosteroids in people with HIV ON antiretroviral treatment with TB pericarditis and suggest assessing on a case-by-case basis pending further study.

Treatment

- A standard 6-month anti-TB treatment course for microbiologically confirmed, drug-susceptible disease is recommended for most forms of extra-pulmonary TB. In patients with meningitis or bone and joint TB, treatment may be extended to 12 months.
- In the absence of culture confirmation and drug susceptibility results, the continuation phase of empiric anti-TB treatment should include a third agent such as ethambutol.
- Empiric anti-TB therapy should be considered in suspected life-threatening extra-pulmonary TB while appropriate diagnostic samples are being obtained.

1. Introduction

Internationally, pulmonary TB is defined as disease involving the lung parenchyma or tracheobronchial tree. Extra-pulmonary TB (EPTB) is defined as disease involving any other organs and includes pleural and isolated intrathoracic lymph node TB.¹ This differs from the terminology for reporting in Canada, which classifies TB as either respiratory (lungs and the conducting airways, pleural, fibrosis of the lung, bronchiectasis, pneumonia, pneumothorax, primary, intrathoracic and mediastinal lymph nodes, isolated tracheal or bronchial, laryngitis, nasopharynx, nose and sinus TB) or nonrespiratory (all other disease sites not listed).² This chapter focuses on the diagnosis and treatment of EPTB.

Over the last 10 years in Canada, the incidence of EPTB has remained stable, similar to other low TB-incidence, and low HIV-TB co-infection countries.³ Peripheral TB lymphadenitis, pleural TB and abdominal TB have remained the three most common types of EPTB (see [Chapter 1: Epidemiology of Tuberculosis in Canada](#)). Impaired host immune status is a risk factor for EPTB and increases associated mortality. Organ transplant recipients, persons with advanced HIV and those undergoing chemotherapy for malignancies are at particularly high risk of life-threatening disease.^{4–9}

NEW AND UPDATED RECOMMENDATIONS

1. We conditionally recommend against routine adjunctive corticosteroid use for pleural TB.
2. We conditionally recommend against routine therapeutic thoracentesis/chest tube drainage for pleural TB-associated effusions.
3. We conditionally recommend using higher doses of rifampin (greater than 15 mg/kg/day orally up to maximum dose 35 mg/kg/day or 15 mg/kg/day



Table 1. Sensitivity and specificity of diagnostic tests in extra-pulmonary tuberculosis compared to mycobacterial culture as gold standard unless otherwise stated.

| Site | Specimen type | Direct stain (ZN) | | Culture | | GeneXpert ^a | | GeneXpert Ultra ^a | | Histopathology and/or cytology | |
|------------------------------|----------------------------------|------------------------|-------------|------------------------|-------------|--------------------------|--------------------------|------------------------------|--------------------------|--------------------------------|---------------------|
| | | Sensitivity | Specificity | Sensitivity | Specificity | Sensitivity ^b | Specificity ^b | Sensitivity ^b | Specificity ^b | Sensitivity | References |
| TB lymphadenitis | FNA | 0.23-0.37 | | 0.17-0.67 | | 0.89 | 0.86 | INS | | 0.62-0.80 ^d | 10-14 |
| | Excisional biopsy | 0.35-0.53 | | 0.71-0.80 | | 0.82 ^c | 0.80 ^c | | | 0.85-1.00 ^d | |
| | Pleural Fluid | 0-0.10 | | 0.10-0.63 | | 0.50 | 0.99 | 0.75 | 0.87 | N/A | |
| Pleural TB | Closed pleural biopsy | 0.13-0.39 | | 0.39-0.67 | | 0.19 ^d | 0.99 ^d | | | | 14-24 |
| | Thorascopic pleural biopsy | INS | | 0.41-0.76 | | 0.31 | 0.97 | INS | | 0.69-0.97 ^d | |
| | CSF | 0.05-0.83 ^g | | 0.40-0.87 ^g | | 0.52 ^e | 1.0 ^e | | | 1.0 ^{df} | |
| CNS - meningitis | | | | | | 0.71 | 0.97 | 0.89 ^h | 0.91 ^h | N/A | 23,25-35 |
| | | | | | | | | 0.63 ^{dh} | 0.99 ^{dh} | | |
| | | | | | | | | | | | |
| CNS - tuberculoma | FNA | INS | | INS | | INS | | INS | | 0.85-0.92 ^d | 23,36-49 |
| | Biopsy (excisional/stereotactic) | 0.17-0.38 ^f | | 0.5-0.8 ^f | | 0.87-1 ^f | 0.89-0.96 ^f | INS | | 1 ^f | |
| | Feces | 0-0.11 ⁱ | | 0.44-0.5 ⁱ | | 0.39 ^d | 0.86 ^d | INS | | INS | |
| Abdominal TB | Ascitic fluid | 0.0-0.06 | | 0.17-0.80 | | 0.5 | 0.98 | INS | | | 0.76-1 ^d |
| | Peritoneal biopsy | 0.02-0.20 | | 0.34-0.92 | | 0.38-0.5 ^{df} | 0.92-1.0 ^{df} | INS | | | |
| | Intestinal biopsy | 0.03-0.14 | | 0.36-0.40 | | 0.08-0.32 ^{df} | 1.0 ^{df} | INS | | 0.16-0.7 ^d | |
| GU TB - renal | Urine | 0.15-0.30 | | 0.80-0.90 | | 0.83 | 0.98 | 1.00 ^j | 1.00 ^j | N/A | 23,25,50-71 |
| | FNA/biopsy | 0.29-0.44 | | 0.95-1.0 ^{hj} | | | | INS | | 0.88-0.95 ^f | |
| | Urine | 0.20-0.24 | | 0.63-0.93 ^b | | 0.83 | 0.98 | INS | | N/A | |
| GU TB - scrotal | FNA/Biopsy | 0.25-0.75 | | 0.8 ^e | | INS | | INS | | 0.95 ^{de} | N/A |
| | Menstrual fluid | 0.03-0.05 | | 0.06-0.19 | | INS | | INS | | N/A | |
| | Endometrial biopsy | 0.05-0.57 | | 0.06-0.46 | | INS | | INS | | 0.07-0.71 ^d | |
| Bone and Joint TB | FNA bone/synovial tissue | 0.30-0.36 | | 0.37-0.97 | | 0.95 | 0.85 | 0.96 ^k | 0.97 ^k | 0.56-0.97 | 23,50-53,72-80 |
| | Synovial fluid | 0.08-0.26 | | 0.50-0.86 | | 0.97 | 0.90 | 0.96 ^k | 0.97 ^k | n/a | |
| | Paraspinal fluid | 0.52-0.59 | | 0.9-0.93 | | 0.97 | 0.90 | 0.96 ^k | 0.97 ^k | n/a | |
| Pericardial TB | Pericardial fluid | 0-0.42 ⁱ | | 0.20-0.86 | | 0.61 | 0.90 | INS | | N/A | 13,14,23,81-88 |
| | Pericardial biopsy | 0.38-0.40 ^j | | 0.22-1.0 | | 0.71 ^{hj} | 1.0 ^{hj} | | | 0.34-0.87 ^d | |
| | Sputum | 0.31-0.37 | | 0.32-0.90 | | | | INS | | INS | |
| Disseminated TB ^l | Bronchial wash | 0.20-0.55 | | 0.07-0.71 | | INS | | INS | | INS | 23,89-96 |
| | Lung biopsy | 0.25-0.43 | | 0.42-0.54 | | INS | | INS | | 0.63-0.95 | |
| | Liver biopsy | 0.13-0.40 | | 0.33-0.54 | | INS | | INS | | 0.88-1.00 | |
| | Bone marrow | 0.16-0.25 | | 0.21-0.67 | | INS | | INS | | 0.56-0.67 | INS |
| | Urine | 0.00-0.18 | | 0.33-0.67 | | INS | | INS | | INS | |
| | Blood | INS | | 0.20-0.65 | | 0.07-0.56 | 0.94-1.0 | INS | | INS | |

Abbreviations: TB, tuberculosis; ZN, Ziehl-Neelsen; SN, sensitivity; SP, specificity; FNA, fine-needle aspiration; INS, insufficient reported data; CNS, central nervous system; CSF, cerebrospinal fluid; GU, genitourinary; N/A, nonapplicable; HIV, human immunodeficiency virus.

Reported data from single studies and meta-analyses included a minimum of 50 patients unless otherwise noted.

Reported data from single studies come from HIV-negative cohorts or studies with a TB-HIV coinfection rate of less than 5% (where reported and if available). Data from meta-analyses that included studies with HIV prevalence greater than 5% and that reported a difference in diagnostics based on HIV status are noted.

^aGeneXpert and GeneXpert Ultra data provided as data from individual NAAT available in single site small studies.

^bPoint estimates used for data provided from meta-analyses.

^cData from Cochrane review does not define method of obtaining biopsy sample.¹⁴

^dComposite reference standard definition: Positive result as the presence of granulomatous inflammation or a positive microbiologic result +/- clinical diagnosis of TB and/or improvement on TB treatment.

^eSingle study data.

^fData limited to two studies (combined n > 50).

^gHigher range sensitivity with larger volume (10 cc, 3 samples) concentrated samples (centrifuged at 3000g).

^hMajority of people with HIV and from single study.²²

ⁱMajority of studies do not explicitly state if abdominal TB or intestinal TB present (many state disseminated disease). Presence of stool AFB and positive culture/Xpert documented in isolated pulmonary disease (primarily in children and HIV/AIDS studies).

^jLess than 50 patients.

^kBiopsy and fluid samples combined.

^lDisseminated TB includes miliary TB, as some included autopsy data did not allow differentiation.

2. General diagnostic considerations

A high index of suspicion is paramount to the diagnosis of EPTB. Any delay in diagnosis could increase the risk of morbidity and mortality.¹⁸ In at-risk patients with fever of unknown origin and site-specific signs and symptoms, or patients with biopsy-proven granulomatous inflammation, appropriate steps should be taken to confirm the diagnosis of TB, including repeat sampling if mycobacterial cultures were not obtained.

Gold-standard phenotypic drug susceptibility testing can only proceed with a viable culture, the results of which can have important treatment implications.^{7,19,20} This point cannot be overemphasized: with the rising incidence of drug-resistant TB, especially in foreign-born residents of Canada, it is difficult to provide appropriate treatment when mycobacterial cultures and drug susceptibility test results are not available.

The clinical specimens obtained for diagnostic purposes will depend upon the suspected anatomic site of involvement. In general, tissue biopsy yields positive nucleic acid amplification test (NAAT) and culture results more often than fluid aspiration; both are superior to swabs (see Table 1 for diagnostic yield estimates). Biopsy material for mycobacterial culture should be submitted fresh or in a small amount of sterile saline.^{19,20} Histopathologic examination requires the specimen to be placed in formalin, which kills the mycobacteria and prevents further culture confirmation.^{19,20} Common histopathologic findings include necrotizing and non-necrotizing granulomas (Table 1). Loss of host immune function can result in histopathologic findings demonstrating greater suppurative response and less well-formed granulomas.⁸⁸

Good practice statement

- Every effort should be made to obtain clinical samples for both mycobacteriologic (acid-fast bacilli smear, culture, nucleic acid amplification test) and histopathologic tests. If the specimen is insufficient for all testing, mycobacterial culture should be prioritized given it has the highest diagnostic yield and allows for gold-standard phenotypic drug testing.

Anywhere from 10 to 50% of patients with EPTB also have pulmonary involvement; it may, therefore, be possible to confirm a diagnosis of TB (and obtain drug susceptibility testing) with sputa assessments, averting the need for more invasive sampling.²⁰

Good practice statement

- Every person with presumed extra-pulmonary TB should be assessed for pulmonary TB to assess infectiousness and potentially assist with diagnosis.

3. Clinical presentations by organ-specific site

Clinical presentation, investigations and management by organ specific site are reviewed in the following sections.

Evidence for treatment duration and other adjunct measures for each organ site is summarized in Table 3. Examples of published corticosteroid regimens for relevant organ sites are described in Table 2.

3.1. Peripheral TB lymphadenitis

Almost all forms of TB involve regional lymphatics and nodes. This section will focus on extrathoracic lymph nodes and, specifically, peripheral TB lymphadenitis.

Mycobacterial involvement of the lymph glands can be secondary to infection from *Mycobacterium tuberculosis* (*M. tuberculosis*) as well as other non-TB mycobacteria (NTM).⁹⁷ Unilateral cervical chain involvement is the most common site of TB lymphadenitis (45 to 80%) but lymphadenitis can occur in the supraclavicular and axillary regions, as well as a variety of other nodal sites.^{98–100} Presentation can be at a single nodal site or in multiple sites.

In general, the disease is most often indolent, and the patient usually presents with an isolated, unilateral, non-tender neck mass.⁹⁸ The term “scrofula” has been used historically to describe TB involvement of a cervical lymph node with sinus tract formation or ulceration of the overlying skin. Non-nodal symptoms are rare, except in people with HIV.^{101,102}

Fine-needle aspiration (FNA) biopsy of affected lymph nodes is a useful initial procedure (see Table 1). If FNA is nondiagnostic, the highest-yield procedure is an excisional lymph node biopsy. If there is high suspicion for an alternate diagnosis (eg, lymphoma), clinicians may choose to pursue excisional biopsy as the initial diagnostic test. Incisional biopsies are discouraged for initial testing because of the risk of sinus tract formation at the biopsy site in mycobacterial disease.¹⁰³ Swabs are discouraged because of the limited material obtained and because the hydrophobic nature of the mycobacterial cell wall inhibits the transfer of organisms from the swab to the culture media.¹⁰⁴ Specimens should be submitted for both mycobacteriologic (smear, culture and NAAT) and histopathologic analysis. Differentiation of *M. tuberculosis* from *M. avium complex* is important, as treatment of the 2 conditions is different.

3.1.1. TB lymphadenitis treatment

Two small randomized controls trials that compared 6- versus 9-month treatment courses did not report any difference in treatment completion or relapse rates.^{105,106} Three systematic reviews that also included observational studies reported the same findings.^{107–109}

Recommendation

- We conditionally recommend six months of standard anti-TB therapy for treating drug-susceptible TB lymphadenitis (poor evidence).

In up to 23% of patients, nodes can appear afresh or enlarge during treatment, possibly as an immune response.

This usually will resolve without change in regime or additional therapy and should not be considered evidence of treatment failure.¹¹⁰ At the end of treatment, up to 34% of patients may be left with residual nodes and, if after treatment, the nodes enlarge or reappear, this is usually transient.¹⁰⁷ Such events do not necessarily imply relapse, but repeat FNA for mycobacterial culture should be considered to assess this possibility if the nodes are persistent.^{111,112}

Good practice statement

- **In TB lymphadenitis, surgical/drainage procedures, other than diagnostic, should be reserved for the relief of discomfort caused by enlarged nodes or tense, fluctuant nodes.**

3.2. Pleural TB

TB pleural-space disease can range from simple TB pleural effusion to TB empyema. Fever, unilateral pleuritic chest pain and cough are the most frequent presenting symptoms in TB pleural effusion. Dyspnea, night sweats and weight loss are also common.¹¹³ TB pleural effusion is typically paucibacillary and often culture negative.¹¹³ In some patients, TB pleural effusion may spontaneously resolve within 2 to 4 months without treatment; however, up to 65% will subsequently develop active TB within the following 5 years.¹¹⁴

TB empyema is less common than TB pleural effusion and characterized by purulent pleural space and presence of acid-fast bacilli on fluid microscopy.¹¹⁵

TB pleural effusions are usually unilateral and can occur on either side of the chest.^{15,115,116} Effusions are usually small to moderate in size but in some cases can occupy over two-thirds of the hemithorax.^{15,116} Co-existing parenchymal abnormalities on chest x-ray are reported in 19 to 67% of patients.^{17,116,117} Chest computed tomography (CT) imaging usually demonstrates smooth pleural thickening with an associated effusion. Parenchymal abnormalities on chest CT are reported in up to 86% of cases.^{117,118} Pleural ultrasound can assist in diagnostic procedures for pleural TB, but imaging findings are nonspecific.¹¹⁹

Even in patients without parenchymal abnormalities on chest x-ray, the yield of induced sputum culture for *M. tuberculosis* is over 50%.^{15,120} TB pleural fluid is exudative and typically straw colored. Pleural fluid glucose may be low or normal. Pleural fluid pH is usually above 7.3. The majority of TB pleural effusions are lymphocyte predominant. However, a neutrophil predominance may be seen in the very early stages of infection or if TB empyema develops.^{113,121} Pleural fluid adenosine deaminase (ADA) testing is not used in Canada (see Chapter 3: Diagnosis of Tuberculosis Disease and Drug-resistant Tuberculosis). Diagnosis is based on bacteriologic confirmation (positive acid-fast bacilli [AFB] smear and/or positive culture and/or NAAT), histologic confirmation (granulomas with or without necrosis, and with or without positive AFB smears on pleural biopsy) or typical radiological features with confirmed TB in other sites. The diagnostic

yield(s) for pleural TB microbiologic investigations are summarized in Table 1.

Good practice statement

- **In patients with suspected TB pleural effusion, pleural fluid should be sent for cell count and differential, protein, glucose, pH, lactate dehydrogenase, acid-fast bacilli smear and culture, Gram stain and culture, nucleic acid amplification test and cytology.**

If pleural and sputum samples are nondiagnostic, pleural biopsy should be considered for definitive diagnosis, given higher diagnostic yield from tissue samples. Timing of pleural biopsy (waiting for final pleural/sputa mycobacterial culture results versus expedited biopsy if NAAT and smears are negative) should be determined on a case-by-case basis, based on patient's clinical scenario.

Biopsy samples should be sent for AFB smear, NAAT and culture (in saline) and for histopathology in cytolyte/formalin. Either image-guided closed or thoracoscopic pleural biopsy can be performed, dependent on local resources and expertise.

3.2.1. Pleural TB treatment

Data from multiple observational studies has demonstrated high rates of treatment completion with low relapse rate with 6 months of therapy. Medical treatment of TB pleural effusion results in successful outcomes in more than 85% of patients.^{122–125}

Based on data from a Cochrane review, adjunctive corticosteroids may accelerate the resolution of pleural fluid and reduce pleural thickening. However, there is no evidence that steroids impact lung function and steroids are associated with adverse effects.¹²⁶

Two small randomized trials have compared therapeutic drainage (thoracentesis or chest tube) and found conflicting results.^{123,127} In one study, chest tube drainage did not reduce residual pleural thickening or improve end-of-treatment forced vital capacity (FVC).¹³³ In the second study, thoracentesis was associated with less residual fluid and a higher FVC at the end of treatment, though the difference was minimal and likely not clinically significant.¹³⁷

Recommendations

- **We conditionally recommend six months of standard anti-TB therapy for treating drug susceptible pleural TB (poor evidence).**
- **We conditionally recommend against routine adjunctive corticosteroid use for pleural TB (poor evidence).**
- **We conditionally recommend against routine therapeutic thoracentesis/chest tube drainage for pleural TB-associated effusions (poor evidence).**

Good practice statement

- **In patients with significant dyspnea, therapeutic thoracentesis may be considered on a case-by-case basis to relieve symptoms.**

Fevers usually resolve within 2 weeks of treatment initiation but may take up to 2 months for full resolution in some cases.^{113,128} Paradoxical radiologic worsening may occur early after initiation of therapy in up to 26% of patients.¹¹⁸ In most cases, pleural fluid resolves after 6 weeks but may last up to 3 months.¹²⁸

TB empyema is diagnosed based on the presence of purulent fluid in a patient with TB pleural space infection.¹¹⁵ Based on expert opinion, drainage and/or decortication is recommended in addition to standard anti-TB therapy.^{113,129} There are limited data to guide the optimal composition/duration of therapy or the use of adjunctive treatments such as intrapleural thrombolytics.^{129,130} Consultation with a TB expert is recommended in this setting.

3.3. Abdominal TB

Abdominal TB can be subdivided into clinically distinct or overlapping presentations, comprising gastrointestinal disease (luminal), peritoneal, visceral, and abdominal lymph node disease. Gastrointestinal and peritoneal forms of TB disease occur most frequently and are the focus of this section. Isolated visceral and abdominal lymph node disease are uncommon forms of abdominal TB.^{131,132}

3.3.1. Gastrointestinal TB

TB may affect any part of the gastrointestinal tract. Ileocecal and jejunoileal involvement are most common (up to 75% of cases), followed by colorectal disease (majority on right side).^{131–134} Gastrointestinal TB, particularly ileocecal disease, may present with clinical and radiographic features that are indistinguishable from Crohn's disease, such as chronic abdominal pain, constitutional symptoms and a right lower quadrant mass.^{135,136} Mesenteric lymph node enlargement is more commonly found on diagnostic imaging in patients with gastrointestinal TB than in patients with inflammatory bowel disease.^{137–140}

Diagnostic testing for gastrointestinal TB includes stool studies for *M. tuberculosis* and, where available, endoscopy for biopsies with multiple dedicated samples sent for both culture/NAAT in saline and histopathology to maximize diagnostic yield. Collectively, these tests may support or confirm diagnosis of TB in up to 70% of cases (see Table 1).^{133,141–143}

If endoscopy is nondiagnostic, laparoscopy/laparotomy can be considered for definitive diagnosis. An empiric trial of anti-TB therapy may be required in some cases. However, in addition to the usual concerns regarding empiric therapy, partial short-term clinical response of Crohn's disease to anti-TB therapy is well described and may confound diagnosis.¹⁴⁴ Such delays in Crohn's disease treatment have been shown to worsen long-term outcomes (increased rates of stricture and future surgery).^{135,144,145} When empiric TB therapy is pursued, end-of-treatment endoscopy may be helpful in differentiating TB disease as significant or complete mucosal healing is reported in more than 75% of cases.^{133,143}

3.3.2. Peritoneal TB

TB involving the peritoneum presents most commonly (in more than 60% of cases) with abdominal swelling secondary to ascites, often concurrent with abdominal pain, fevers and/or weight loss.^{38,40} Individuals with chronic liver disease (particularly alcoholic liver disease), chronic renal disease and HIV are at increased risk.^{7,18,146}

Radiologic assessment can be helpful but is not diagnostic in peritoneal TB.^{137,140,147} Diagnosis of peritoneal TB typically starts with percutaneous sampling of ascites for fluid analysis, microscopy and culture. Assessment of ascitic fluid classically demonstrates a proteinaceous exudative pattern (protein greater than 30 grams per L) with a predominance of lymphocytes (greater than 70%) and a low (less than 11 grams per L) serum ascites albumin gradient (SAAG). However, when TB peritonitis complicates chronic peritoneal dialysis or decompensated cirrhosis, it may not have this typical ascites profile. Ascitic fluid is rarely AFB-smear positive but may culture *M. Tuberculosis* in up to 80% of cases. Larger volume (greater than 1 liter) and concentration of samples are reported to increase culture yield.^{38,148} If ascitic fluid sampling is nondiagnostic, peritoneal biopsy (diagnostic image-guided or laparoscopic) for AFB smear, culture, NAAT and histopathology should be considered, as diagnostic yield for peritoneal tissues is significantly higher than ascitic fluid alone (See Table 1).^{38,41,149} ADA testing is not used in Canada (see Chapter 3: Diagnosis of Tuberculosis Disease and Drug-resistant Tuberculosis).

It is important to recognize that peritoneal and other forms of TB can cause an elevation in serum tumor marker CA 125 levels. Given shared radiographic findings with peritoneal carcinomatosis of metastatic ovarian cancer, there are numerous case reports of misdiagnosis of malignancy, underscoring the importance of pursuing tissue and/or culture diagnosis of peritoneal TB.^{39,150}

3.3.3. Abdominal TB treatment

Three clinical trials evaluating treatment of abdominal TB consistently found that 6 months of standard anti-TB treatment is adequate in individuals with drug-susceptible abdominal TB and extension of treatment does not significantly improve clinical cure or relapse risk. The available evidence is limited to HIV-negative individuals and strongest for gastrointestinal forms of abdominal TB.¹⁵¹

Surgery should generally be reserved for abdominal TB cases with serious complications, such as perforation, bleeding or obstruction.¹⁵²

Recommendation

- We conditionally recommend six months of standard anti-TB therapy for all forms of drug-susceptible abdominal TB (poor evidence).

3.4. Bone and joint TB

3.4.1. Spinal/vertebral disease

Spinal or vertebral TB (Pott's disease) involvement is noted in approximately 50% of bone and joint TB cases.^{52,153–155}

Most patients present with slowly progressive back pain.^{156–157} Fever and constitutional symptoms are not common unless there is concurrent extraspinal or disseminated disease. Given nonspecific complaints, the diagnosis of spinal TB usually is not made until several months after the beginning of symptoms.^{72,73,80,154,155,156–162} Radiographic findings include loss of vertebral body height and scalloping of vertebral bodies by paraspinal fluid collections; however, these findings are insensitive.¹⁶³ CT and magnetic resonance imaging (MRI) imaging are more sensitive for detection of vertebral and soft-tissue abnormalities associated with spinal TB.¹⁶³ Findings include anterior vertebral involvement of thoracic or lumbar vertebrae adjacent to the endplate with evidence of marrow edema with minimal sclerosis; discitis of intervening disks with preservation of the disk until late in disease; and large paraspinal abscesses (calcification being suggestive of TB). MRI is the modality of choice for assessing spinal cord involvement or damage.^{80,159,160,163–168}

CT-guided needle biopsy of vertebrae and/or aspiration of paraspinal fluid collections have the highest diagnostic yield, outside of surgery, and are the recommended initial diagnostic sampling method (Table 1).^{23,72,73,77} If CT guided sampling cannot be performed or is nondiagnostic, a surgical biopsy can be obtained for definitive diagnosis and to assess for etiologies other than TB osteomyelitis.^{23,50,72,78,169–174} It is important to assess the patient for other manifestations of TB disease, as studies have demonstrated that one-third of patients with spinal TB had evidence of TB elsewhere, and the diagnosis of TB disease was made in one-quarter of patients by obtaining nonspinal specimens.⁷³

3.4.2. Joint TB (TB arthritis)

Joint TB is usually a mono-arthritis affecting large, weight-bearing joints such as the hip or knee. Symptoms can include swelling, pain and loss of function. Focal signs typically associated with septic arthritis, such as local erythema and warmth, are often missing, as are constitutional symptoms. Cartilage erosion, deformity and draining sinuses have been associated with late presentation. *M. tuberculosis* has also been associated with prosthetic joint infections. Osteomyelitis affecting other sites in the skeleton is uncommon but has been described.^{52,175–178} Multifocal presentations can occur in 15% to 20% of cases, often in immune-suppressed individuals, and can be misinterpreted as metastases or inflammatory arthritis.^{74,179}

Radiologic findings suggestive of joint TB include synovial thickening and joint effusion, however differentiation from other arthritic conditions can be difficult. MRI changes suggestive of joint TB include moderate but uniform thickening of the synovium, as compared with the larger and irregular thickening seen in rheumatoid arthritis. Adjacent fasciitis and cellulitis can be seen in both TB and pyogenic arthritis but are more indicative of a pyogenic arthritis.^{153,164,177–181}

Synovial fluid aspirate is a reasonable first step in obtaining a diagnosis of joint TB.^{79,153,178,182} Typical synovial fluid findings are that of an elevated white blood cell count (WBC) (10,000 to 20,000 WBC per ml with neutrophil predominance), decreased glucose (less than 2.2 mmol per L) and

elevated protein (greater than 25 g per L).^{79,153,157,178} The yield of synovial-fluid AFB smear is low (19%); however, NAAT and mycobacterial culture synovial-fluid sensitivity is relatively high at 75 to 85%.^{23,50,51,74,75,79,153,178,182} Synovial biopsy with sampling for NAAT, mycobacterial smear and culture, as well as histopathology, has high diagnostic yield (94%) and should be obtained if joint TB is still a diagnostic consideration and synovial fluid mycobacteriology assessment is negative (Table 1).^{23,50,51,74,75,79,153,178,182}

3.4.3. Bone and joint TB treatment

In a review of bone and joint TB treatment with isoniazid (INH) and rifampin (RMP) anti-TB therapy for individuals with drug-susceptible TB (majority of studies were observational cohort in adults and children), a risk of relapse was 1.35% with 6 months of anti-TB therapy (539 individuals studied), 0.86% with 6–12 months of anti-TB therapy (437 individuals studied) and 0.51% with greater than 12 months of anti-TB therapy (1,386 individuals studied).¹⁸³ Similar outcomes were identified in 2 small randomized controlled trials.^{184–186}

Increased risk of failure/relapse has been associated with extensive disease at the outset of treatment (radiographically defined or with evidence of smear positivity at beginning of therapy), evidence of sclerotic bony disease on imaging, and Erythrocyte sedimentation rate (ESR)/C-reactive protein (CRP) (both measures of inflammatory response) elevation at end of treatment.^{162,183} The definition of cure is difficult in bone and joint TB, and follow-up samples are infrequently obtained to demonstrate lack of mycobacterial growth. Alternative definitions of cure have utilized radiologic markers; however, vertebral x-rays may never return to baseline and studies in spinal TB have shown that 50% of patients will have MRI evidence of TB activity even at the end of 12 months of treatment.^{165,168,187} Studies utilizing vertebral CT positron emission tomography (PET) imaging have suggested decrease in PET activity may predict when TB treatment for spinal TB can be safely discontinued, but there is not enough evidence to recommend currently.¹⁸⁸ Routine surgery for bone and joint TB is not required to achieve cure, but should be considered to treat complications of vertebral TB (neurologic compromise) and joint disease (pain and immobility).^{161,185,189}

Recommendations

- We conditionally recommend 6 months of standard anti-TB therapy for individuals with drug-susceptible bone and joint TB with extension to 9–12 months in individuals with markers of increased risk of failure/relapse; diagnostic biopsy sample smear positive for acid-fast bacilli and/or elevated erythrocyte sedimentation rate/C-reactive protein (ESR/CRP) at planned end of treatment (*poor evidence*).
- We conditionally recommend against routine surgical intervention as part of treatment in spinal TB. Surgical treatment of spinal TB should be considered in those with progressive neurologic deterioration and in those less than 15 years of age with significant kyphosis (*poor evidence*).

- **We conditionally recommend against routine surgical intervention as part of treatment in joint TB. Surgical treatment should be considered to prevent extension of disease and to provide relief of pain and immobility after control of infection is established (poor evidence).**

3.5. Central nervous system (CNS) TB

CNS TB refers to the clinical and pathological spectrum of TBM, spinal TB arachnoiditis and tuberculoma. The epidemiology of CNS TB varies by regional TB prevalence. In areas of high TB prevalence, CNS TB occurs more commonly among children and young adults, whereas TBM predominates and occurs more commonly in adults as reactivation disease in low-prevalence settings. In one retrospective Canadian series, 75% of CNS TB disease was adult TBM.^{190–192}

3.5.1. TBM

TBM is the most severe and rapidly progressive form of TB, meaning suspected cases must be treated as a medical emergency. Young children (less than 5 years) and people with HIV are at greatest risk of TBM. Although classically, TBM is described as a slow progressive meningitis syndrome, 50% of people with TBM are ill for less than 2 weeks before diagnosis.¹⁹³ TBM may have a prodrome of headache, malaise, fever and personality changes, followed by meningismus, cranial nerve palsies and confusion.¹⁹⁴

Prior to the development of anti-TB therapy, TBM was universally fatal. Despite major advances in diagnosis and treatment, global mortality estimates for TBM remain high (20–40%), with up to 50% of survivors suffering permanent neurologic deficits and long-term disability.^{194–197} Unfavorable outcomes in TBM correlate with older age, immunosuppression, presence of hydrocephalus and/or vasculitis and more advanced clinical stage at time of presentation.^{193,198,199} Prognostic models have been developed for the prediction of unfavorable outcomes among adults (greater than 14 years of age) with TBM.^{198,200}

Neurologic imaging in TBM can support the diagnosis; inform prognosis; and identify a need for neurosurgical intervention. Basal meningeal enhancement on contrast-enhanced CT has high specificity for TBM (greater than 90% in adults and children), further increased by additional findings of infarcts and hydrocephalus that may become more apparent in later-stage disease.^{194,201} Gadolinium-enhanced MRI of the brain provides improved identification of early disease by visualization of localized leptomeningeal disease, characterization of ischemia and early infarction and superior evaluation of cranial nerve and brain stem involvement, compared with CT. However, both CT and MRI lack diagnostic sensitivity. Typical findings may be less prominent or absent in 15–30% of children, those with early disease, and those with advanced HIV.^{202,203}

Lumbar puncture should be performed when feasible as the preferred diagnostic test for TBM. Early in the disease,

cerebrospinal fluid (CSF) measurements are often normal. With disease progression, opening pressure becomes elevated, with low glucose levels (less than 2.5 mmol per L), elevated protein (greater than or equal to 0.5 g per L) and a moderate pleocytosis with lymphocyte/mononuclear cell predominance (100–500 total white cells per mL).²⁰⁴ Identification of *M. tuberculosis* in CSF is the standard for definite diagnosis of TBM (Table 1). However, the paucibacillary nature of TBM limits the diagnostic yield of microbial tests. Large volume sampling (7 to 15 mL, 3 serial samples), concentration of samples, experienced laboratory-technician review and the inclusion of more gene targets in NAATs can maximize diagnostic yield.^{23,205–207} Based on the insensitivity of CSF microscopy and diagnostic delay of culture-based methods, NAAT testing should be performed if adequate volume of CSF is available to assist with rapid diagnosis.^{208,209} Among commercially available NAATs, GeneXpert ULTRA has the best-available evidence supporting high sensitivity and should be used when possible.³²

3.5.2. TBM treatment

Clinical suspicion of TBM should be based upon presence of epidemiological risk factors and clinical features for TB, as well as relative suspicion of alternate diagnosis. Treatment started during early-stage TBM disease may decrease mortality to less than 10%.^{206,210–212}

Good practice statement

- **In cases with high suspicion of TB meningitis, we suggest initiating empiric therapy immediately while awaiting diagnostic results, to reduce mortality and prevent complications.**

An optimal anti-TB therapy regimen for TBM remains uncertain. Current guidelines from the American Thoracic Society/Centers for Disease Control and Prevention/Infectious Diseases Society of America, National Institute for Health and Care Excellence, American Academy of Pediatrics and World Health Organization (WHO) recommend standard dose RMP and INH with pyrazinamide (PZA) during initial two-month intensive phase, with lack of consensus on the best fourth agent.^{213–218}

Based on relatively low CNS penetration of RMP, a series of clinical trials have evaluated the clinical and pharmacokinetic outcomes of higher dose (“intensified”) RMP as part of intensive-phase regimens for TBM (in these studies high dose RMP was not given throughout full duration of treatment). Although published studies have not identified an optimal dose or conclusively demonstrated clinical benefit, there are now compelling and consistent findings for both RMP CNS-concentration response correlation and safety studies supporting use of higher-dose RMP regimens.^{219–226}

Recommendation

- **We conditionally recommend using higher doses of rifampin (greater than 15 mg/kg/day orally, up to a**

maximum dose of 35 mg/kg/day orally or 20 mg/kg/day IV) along with standard dose isoniazid, pyrazinamide and ethambutol during the intensive phase treatment for drug-susceptible TB meningitis (poor evidence).

In the absence of clinical trial data regarding the optimal fourth drug in a TBM regimen, the American Pediatric Association and the WHO advise the use of streptomycin or ethionamide in place of ethambutol (EMB), based on low CNS penetration of EMB and difficulty monitoring EMB toxicity in children.²²⁷ Although fluoroquinolones have a favorable CNS-penetration profile, clinical trials evaluating their use as a fourth agent for TBM in adults and children have not shown benefit outside of drug-resistant disease and raise concern about adverse effects.^{228–231} Second-line drugs with favorable CNS-penetration profiles (linezolid, ethionamide) lack adequate quality clinical data to support standard use at this time.^{232,233,234,235,219}

Recommendation

- **We conditionally recommend against using fluoroquinolone for TB meningitis unless there is a concern about drug resistance (poor evidence).**

No clinical trials have directly assessed duration of anti-TB treatment for CNS TB. A review of TBM observational studies did not find increased relapse among those treated for 6 months.²³⁶ However, given concerns about variable CNS drug penetration and potential for significant harm from relapsed disease if undertreated, current American, British and WHO guidelines continue to recommend a minimum of nine months and up to 12 months of anti-TB therapy for all forms of drug-susceptible CNS-TB.^{190,214,217,237,238} Based on poor evidence for extended duration of therapy, if issues with anti-TB therapy tolerance occur, then continuation beyond 6 months should be reevaluated in consultation with an expert in TB.

Recommendation

- **We conditionally recommend, given the severity of disease and risk of morbidity with inadequate treatment of central nervous system TB, extending treatment to 9–12 months for drug-susceptible disease (poor evidence).**

In addition to anti-TB therapy, measures to mitigate elevated intracranial pressure and prevent hydrocephalus and infarction are important. Adjunctive steroids have been shown to reduce short-term mortality of TBM but have not been demonstrated to reduce disabling neurologic defects or longer-term survival.^{194,197,239,240} (Table 2 Corticosteroid Dosing).

Recommendation

- **We strongly recommend that all individuals presenting with TB meningitis receive a course of corticosteroids guided by disease severity (good evidence).**

Several small controlled trials using aspirin (ASA), in addition to anti-TB therapy and corticosteroids, for TBM

in both adults and children have suggested potential benefit (reduction of new infarctions and decrease in 60-day mortality) without added harm. Studied ASA regimens ranged from 81 to 1000 mg per day for the initial 1–2 months of treatment.^{241,242} Given the preliminary nature of these findings, recommendation for the addition of ASA to TBM treatment cannot be made at this time but may change with results from pending studies in both adults and children.^{243,244}

Several large clinical studies evaluating TBM treatment are currently underway (INTENSE-TBM, HARVEST, and TBM-KIDS) and guideline recommendations will be refined as data from the trials become available.^{219–221}

3.5.3. Tuberculoma and spinal TB arachnoiditis

Both tuberculoma and spinal TB arachnoiditis may occur with TBM. Spinal disease (often clinically occult) has been reported in up to 76% of TBM cases and brain tuberculomas in approximately 10% of TBM cases, with even higher prevalence in people with HIV.^{195,242,245} When presenting in isolation, symptoms are more typically subacute and neurologic in nature relating to mass effect (headache and/or focal neurologic signs of hemiplegia, nerve root compression and spinal cord syndromes), in contrast to the more acute infectious and inflammatory presentation of TBM.^{191,212}

Contrast-enhanced CT and gadolinium-enhanced MRI can demonstrate solitary or multiple ring-enhancing lesions suggestive of tuberculoma in the brain and spinal cord. MRI offers improved visualization of spinal TB arachnoiditis and associated epidural space infection, myelitis, spondylitis and nerve root involvement.²⁰¹ When possible, a tissue/microbiologic diagnosis should be pursued prior to treatment initiation (Table 1). If empiric therapy is commenced, it is important to note that paradoxical clinical (30%) and radiographic (65%) worsening during the first 6 months of effective therapy is common and confounds assessment of treatment response. Follow-up neuroimaging should be used primarily to evaluate for alternate diagnoses and complications, and be interpreted in concert with clinical and microbiological information.²⁴⁶

3.5.4. Tuberculoma treatment

There are no controlled trials evaluating optimal management of either tuberculoma or TB arachnoiditis. Recommendations for treatment duration and adjunctive corticosteroids are extrapolated from observational and clinical trial data for TBM.²⁴⁶

Good practice statement

- **In cases of drug-susceptible tuberculoma and spinal arachnoiditis, 9–12 months of standard anti-TB therapy is suggested, given the severity of disease and risk of morbidity with inadequate treatment.**

Extrapolating from other inflammatory parenchymal CNS diseases (ie, cancer) and limited observational data,

corticosteroids may have a limited role for symptom relief in the context of tuberculomas with vasogenic edema-associated neurologic symptoms.¹⁹⁰

Recommendation

- **We conditionally recommend against universal use of adjunctive corticosteroids in cases of central nervous system tuberculoma without meningitis (poor evidence).**

The Tuberculosis Meningitis International Research Consortium has compiled a collection of peer-reviewed, open-access expert reviews to address the rapidly evolving field of CNS-TB diagnosis and management (<https://wellcomeopenresearch.org/collections/tbmeningitis>). A comprehensive review of the management of complications associated with CNS-TB has also recently been published by Donovan et al.²⁴⁷

3.6. Disseminated TB

Disseminated TB is defined as disease occurring in two or more noncontiguous organs or the isolation of *M. tuberculosis* in blood, bone marrow or liver biopsy.²¹ Miliary TB is a distinct subset of disseminated TB, one that comes with increased mortality.²⁴⁸ Hematogenous dissemination of TB in miliary disease causes formation of minute tubercles throughout multiple organs, often resulting in characteristic uniform micronodular (1-5 mm) changes on lung imaging and life-threatening systemic illness.⁹⁵

The clinical presentation of disseminated TB is often nonspecific. Fever, night sweats, anorexia, weight loss and weakness are commonly reported, while respiratory or other organ-specific symptoms occur less frequently.^{94,96,249,250} Clinical presentation is often subacute or chronic, although acute fulminant deterioration (including shock and acute respiratory distress syndrome) has been described.^{251,252} In elderly patients, disseminated TB may mimic metastatic carcinoma, being characterized by progressive wasting alone. The absence of fever and chest radiograph changes in this setting can confound TB diagnostic work-up.^{253,254} Funduscopic exam demonstrating choroidal tubercles is a specific finding and occurs in up to 20% of cases among people with HIV.^{255,256} The nonspecific and often variable presentation of disseminated TB frequently leads to a delay or lack of diagnosis and a high mortality rate.^{96,257}

Good practice statement

- **In severely ill cases, or in those with significant immunocompromising conditions, empiric anti-TB treatment should be considered to prevent morbidity and death while waiting for mycobacterial culture confirmation.**

Laboratory findings in disseminated TB are nonspecific, though hematologic abnormalities are common, and hypoalbuminemia, hypercalcemia and elevated ferritin portend

a worse prognosis.^{95,96} Between 30 and 50% of cases do not have the classic discrete micronodular or “miliary” pattern on chest radiograph.⁹⁵ High-resolution CT is more sensitive than chest radiograph, though not necessarily specific for miliary TB.²⁵⁸ Prompt examination of AFB smear, NAAT and culture of sputum and urine, along with clinical specimens from multiple sites where disease may be clinically or radiographically apparent (pleural, pericardial, peritoneal, CSF, liver, bone marrow, blood) increases the probability of a positive result and may obviate the need for more invasive tissue biopsy testing (Table 1).^{93,94,96,248} Positron emission tomography-computed tomography (PET-CT) scan may aid in the work-up of disseminated TB by accurately mapping involved lymph nodes to improve diagnostic yield of minimally invasive sampling.²⁵⁹ However, limited accessibility may make routine use of PET-CT impractical and delayed access should not delay other investigations or treatment. CNS involvement in disseminated TB may occur in 10-30% of cases, with increased prevalence among people with HIV. Thorough clinical assessment and a low threshold for neuroimaging and lumbar puncture, among those with objective neurologic findings, is advised when disseminated TB is suspected or confirmed.

3.6.1. Disseminated TB treatment

There are no clinical trials specifically evaluating optimal therapy for disseminated TB and observational studies are confounded by variable definitions. Existing guideline recommendations are largely expert opinion based on extrapolation from evidence on treatment of other forms of organ-specific EPTB.^{95,96,213,214}

Recommendation

- **For drug-susceptible disseminated TB without central nervous system involvement, we conditionally recommend treatment with six months of standard anti-TB therapy. Extension to 9-12 months can be considered in immune-compromised patients if the predisposing condition is not modified (eg, people with human immunodeficiency virus not receiving antiretroviral therapy, or continuation of immunosuppressive therapy) (poor evidence).**

3.7. Genitourinary TB

3.7.1. Urinary tract

Urinary tract disease is more commonly seen in men and those with end-stage renal disease requiring dialysis.^{56,260} Most often, onset of the disease is insidious, and patients present with asymptomatic sterile pyuria, gross hematuria, frequency and dysuria.²⁶¹⁻²⁶⁴ Back pain or flank pain resembling acute pyelonephritis often reflects calyceal or ureteral obstruction, though renal colic is uncommon. Bladder involvement (with resultant diminished bladder capacity) may present with complaints of an inability to empty the bladder and may be associated with the development of a secondary bacterial bladder infection.

Ultrasonography, CT and MRI are useful diagnostic modalities for the assessment of genitourinary TB.^{265,266} Radiologic abnormalities associated with urinary tract TB are distorted or eroded calyces, hydronephrosis, renal parenchymal scarring and calcification (all of which can mimic the changes seen in chronic pyelonephritis).^{265,267}

In patients with urinary tract disease, urine samples sent for AFB smear, NAAT and culture will confirm the diagnosis in more than 90% of cases. (Table 1 and Chapter 3: **Diagnosis of Tuberculosis Disease and Drug-resistant Tuberculosis** for details).^{23,54–57,268} Antibiotic therapy with fluoroquinolones, used to treat superimposed bacterial infection, may compromise the laboratory's ability to recover *M. tuberculosis* in urine samples and therefore should be stopped more than 48 hours before urine specimens are collected for mycobacteriologic assessment.²⁶⁸ Occasionally, fine needle aspiration (FNA) of the kidney under ultrasound guidance may be indicated if radiologic assessment is suggestive of renal TB and urine mycobacterial cultures are negative.^{57,58}

3.7.2. Genital tract

Genital tract TB may follow from a renal focus; the diagnosis of genital TB, therefore, should lead to a search for urinary tract disease. However, disease involving the female genital tract or the seminal vesicles in males is most often due to hematogenous or direct spread from neighboring organs; as such a lack of confirmation of urinary TB should not preclude further investigation in genital tract TB.

3.7.2.1 Female genital tract TB. Any site in the female genital tract may be involved; however, for reasons that are unknown, 90–100% of patients with female genital tract TB have fallopian tube infection, and both tubes are usually involved, with resultant high rates of infertility.^{269–271} Female genital tract TB is most commonly diagnosed during a work-up for infertility or during evaluation of abnormal uterine bleeding, pelvic pain or adnexal masses. Other, less common sites of involvement in the female genital tract include cervical or vulvovaginal, which frequently presents as abnormal vaginal bleeding or ulcers.²⁷² The diagnosis of female genital tract TB requires a combination of microbiologic, histologic and radiologic techniques.^{59,60,270,271,273} Findings on hysterosalpingography may suggest TB, although, as with renal TB, imaging is often nonspecific and characteristic findings are typically seen only with more advanced disease.²⁷⁴ Cultures of *M. tuberculosis* can be obtained from several sources, including menstrual fluid, peritoneal fluid, endometrial biopsy or biopsy of abnormal tissue identified during laparoscopy.^{59,60,62,273,275} Small studies have examined the role of NAAT testing in diagnosis of female genital tract TB, with high sensitivity in tissue samples.⁶¹ Even with adequate treatment for female genital tract TB, subsequent fertility rates range between 10% and 30%.^{60,276,277} The importance of confirming a diagnosis of TB-related infertility has been highlighted by cases of congenital TB in those in whom the diagnosis has not been identified prior to in-vitro fertilization.²⁷⁸

3.7.2.2 Male genital tract TB. As with the female genital tract, any site of the male genital tract can be involved. Epididymitis/orchitis is the most common presentation.²⁶⁹ Penile and prostatic involvement are rare.^{269,279,280} Male genital tract TB usually presents with scrotal swelling, sometimes with rectal or pelvic pain and less commonly with hydrocele or, in advanced cases, a discharging sinus (“watering can” perineum).^{269,279} On examination, the epididymis can be rubbery or nodular, and the prostate can be thickened with hard nodules. Between 50% and 75% of patients have palpable thickening of the vas deferens. Urine and discharge from draining sinuses should be sent for AFB smear, NAAT and culture.^{63–66} If this is nondiagnostic, biopsies (FNA or excisional) should be performed for diagnosis.^{63–66}

3.7.3. *Bacillus Calmette-Guérin* (BCG) disease

Individuals who have received intravesical administration of BCG for treatment of bladder cancer are at risk of developing localized genitourinary TB (1% of patients) or spinal or disseminated disease (0.4% of patients).^{281–287} Men are more likely to develop BCG disease, with a median time to diagnosis of 170 days post-installation.²⁸⁴ Diagnosis can be made with urine mycobacterial sampling for genitourinary disease, and with biopsy sampling of affected organ(s) for disseminated disease.

3.7.4. Genitourinary TB treatment

Standard 6-month anti-TB treatment for drug-susceptible genitourinary TB has demonstrated adequate mycobacterial cure rates in observational studies in male genital TB and urinary TB, and is the suggested regime.^{21,288–290} There has been one randomized controlled trial in female genital TB comparing 6 months and 9 months of treatment that demonstrated similar treatment outcomes; therefore 6 months is the suggested treatment duration.²⁹¹ Given the drug-susceptibility profile of BCG mycobacterium (with sensitivity to INH and RMP and resistance to pyrazinamide), individuals with disease caused by BCG are usually treated with 9 months of INH and RMP.^{281,282,285–287}

Recommendations

- **We conditionally recommend six months of standard anti-TB therapy for treatment of drug-susceptible genitourinary TB (poor evidence).**
- **We conditionally recommend a minimum of nine months of treatment with isoniazid and rifampin for BCG disease, given inherent pyrazinamide resistance (poor evidence).**

3.8. TB pericarditis

Common presenting symptoms are nonspecific and result from the underlying infectious process (fever, night sweats),

cardiac compromise (dyspnea, orthopnea) or disease elsewhere (eg, cough). Physical signs vary depending upon the degree of cardiac compromise.^{86,292} Early presentation is associated with a serosanguinous exudative pericardial effusion that may resolve spontaneously over a few weeks or may progress to cardiac tamponade or pericardial constriction.

Imaging modalities for TB pericarditis can include chest radiography, echocardiography, cardiac MRI or CT chest imaging.^{86,292} TB pericardial effusion is more likely than viral/idiopathic pericardial effusion if there is mediastinal lymphadenopathy on CT imaging; however, this does not assist with differentiation from malignant pericardial effusions.^{293,294}

TB pericardial fluid is typically a bloody exudative effusion and often lymphocyte-predominant.⁸⁴ However, similar findings can be found in chronic idiopathic and malignant pericardial effusions.²⁹⁵

Diagnosis of TB pericarditis can be confirmed with sampling of pericardial fluid (60% sensitivity) and/or pericardial tissue (90% sensitivity) for AFB smear, NAAT, culture and histopathologic analysis (Table 1). Given the difficulties in diagnosis and the high morbidity and mortality associated with this condition (80-90% mortality in the pre-antibiotic era), empiric treatment may need to be considered while awaiting the results of microbiologic/histologic testing (especially in the immunocompromised, as typical histopathology findings may not be present).²⁹⁶⁻²⁹⁸

3.8.1. TB pericarditis treatment

Treatment regimen/duration recommendations are based on observational data that has demonstrated a reduction in incidence of constrictive pericarditis and mortality compared to the pre-antibiotic era.^{82,84,296,299-301}

In a Cochrane review that analyzed HIV-negative and people with HIV separately, corticosteroids significantly reduced the risk of death from pericarditis. In people with HIV, steroids did not significantly improve any clinical outcome. Of note, only 20% of the participants with HIV were taking antiretroviral medications. There was not a significant increase in opportunistic infections or malignancy among either People with HIV or HIV-negative participants, although the data are limited. There are minimal data for people with HIV on established antiretroviral treatment. In patients with good antiretroviral drug viral suppression, the data from HIV negative patients may be considered more applicable.²⁹⁶ An initial dose of prednisolone 120 mg PO daily with a taper over six weeks was prescribed in the largest and most recent study included in the 2017 Cochrane meta-analysis (Table 2).^{296,301}

Recommendations

- **We conditionally recommend six months of standard anti-TB therapy for treatment of drug-susceptible TB pericarditis (poor evidence).**
- **We conditionally recommend initial adjunctive corticosteroid treatment in all HIV negative patients with TB pericarditis (poor evidence).**

- **We conditionally recommend against routine use of adjunctive corticosteroids in people with HIV not on antiretroviral treatment (poor evidence).**

Good practice statements

- **We are unable to provide a specific recommendation on use of adjunctive corticosteroids in people with HIV who are on antiretroviral treatment with TB pericarditis given lack of data. We suggest assessing on a case-by-case basis pending further study.**
- **In patients with recurrent pericardial effusions or persistently elevated central venous pressures despite removal of pericardial fluid and use of anti-TB drugs, we suggest referral for consideration of early pericardiectomy.**

3.9. Ocular TB

Ocular involvement is a rare manifestation of EPTB. Although commonly encountered in the context of disseminated disease, ocular TB more typically occurs without clinically apparent systemic disease.^{255,302} In low TB-incidence settings such as Canada, ophthalmologist evaluation of uveitis of unknown etiology with positive immune markers for TB infection (identified through a tuberculin skin test (TST) or interferon-gamma release assay (IGRA)) often leads to a TB center referral for consideration of empiric anti-TB therapy.^{303,304}

TB can affect all parts of the eye and may result from hematogenous spread or adjacent structure extension. Ocular TB can be subcategorized by involvement of the peri-ocular, superficial and intra-ocular structures of the eye, which may occur in isolation or overlap. Intra-ocular TB (IOTB) is the most common form of ocular TB and predominantly manifests as uveitis, primarily involving the vascular posterior portion of the eye (choroid). IOTB may occur in one or both eyes, with variable symptoms ranging from blurred vision to ophthalmalgia, conjunctivitis and vision loss.³⁰² The pathophysiology of IOTB remains incompletely characterized but is posited to involve direct infection of the eye or immune-mediated hypersensitivity reaction triggered by TB infection elsewhere in the body.³⁰⁵

The clinical overlap between IOTB and other, more common infectious and noninfectious causes of uveitis pose a significant diagnostic challenge. Whereas ophthalmic exam findings of choroidal granulomas, occlusive retinal vasculitis and multifocal serpiginoid choroiditis have been proposed as specific to IOTB in high TB-incidence settings, the transferability of these findings to low-incidence setting has been questioned.^{302,306} Confirmation of IOTB (culture, molecular or pathologic) is rarely achieved, given the risks associated with ocular sampling and generally low diagnostic yield of such samples when obtained.^{307,308} Therefore, a presumptive diagnosis of IOTB is often made based upon ophthalmic exam findings, epidemiologic risk factors for TB exposure and

positive immune markers of TB infection (TST and/or IGRA) in the absence of alternate cause.^{309,310} This approach will over-estimate those with IOTB compared with those diagnosed based on polymerase chain reaction (PCR) ocular sampling (46.9% vs 37.7%) or TB culture confirmation at another site (3.8%).^{310–313}

3.9.1. Ocular TB treatment

Consensus guidance has recently been published summarizing clinical evaluation and treatment of ocular TB.³¹¹ Given the treatable nature of ocular TB and risk of vision loss with treatment delay, initiation of anti-TB therapy should not be delayed once adequate work-up is complete.^{304,312} Given that IOTB diagnosis is typically indirect and highly dependent on eye exam findings, clear communication with the TB care provider regarding ophthalmic findings and degree of clinical suspicion for IOTB is essential to ensuring timely and appropriate management.

While 9 or more months of therapy has been proposed to reduce relapse, particularly with evidence of persistent inflammatory changes, intra-ocular inflammatory response to effective therapy may be delayed and ongoing infection indistinguishable from hypersensitivity-associated inflammation.^{303,304,310,312–317}

The role of ocular or systemic anti-inflammatories remains controversial and, when used, should be under the guidance and follow-up of an ophthalmologist.^{304,318} TB preventative therapy should only be used when alternate etiology of ocular disease has been confirmed and is otherwise indicated.³¹¹ EMB can be safely used as part of standard anti-TB therapy for intra-ocular TB, assuming close follow-up to monitor for signs of optic neuropathy, particularly in those with risk factors.^{311,319}

Recommendation

- We conditionally recommend 6 months of standard anti-TB therapy for drug-susceptible (suspected or confirmed) intra-ocular TB (*poor evidence*).

3.10. Dermatologic and other rare manifestations of extra-pulmonary TB

TB involving skin is estimated to occur in 1%-2% of EPTB cases. This may be from direct infection (cutaneous TB) or immune-mediated reactions to TB infection elsewhere in the body (tuberculids).³²⁰ Cutaneous TB typically results from endogenous spread (direct contiguous or lympho-hematogenous) and rarely from exogenous inoculation (TB chancre or verrucosa cutis).³²¹ More common forms of cutaneous TB include scrofuladerma (ulcerative erosions), lupus vulgaris (patches or plaques) and orificialis disease (anal and perianal localizing). Tuberculids are often papulo-nodular in character, ranging from superficial and minimally symptomatic (lichen scrofulosorum) to deeper, painful and, at times, ulcerating panniculitis (erythema induratum of Bazin and papulonecrotic disease).^{320–322}

Diagnosis of cutaneous TB is made by biopsy, with histopathologic assessment and mycobacterial smear and culture. Non-TB mycobacteria and leprosy may also manifest as skin disease, underscoring the diagnostic importance of appropriate tissue-sample testing.

Tubercloid diagnosis can be challenging, often entailing a combination of consistent biopsy histopathology, exclusion of alternate diagnoses, demonstration of TB infection/disease (culture confirmed at another site or supported by chest x-ray, TST and/or IGRA) and response to TB treatment.

Recommendation

- We conditionally recommend 6 months of standard anti-TB therapy for treatment of drug-susceptible TB of the skin, including tuberculids (*poor evidence*).

In addition to more common extra-pulmonary sites previously described, TB may also involve non-nodal glandular tissue (eg, breast), great vessels, endocardium and bone marrow.^{321,323–326} Although rare, life-threatening immune-mediated responses to *M. tuberculosis* may also occur more frequently with EPTB. These include, but are not limited to, hemophagocytic lymphohistiocytosis, acute

Table 2. Examples of published corticosteroid regimens for extra-pulmonary TB.^a

| | TBM | | | | Pericardial TB Adults ⁽³⁰¹⁾ (≥18 years) |
|-----------------|---|--|--|---|---|
| | Children ^(329,330) (≤14 years) | | Adults/Adolescents ⁽¹⁹⁶⁾ (>14 years) | | |
| Clinical Status | All | Alert (GCS 15) | Altered LOC (GCS <15) +/- focal neurologic deficits (8 wk course) | Alert (GCS 15), no focal neurologic deficits (6 wk course) | HIV-negative |
| Week 1 | DEXA 0.6mg/kg/d IV | Prednisolone ^a 4 mg/kg/d PO (max 60 mg/d) | DEXA 0.4 mg/kg/d IV | DEXA 0.3 mg/kg/d IV | Prednisolone PO 120 mg PO |
| Week 2 | " | " | 0.3 mg/kg/d IV | 0.2 mg/kg/d IV | 90 mg PO |
| Week 3 | " | " | 0.2 mg/kg/d IV | 0.1 mg/d PO ^b | 60 mg PO |
| Week 4 | " | " | 0.1 mg/kg/d IV | 3 mg/d PO ^b | 30 mg PO |
| Week 5+ | Tapered through Week 8 | Tapered through Week 8 | 4 mg/d PO ^b decrease by 1 mg/wk | 2 mg/d PO ^b decrease by 1 mg/wk | Week 5 = 15 mg PO Week 6 = 5 mg PO |

Abbreviations: TB, tuberculosis; TBM, tuberculosis meningitis; LOC, level of consciousness; GCS, Glasgow Coma Scale; DEXA, dexamethasone; IV, intravenous; PO, by mouth; BMRC, modified British Medical Research Council TBM severity grading; GI, gastrointestinal.¹⁹⁹

^aAll corticosteroid doses assume concurrent use of rifampin. If rifampin is not part of TB regimen, then dose adjustment advised.

^bcontinue as IV if significant GI intolerance or compromise, early switch to PO may be considered in those with prompt improvement and GI tolerance.

Table 3. Summary of evidence for duration of treatment in extra-pulmonary TB with adjunct measures.

| Site | Duration TB treatment | Level of guidance | Evidence ^a | | | Adjunct measures | | |
|------------------------|---|---------------------------|-----------------------|---|--------------|--|---|---|
| | | | Type of Studies | Number | | Consistency | Corticosteroids (ref) | Other Measures (ref) |
| | | | | Studies (refs) | Participants | | | |
| Pleural | 6 months | Conditional | OBS | 4 ⁽¹²²⁻¹²⁵⁾ | 454 | Good | Do not use ⁽¹²⁶⁾ | Drainage not necessary ^(123,127) |
| Lymph Node | 6 months | Conditional | RCT | 2 ^(105,106) | 290 | Good | ^b | Surgery not necessary ⁽⁹⁸⁾ |
| Abdominal (all forms) | 6 months | Conditional | SRMA | 3 ⁽¹⁵¹⁾ | 328 | Good | | |
| Bone and Joint | 6 months, consider 9-12 months with markers of severe disease | Conditional | RCT/OBS/Review | Review 77 studies ⁽¹⁸³⁾ 2 RCTs ^(184,186) | 2,889 293 | Good, two studies with higher relapse rates Dutt et al. ⁽³³¹⁾ twice weekly self-administered therapy for 9 months and Ramachandran et al. ⁽³³²⁾ primarily based on lack of radiograph response | | Surgical intervention is not routinely recommended as part of treatment in spinal tuberculosis. Surgical treatment of spinal TB should be considered in those with neurologic deterioration and in those less than 15 years of age with significant kyphosis ^(161,185,189) |
| Central Nervous System | TBM 9-12 months | Conditional | OBS/SR | 18 ⁽²³⁶⁾ | 2,098 | Good | Use for all individuals, dosing guided by age group and disease severity. ⁽²³⁹⁾ See Table 2 for dosing guidance | Higher dose rifampin (IV or oral up to maximum dose 35 mg/kg/day) advised during intensive phase of therapy. ^(223-225,229,333,334) Routine use fluoroquinolone not advised unless concern about drug resistant TBM ^(228,231) |
| Disseminated | 6 months | Conditional | Expert opinion | | | | | |
| Genitourinary | 6 months | urinary female genital | OBS RCT | 2 ^(289,335) 1 ⁽²⁹¹⁾ | 200 175 | Good N/A – one study | Do not use unless clinical significant mass effect ^(28,246) | |
| Pericardial | 6 months | Conditional | OBS | 4 ^(82,299-301) | 660 | Good | Use for HIV negative populations. People with HIV not on ARV-do not use. ⁽²⁹⁶⁾ Limited data for People with HIV on ARV | |
| Ocular | 6 months | Conditional | OBS | 4 ^(303,314,316,317) | 426 | Poor | | |
| Cutaneous | 6 months | Conditional | OBS | Case reports and reviews | | | | |

Abbreviations: ARV, anti-retrovirals; TB, tuberculosis; OBS, observational study; RCT, randomized clinical trial; SR, systematic review; SRMA, systematic review meta-analysis.

^a(-) Insufficient data.^bOne cohort in SR was unpublished.

respiratory distress syndrome and severe hypersensitivity reaction to BCG therapy.^{251,281,327,328}

Disclosure statement

The CTS TB Standards editors and authors declared potential conflicts of interest at the time of appointment and these were updated throughout the process in accordance with the CTS Conflict of Interest Disclosure Policy. Individual member conflict of interest statements are posted on the CTS website.

Funding

The 8th edition Canadian Tuberculosis Standards are jointly funded by the Canadian Thoracic Society (CTS) and the Public Health Agency of Canada, edited by the CTS and published by the CTS in collaboration with AMMI Canada. However, it is important to note that the clinical recommendations in the Standards are those of the CTS. The CTS TB Standards editors and authors are accountable to the CTS CRGC and the CTS Board of Directors. The CTS TB Standards editors and authors are functionally and editorially independent from any funding sources and did not receive any direct funding from external sources.

The CTS receives unrestricted grants which are combined into a central operating account to facilitate the knowledge translation activities of the CTS Assemblies and its guideline and standards panels. No corporate funders played any role in the collection, review, analysis or interpretation of the scientific literature or in any decisions regarding the recommendations presented in this document.

References

- World Health Organization. *Definitions and Reporting Framework for Tuberculosis—2013 Revision: updated December 2014 and January 2020*. 2013. <https://apps.who.int/iris/handle/10665/79199>. Accessed March 2, 2022.
- Public Health Agency of Canada. personal communication. 2020.
- Vachon J, Gallant V, Siu W. Tuberculosis in Canada, 2016. *Can Commun Dis Rep*. 2018;44(3-4):75–81. doi:10.14745/ccdr.v44i34a01.
- Aguado JM, Herrero JA, Gavalda J, et al. Clinical presentation and outcome of tuberculosis in kidney, liver, and heart transplant recipients in Spain. Spanish Transplantation Infection Study Group, GESITRA. *Transplantation*. 1997;63(9):1278–1286. doi:10.1097/00007890-199705150-00015.
- Russo RL, Duley FL, Suganuma L, França IL, Yasuda MA, Costa SF. Tuberculosis in hematopoietic stem cell transplant patients: case report and review of the literature. *Int J Infect Dis*. 2010;14 Suppl 3:e187–e191. doi:10.1016/j.ijid.2009.08.001.
- Dobler CC, Cheung K, Nguyen J, Martin A. Risk of tuberculosis in patients with solid cancers and haematological malignancies: a systematic review and meta-analysis. *Eur Respir J*. 2017;50(2):1700157. doi:10.1183/13993003.00157-2017.
- Kingkaew N, Sangtong B, Amnuaiophon W, et al. HIV-associated extrapulmonary tuberculosis in Thailand: epidemiology and risk factors for death. *Int J Infect Dis*. 2009;13(6):722–729. doi:10.1016/j.ijid.2008.11.013.
- Jones BE, Young SM, Antoniskis D, Davidson PT, Kramer F, Barnes PF. Relationship of the manifestations of tuberculosis to CD4 cell counts in patients with human immunodeficiency virus infection. *Am Rev Respir Dis*. 1993;148(5):1292–1297. doi:10.1164/ajrccm/148.5.1292.
- Leeds IL, Magee MJ, Kurbatova EV, et al. Site of Extrapulmonary Tuberculosis is Associated with HIV Infection. *Clinical Infectious Diseases*. 2012;55(1):75–81. doi:10.1093/cid/cis303.
- Lau SK, Wei WI, Hsu C, Engzell UC. Efficacy of fine needle aspiration cytology in the diagnosis of tuberculous cervical lymphadenopathy. *J Laryngol Otol*. 1990;104(1):24–27. doi:10.1017/s0022215100111697.
- Asimacopoulos EP, Berry M, Garfield B, et al. The diagnostic efficacy of fine-needle aspiration using cytology and culture in tuberculous lymphadenitis. *Int J Tuberc Lung Dis*. 2010;14(1):93–98.
- Polesky A, Grove W, Bhatia G. Peripheral tuberculous lymphadenitis: epidemiology, diagnosis, treatment, and outcome. *Medicine (Baltimore)*. 2005;84(6):350–362. doi:10.1097/01.md.0000189090.52626.7a.
- Menzies D, Wong T, eds. Canadian Tuberculosis Standards. 7th ed.. Chapter 7: Nonrespiratory Tuberculosis. Canadian Thoracic Society, Canadian Lung Association, Public Health Agency of Canada. *Can Respir J*. 2013;20(Suppl A):54A–64A.
- Kohli M, Schiller I, Dendukuri N, et al. Xpert MTB/RIF Ultra and Xpert MTB/RIF assays for extrapulmonary tuberculosis and rifampicin resistance in adults. *Cochrane Database Syst Rev*. 2021;1:CD012768. doi:10.1002/14651858.CD012768.pub3.
- Conde MB, Loivos AC, Rezende VM, et al. Yield of sputum induction in the diagnosis of pleural tuberculosis. *Am J Respir Crit Care Med*. 1 2003;167(5):723–5. doi:10.1164/rccm.2111019.
- Ruan SY, Chuang YC, Wang JY, et al. Revisiting tuberculosis pleurisy: pleural fluid characteristics and diagnostic yield of mycobacterial culture in an endemic area. *Thorax*. Sep 2012;67(9):822–827. doi:10.1136/thoraxjnl-2011-201363.
- Seibert AF, Haynes J, Jr, Middleton R, Bass JB, Jr. Tuberculous pleural effusion. Twenty-year experience. *Chest*. Apr. 1991;99(4):883–886. doi:10.1378/chest.99.4.883.
- Akhter N, Sumalani KK, Chawla D, Rizvi A. N. Comparison between the diagnostic accuracy of Xpert MTB/Rif assay and culture for pleural tuberculosis using tissue biopsy. *ERJ Open Res*. 2019;5(3):00065–2019. doi:10.1183/23120541.00065-2019.
- Diacon AH, Van de Wal BW, Wyser C, et al. Diagnostic tools in tuberculous pleurisy: a direct comparative study. *Eur Respir J*. 2003;22(4):589–591. doi:10.1183/09031936.03.00017103a.
- Casalini AG, Mori PA, Majori M, et al. Pleural tuberculosis: medical thoracoscopy greatly increases the diagnostic accuracy. *ERJ Open Res*. 2018;4(1):00046–2017. doi:10.1183/23120541.00046-2017.
- Lewinsohn DM, Leonard MK, LoBue PA, et al. Official American Thoracic Society/Infectious Diseases Society of America/Centers for Disease Control and Prevention Clinical Practice Guidelines: Diagnosis of Tuberculosis in Adults and Children. *Clin Infect Dis*. 2017;64(2):e1–e33. doi:10.1093/cid/ciw694.
- Chan CH, Arnold M, Chan CY, Mak TW, Hoheisel GB. Clinical and pathological features of tuberculous pleural effusion and its long-term consequences. *Respiration*. 1991;58(3-4):171–175. doi:10.1159/000195921.
- Kohli M, Schiller I, Dendukuri N, et al. Xpert((R)) MTB/RIF assay for extrapulmonary tuberculosis and rifampicin resistance. *Cochrane Database Syst Rev*. 2018;8:CD012768. doi:10.1002/14651858.CD012768.pub2.
- Jiang J, Yang J, Shi Y, et al. Head-to-head comparison of the diagnostic accuracy of Xpert MTB/RIF and Xpert MTB/RIF Ultra for tuberculosis: a meta-analysis. *Infect Dis (Lond)*. 2020; 52(11):763–775. doi:10.1080/23744235.2020.1788222.
- Armand S, Vanhuls P, Delcroix G, Courcol R, Lemaitre N. Comparison of the Xpert MTB/RIF test with an IS6110-TaqMan real-time PCR assay for direct detection of Mycobacterium tuberculosis in respiratory and nonrespiratory specimens. *J Clin Microbiol*. 2011;49(5):1772–1776. doi:10.1128/jcm.02157-10.
- Lang AM, Feris-Iglesias J, Pena C, et al. Clinical evaluation of the Gen-Probe Amplified Direct Test for detection of

- Mycobacterium tuberculosis complex organisms in cerebrospinal fluid. *J Clin Microbiol*. 1998;36(8):2191–2194. doi:10.1128/JCM.36.8.2191-2194.1998.
27. Pai M, Flores LL, Pai N, Hubbard A, Riley LW, Colford JM. Diagnostic accuracy of nucleic acid amplification tests for tuberculous meningitis: a systematic review and meta-analysis. *Lancet Infect Dis*. 2003;3(10):633–643. doi:10.1016/s1473-3099(03)00772-2.
 28. Thwaites G, Fisher M, Hemingway C, et al. British Infection Society guidelines for the diagnosis and treatment of tuberculosis of the central nervous system in adults and children. *J Infect*. 2009;59(3):167–187. doi:10.1016/j.jinf.2009.06.011.
 29. Bouchama A, Al-Kawi MZ, Kanaan I, et al. Brain biopsy in tuberculoma: the risks and benefits. *Neurosurgery*. 1991;28(3):405–409.
 30. Rajshekhar V, Chandy MJ. CT-guided stereotactic surgery in the management of intracranial tuberculomas. *Br J Neurosurg*. 1993;7(6):665–671. doi:10.1019/02688699308995096.
 31. Mohanty A, Santosh V, Anandh B, et al. Diagnostic efficacy of stereotactic biopsies in intracranial tuberculomas. *Surg Neurol*. 1999;52(3):252–257. discussion 257–8. doi:10.1016/s0090-3019(99)00089-0.
 32. Bahr NC, Nuwagira E, Evans EE, ASTRO-CM Trial Team, et al. Diagnostic accuracy of Xpert MTB/RIF Ultra for tuberculous meningitis in HIV-infected adults: a prospective cohort study. *Lancet Infect Dis*. 2018;18(1):68–75. doi:10.1016/s1473-3099(17)30474-7.
 33. Erdem H, Ozturk-Engin D, Elaldi N, et al. The microbiological diagnosis of tuberculosis meningitis: results of Haydarpasa-1 study. *Clin Microbiol Infect*. 2014;20(10):O600–8. doi:10.1111/1469-0691.12478.
 34. Heemskerck AD, Donovan J, Thu DDA, et al. Improving the microbiological diagnosis of tuberculous meningitis: A prospective, international, multicentre comparison of conventional and modified Ziehl-Neelsen stain, GeneXpert, and culture of cerebrospinal fluid. *J Infect*. 2018;77(6):509–515. doi:10.1016/j.jinf.2018.09.003.
 35. Maiti S, Banerjee M, Das R. Down to the ground zero: Perspective from tissue diagnosis in neurological TB. *International Journal of Clinical and Biomedical Research*. 2020;6(2):25–29. doi:10.31878/ijcbr.2020.62.07.
 36. Giouleme O, Paschos P, Katsaros M, et al. Intestinal tuberculosis: a diagnostic challenge-case report and review of the literature. *Eur J Gastroenterol Hepatol*. 2011;23(11):1074–1077. doi:10.1097/MEG.0b013e32834a9470.
 37. Singh MM, Bhargava AN, Jain KP. Tuberculous peritonitis. An evaluation of pathogenetic mechanisms, diagnostic procedures and therapeutic measures. *N Engl J Med*. 1969;281(20):1091–1094. doi:10.1056/nejm196911132812003.
 38. Sanai FM, Bzeizi KI. Systematic review: tuberculosis peritonitis-presenting features, diagnostic strategies and treatment. *Aliment Pharmacol Ther*. 2005;22(8):685–700. doi:10.1111/j.1365-2036.2005.02645.x.
 39. Chau TN, Leung VK, Wong S, et al. Diagnostic challenges of tuberculosis peritonitis in patients with and without end-stage renal failure. *Clin Infect Dis*. 2007;45(12):e141–e146. doi:10.1086/523727.
 40. Yeh HF, Chiu TF, Chen JC, Ng CJ. Tuberculous peritonitis: analysis of 211 cases in Taiwan. *Dig Liver Dis*. 2012;44(2):111–117. doi:10.1016/j.dld.2011.08.020.
 41. Que Y, Wang X, Liu Y, Li P, Ou G, Zhao W. Ultrasound-guided biopsy of greater omentum: an effective method to trace the origin of unclear ascites. *Eur J Radiol*. May 2009;70(2):331–335. doi:10.1016/j.ejrad.2008.01.036.
 42. Vardareli E, Kebapci M, Saricam T, Pasaoglu O, Acikalin M. Tuberculous peritonitis of the wet ascitic type: clinical features and diagnostic value of image-guided peritoneal biopsy. *Dig Liver Dis*. 2004;36(3):199–204. doi:10.1016/j.dld.2003.10.016.
 43. Chow KM, Chow VC, Szeto CC. Indication for peritoneal biopsy in tuberculous peritonitis. *Am J Surg*. 2003;185(6):567–573. doi:10.1016/s0002-9610(03)00079-5.
 44. Kumar S, Bopanna S, Kedia S, et al. Evaluation of Xpert MTB/RIF assay performance in the diagnosis of abdominal tuberculosis. *Intest Res*. 2017;15(2):187–194. doi:10.5217/ir.2017.15.2.187.
 45. Talib A, Bhatta S, Mehmood K, et al. GeneXpert in stool: Diagnostic yield in Intestinal Tuberculosis. *J Clin Tuberc Other Mycobact Dis*. 2019;17:100131 doi:10.1016/j.jctube.2019.100131.
 46. Liu R, Li J, Tan Y, et al. Multicenter evaluation of the acid-fast bacillus smear, mycobacterial culture, Xpert MTB/RIF assay, and adenosine deaminase for the diagnosis of tuberculosis peritonitis in China. *Int J Infect Dis*. 2020;90:119–124. doi:10.1016/j.ijid.2019.10.036.
 47. Bellam BL, Mandavdhare HS, Sharma K, et al. Utility of tissue Xpert-Mtb/Rif for the diagnosis of intestinal tuberculosis in patients with ileocolonic ulcers. *Ther Adv Infect Dis*. 2019;6:2049936119863939 doi:10.1177/2049936119863939.
 48. Bera C, Michael JS, Burad D, et al. Tissue Xpert MTB/Rif assay is of limited use in diagnosing peritoneal tuberculosis in patients with exudative ascites. *Indian J Gastroenterol*. 2015;34(5):395–398. doi:10.1007/s12664-015-0599-7.
 49. Horvath KD, Whelan RL. Intestinal tuberculosis: return of an old disease. *Am J Gastroenterol*. 1998;93(5):692–696. doi:10.1111/j.1572-0241.1998.207_a.x.
 50. Penz E, Boffa J, Roberts DJ, et al. Diagnostic accuracy of the Xpert(R) MTB/RIF assay for extra-pulmonary tuberculosis: a meta-analysis. *Int j Tuberc Lung Dis*. 2015;19(3):278–284. i-iii. doi:10.5588/ijtld.14.0262.
 51. Denkinger CM, Schumacher SG, Boehme CC, Dendukuri N, Pai M, Steingart KR. Xpert MTB/RIF assay for the diagnosis of extrapulmonary tuberculosis: a systematic review and meta-analysis. *Eur Respir J*. 2014;44(2):435–446. doi:10.1183/09031936.00007814.
 52. Iseman M. *A Clinician's Guide to Tuberculosis*. Philadelphia, PA: Lippincott Williams & Wilkins; 2000.
 53. Zeka AN, Tasbakan S, Cavusoglu C. Evaluation of the GeneXpert MTB/RIF assay for rapid diagnosis of tuberculosis and detection of rifampin resistance in pulmonary and extrapulmonary specimens. *J Clin Microbiol*. 2011;49(12):4138–4141. doi:10.1128/jcm.05434-11.
 54. Bentz RR, Dimcheff DG, Nemiroff MJ, Tsang A, Weg JG. The incidence of urine cultures positive for Mycobacterium tuberculosis in a general tuberculosis patient population. *Am Rev Respir Dis*. 1975;111(5):647–650. doi:10.1164/arrd.1975.111.5.647.
 55. Lattimer JK, Reilly RJ, Segawa A. The significance of the isolated positive urine culture in genitourinary tuberculosis. *J Urol*. 1969;102(5):610–613. doi:10.1016/s0022-5347(17)62211-0.
 56. Hsu HL, Lai CC, Yu MC, et al. Clinical and microbiological characteristics of urine culture-confirmed genitourinary tuberculosis at medical centers in Taiwan from 1995 to 2007. *Eur J Clin Microbiol Infect Dis*. 2011;30(3):319–326. doi:10.1007/s10096-010-1083-z.
 57. Baniel J, Manning A, Leiman G. Fine needle cytodiagnosis of renal tuberculosis. *J Urol*. 1991;146(3):689–691. doi:10.1016/S0022-5347(17)37894-1.
 58. Das KM, Vaidyanathan S, Rajwanshi A, Indudhara R. Renal tuberculosis: diagnosis with sonographically guided aspiration cytology. *AJR Am J Roentgenol*. 1992;158(3):571–573. doi:10.2214/ajr.158.3.1738997.
 59. Turkmen IC, Bassullu N, Comunoglu C, et al. Female genital system tuberculosis: a retrospective clinicopathological study of 1,548 cases in Turkish women. *Arch Gynecol Obstet*. 2012;286(2):379–384. doi:10.1007/s00404-012-2281-y.
 60. Margolis K, Wranz PA, Kruger TF, Joubert JJ, Odendaal HJ. Genital tuberculosis at Tygerberg Hospital-prevalence, clinical presentation and diagnosis. *S Afr Med J*. 1992;81(1):12–15.
 61. Sharma JB, Kriplani A, Dharmendra S, Chaubey J, Kumar S, Sharma SK. Role of Gene Xpert in diagnosis of female genital tuberculosis: a preliminary report. *Eur J Obstet Gynecol Reprod Biol*. 2016;207:237–238. doi:10.1016/j.ejogrb.2016.10.045.
 62. Thangappah RB, Paramasivan CN, Narayanan S. Evaluating PCR, culture & histopathology in the diagnosis of female genital tuberculosis. *Indian J Med Res*. 2011;134:40–46.

63. Gómez García I, Gómez Mampaso E, Burgos Revilla J, et al. Tuberculous orchiepididymitis during 1978-2003 period: review of 34 cases and role of 16S rRNA amplification. *Urology*. 2010;76(4):776-781. doi:10.1016/j.urology.2010.01.033.
64. Lee IK, Yang WC, Liu JW. Scrotal tuberculosis in adult patients: a 10-year clinical experience. *Am J Trop Med Hyg*. 2007;77(4):714-718.
65. Gorse GJ, Belshe RB. Male genital tuberculosis: a review of the literature with instructive case reports. *Rev Infect Dis*. 1985;7(4):511-524. Jul-Augdoi:10.1093/clinids/7.4.511.
66. Madeb R, Marshall J, Nativ O, Erturk E. Epididymal tuberculosis: case report and review of the literature. *Urology*. 2005;65(4):798. doi:10.1016/j.urology.2004.10.029.
67. Altez-Fernandez C, Ortiz V, Mirzazadeh M, Zegarra L, Seas C, Ugarte-Gil C. Diagnostic accuracy of nucleic acid amplification tests (NAATs) in urine for genitourinary tuberculosis: a systematic review and meta-analysis. *BMC Infect Dis*. 2017;17(1):390. doi:10.1186/s12879-017-2476-8.
68. Chen Y, Wu P, Fu L, Liu YH, Zhang Y, Zhao Y. Multicentre evaluation of Xpert MTB/RIF assay in detecting urinary tract tuberculosis with urine samples. *Sci Rep*. 2019;9(1):11053. doi:10.1038/s41598-019-47358-3.
69. Perez-Risco D, Rodriguez-Temporal D, Valledor-Sanchez I, Alcaide F. Evaluation of the Xpert MTB/RIF Ultra Assay for Direct Detection of Mycobacterium tuberculosis Complex in Smear-Negative Extrapulmonary Samples. *J Clin Microbiol*. 2018;56(9). doi:10.1128/JCM.00659-18.
70. Kashyap B, Kaur T, Jhamb R, Kaur I. Evaluating the utility of menstrual blood versus endometrial biopsy as a clinical sample in the diagnosis of female genital tuberculosis. *Asian J Med Res*. 2012;1:45-49.
71. Lenk S, Schroeder J. Genitourinary tuberculosis. *Curr Opin Urol*. 2001;11(1):93-98. doi:10.1097/00042307-200101000-00014.
72. Colmenero JD, Ruiz-Mesa JD, Sanjuan-Jimenez R, Sobrino B, Morata P. Establishing the diagnosis of tuberculous vertebral osteomyelitis. *Eur Spine J*. 2013;22 Suppl 4:579-86. doi:10.1007/s00586-012-2348-2.
73. Cormican L, Hammal R, Messenger J, Milburn HJ. Current difficulties in the diagnosis and management of spinal tuberculosis. *Postgrad Med J*. 2006;82(963):46-51. doi:10.1136/pgmj.2005.032862.
74. Garrido G, Gomez-Reino JJ, Fernandez-Dapica P, Palenque E, Prieto S. A review of peripheral tuberculosis arthritis. *Semin Arthritis Rheum*. 1988;18(2):142-149. doi:10.1016/0049-0172(88)90007-8.
75. Ellis ME, el-Ramahi KM, Al-Dalaan AN. Tuberculosis of peripheral joints: a dilemma in diagnosis. *Tuber Lung Dis*. 1993;74(6):399-404. doi:10.1016/0962-8479(93)90084-b.
76. Sant M, Bajaj H. Role of histopathology in the diagnosis of tuberculosis synovitis. *J Indian Med Assoc*. 1992;90(10):263-264.
77. Mondal A. Cytological diagnosis of vertebral tuberculosis with fine-needle aspiration biopsy. *J Bone Joint Surg Am*. 1994;76(2):181-184. doi:10.2106/00004623-199402000-00003.
78. Sun Q, Wang S, Dong W, et al. Diagnostic value of Xpert MTB/RIF Ultra for osteoarticular tuberculosis. *J Infect*. 2019;79(2):153-158. doi:10.1016/j.jinf.2019.06.006.
79. Wallace R, Cohen AS. Tuberculous arthritis: A report of two cases with review of biopsy and synovial fluid findings. *Am J Med*. 1976;61(2):277-282. doi:10.1016/0002-9343(76)90177-7.
80. Pertuiset E, Beaudreuil J, Liote F, et al. Spinal tuberculosis in adults. A study of 103 cases in a developed country, 1980-1994. *Medicine (Baltimore)*. 1999;78(5):309-320. doi:10.1097/00005792-199909000-00003.
81. Quale JM, Lipschik GY, Heurich AE. Management of tuberculous pericarditis. *Ann Thorac Surg*. 1987;43(6):653-655. doi:10.1016/s0003-4975(10)60243-3.
82. Strang JI, Kakaza HH, Gibson DG, et al. Controlled clinical trial of complete open surgical drainage and of prednisolone in treatment of tuberculosis pericardial effusion in Transkei. *Lancet*. 1988;2(8614):759-764. doi:10.1016/S0140-6736(88)92415-4.
83. Komsuoğlu B, Göldeli O, Kulan K, Komsuoğlu SS. The diagnostic and prognostic value of adenosine deaminase in tuberculous pericarditis. *Eur Heart J*. 1995;16(8):1126-1130. doi:10.1093/oxfordjournals.eurheartj.a061057.
84. Mayosi BM, Burgess LJ, Doubell AF. Tuberculous pericarditis. *Circulation*. 2005;112(23):3608-3616. doi:10.1161/circulationaha.105.543066.
85. Fowler NO, Manitsas GT. Infectious pericarditis. *Prog Cardiovasc Dis*. 1973;16(3):323-336. doi:10.1016/S0033-0620(73)80004-0.
86. Trautner BW, Darouiche RO. Tuberculous pericarditis: optimal diagnosis and management. *Clin Infect Dis*. 2001;33(7):954-961. doi:10.1086/322621.
87. Cegielski JP, Devlin BH, Morris AJ. Comparison of PCR, culture, and histopathology for diagnosis of tuberculosis pericarditis. *J Clin Microbiol*. 1997;35(12):3254-7. doi:10.1128/JCM.35.12.3254-3257.1997.
88. Yu G, Ye B, Chen D, et al. Comparison between the diagnostic validities of Xpert MTB/RIF and interferon-gamma release assays for tuberculosis pericarditis using pericardial tissue. *PLoS One*. 2017;12(12):e0188704. doi:10.1371/journal.pone.0188704.
89. Prout S, Benatar SR. Disseminated tuberculosis. A study of 62 cases. *S Afr Med J*. 1980;58(21):835-842.
90. Kim JH, Langston AA, Gallis HA. Miliary tuberculosis: epidemiology, clinical manifestations, diagnosis, and outcome. *Rev Infect Dis*. 1990;12(4):583-590. doi:10.1093/clinids/12.4.583.
91. Mert A, Bilir M, Tabak F, et al. Miliary tuberculosis: clinical manifestations, diagnosis and outcome in 38 adults. *Respirology*. 2001;6(3):217-224. doi:10.1046/j.1440-1843.2001.00328.x.
92. Maartens G, Willcox PA, Benatar SR. Miliary tuberculosis: rapid diagnosis, hematologic abnormalities, and outcome in 109 treated adults. *Am J Med*. 1990;89(3):291-296. doi:10.1016/0002-9343(90)90340-J.
93. Hussain SE, Irfan M, Abbasi M, et al. Clinical characteristics of 110 miliary tuberculosis patients from a low HIV prevalence country. *Int J Tuberc Lung Dis*. 2004;8(4):493-499.
94. Mert A, Arslan F, Kuyucu T, et al. Miliary tuberculosis: Epidemiological and clinical analysis of large-case series from moderate to low tuberculosis endemic Country. *Medicine (Baltimore)*. 2017;96(5):e5875. doi:10.1097/MD.0000000000005875.
95. Sharma SK, Mohan A. Miliary Tuberculosis. *Microbiol Spectr*. 2017;5(2). doi:10.1128/microbiolspec.TNMI7-0013-2016.
96. Wang JY, Hsueh PR, Wang SK, et al. Disseminated tuberculosis: a 10-year experience in a medical center. *Medicine (Baltimore)*. Jan 2007;86(1):39-46. doi:10.1097/MD.0b013e318030b605.
97. Dankner WM, Davis CE. Mycobacterium bovis as a significant cause of tuberculosis in children residing along the United States-Mexico border in the Baja California region. *Pediatrics*. 2000;105(6):E79-e79. doi:10.1542/peds.105.6.e79.
98. Fontanilla JM, Barnes A, von Reyn CF. Current diagnosis and management of peripheral tuberculous lymphadenitis. *Clin Infect Dis*. 2011;53(6):555-562. doi:10.1093/cid/cir454.
99. Shriner KA, Mathisen GE, Goetz MB. Comparison of mycobacterial lymphadenitis among persons infected with human immunodeficiency virus and seronegative controls. *Clin Infect Dis*. 1992;15(4):601-605. doi:10.1093/clind/15.4.601.
100. Cook VJ, Manfreda J, Hershfield ES. Tuberculous lymphadenitis in Manitoba: incidence, clinical characteristics and treatment. *Can Respir J*. 2004;11(4):279-286. doi:10.1155/2004/826501.
101. Lee MP, Chan JW, Ng KK, Li PC. Clinical manifestations of tuberculosis in HIV-infected patients. *Respirology*. 2000;5(4):423-426. doi:10.1111/j.1440-1843.2000.00287.x.
102. Long R, Boffa J. High HIV-TB co-infection rates in marginalized populations: evidence from Alberta in support of screening TB patients for HIV. *Can J Public Health*. 2010;101(3):202-204.
103. Martin T, Hoepfner VH, Ring ED. Superficial mycobacterial lymphadenitis in Saskatchewan. *CMAJ*. 1988;138(5):431-434.
104. Metchnikoff BG, Nolte FS, and Wallace RJ. Jr. Manual of clinical microbiology. In: *Mycobacterium*. 7th ed. Washington, D.C.: American Society for Microbiology; 1999:399-437.
105. Campbell IA, Ormerod LP, Friend JA, Jenkins PA, Prescott RJ. Six months versus nine months chemotherapy for tuberculosis

- of lymph nodes: final results. *Respir Med.* 1993;87(8):621–623. doi:10.1016/s0954-6111(05)80265-3.
106. Yuen AP, Wong SH, Tam CM, Chan SL, Wei WI, Lau SK. Prospective randomized study of thrice weekly six-month and nine-month chemotherapy for cervical tuberculous lymphadenopathy. *Otolaryngol Head Neck Surg.* 1997;116(2):189–192. doi:10.1016/s0194-5998(97)70323-1.
 107. van Loenhout-Rooyackers JH, Laheij RJ, Richter C, Verbeek AL. Shortening the duration of treatment for cervical tuberculous lymphadenitis. *Eur Respir J.* 2000;15(1):192–195. doi:10.1183/09031936.00.15119200.
 108. McMaster P, Isaacs D. Critical review of evidence for short course therapy for tuberculous adenitis in children. *Pediatr Infect Dis J.* 2000;19(5):401–404. doi:10.1097/00006454-200005000-00003.
 109. Donald PR. The chemotherapy of tuberculous lymphadenopathy in children. *Tuberculosis (Edinb).* 2010;90(4):213–224. doi:10.1016/j.tube.2010.05.001.
 110. Hawkey CR, Yap T, Pereira J, et al. Characterization and management of paradoxical upgrading reactions in HIV-uninfected patients with lymph node tuberculosis. *Clin Infect Dis.* 2005;40(9):1368–1371. doi:10.1086/429317.
 111. Blaikley JF, Khalid S, Ormerod LP. Management of peripheral lymph node tuberculosis in routine practice: an unselected 10-year cohort. *Int J Tuberc Lung Dis.* 2011;15(3):375–378.
 112. Seok H, Jeon JH, Oh KH, et al. Characteristics of residual lymph nodes after six months of antituberculous therapy in HIV-negative individuals with cervical tuberculous lymphadenitis. *BMC Infect Dis.* 2019;19(1):867. doi:10.1186/s12879-019-4507-0.
 113. Shaw JA, Diacon AH, Koegelenberg CFN. Tuberculous pleural effusion. *Respirology.* Oct 2019;24(10):962–971. doi:10.1111/resp.13673.
 114. Antonangelo L, Faria CS, Sales RK. Tuberculous pleural effusion: diagnosis & management. *Expert Rev Respir Med.* 2019;13(8):747–759. doi:10.1080/17476348.2019.1637737.
 115. Wen P, Wei M, Han C, He Y, Wang MS. Risk factors for tuberculous empyema in pleural tuberculosis patients. *Sci Rep.* 2019;9(1):19569. doi:10.1038/s41598-019-56140-4.
 116. Valdes L, Alvarez D, San JE, et al. Tuberculous pleurisy: a study of 254 patients. *Arch Intern Med.* 1998;158(18):2017–2021. doi:10.1001/archinte.158.18.2017.
 117. Kim HJ, Lee HJ, Kwon SY, et al. The prevalence of pulmonary parenchymal tuberculosis in patients with tuberculous pleuritis. *Chest.* 2006;129(5):1253–1258. doi:10.1378/chest.129.5.1253.
 118. Ko JM, Park HJ, Kim CH. Pulmonary changes of pleural TB: up-to-date CT imaging. *Chest.* 2014;146(6):1604–1611. doi:10.1378/chest.14-0196.
 119. Di Gennaro F, Pisani L, Veronese N, et al. Potential Diagnostic Properties of Chest Ultrasound in Thoracic Tuberculosis-A Systematic Review. *IJERPH.* Oct 12 2018;15(10):2235. doi:10.3390/ijerph15102235.
 120. Keshishyan S, Kaul V, Gupta A, Ahn C, Aronow WS, Epelbaum O. Sputum culture for the diagnosis of tuberculous pleural effusion: analysis of absolute and incremental yields. *Adv Respir Med.* 2019;87(5):281–288. doi:10.5603/ARM.2019.0050.
 121. Gopi A, Madhavan SM, Sharma SK, Sahn SA. Diagnosis and treatment of tuberculous pleural effusion in 2006. *Chest.* 2007;131(3):880–889. doi:10.1378/chest.06-2063.
 122. Cañete C, Galarza I, Granados A, Farrero E, Estopà R, Manresa F. Tuberculous pleural effusion: experience with six months of treatment with isoniazid and rifampicin. *Thorax.* 1994;49(11):1160–1161. doi:10.1136/thx.49.11.1160.
 123. Lai YF, Chao TY, Wang YH, Lin AS. Pigtail drainage in the treatment of tuberculosis pleural effusions: a randomised study. *Thorax.* 2003;58(2):149–151. doi:10.1136/thorax.58.2.149.
 124. Garcia-Rodriguez JF, Valcarce-Pardeiro N, Alvarez-Diaz H, Marino-Callejo A. Long-term efficacy of 6-month therapy with isoniazid and rifampin compared with isoniazid, rifampin, and pyrazinamide treatment for pleural tuberculosis. *Eur J Clin Microbiol Infect Dis.* 2019;38(11):2121–2126. doi:10.1007/s10096-019-03651-7.
 125. Wyser C, Walzl G, Smedema JP, Swart F, van Schalkwyk EM, van de Wal BW. Corticosteroids in the treatment of tuberculous pleurisy. A double-blind, placebo-controlled, randomized study. *Chest.* 1996;110(2):333–338. doi:10.1378/chest.110.2.333.
 126. Ryan H, Yoo J, Darsini P. Corticosteroids for tuberculous pleurisy. *Cochrane Database Syst Rev.* 2017;3:CD001876. doi:10.1002/14651858.CD001876.pub3.
 127. Bhuniya S, Arunabha DC, Choudhury S, Saha I, Roy TS, Saha M. Role of therapeutic thoracentesis in tuberculosis pleural effusion. *Ann Thorac Med.* 2012;7(4):215–219. doi:10.4103/1817-1737.102176.
 128. Light RW. Update on tuberculous pleural effusion. *Respirology.* 2010;15(3):451–458. doi:10.1111/j.1440-1843.2010.01723.x.
 129. Ferrer J. Tuberculous pleural effusion and tuberculous empyema. *Semin Respir Crit Care Med.* 2001;22(6):637–646. doi:10.1055/s-2001-18800.
 130. Long R, Barrie J, Stewart K, Peloquin CA. Treatment of a tuberculous empyema with simultaneous oral and intrapleural antituberculosis drugs. *Can Respir J.* 2008;15(5):241–243. doi:10.1155/2008/747206.
 131. Cho J-K, Choi YM, Lee SS, et al. Clinical features and outcomes of abdominal tuberculosis in southeastern Korea: 12 years of experience. *BMC Infect Dis.* 2018;18(1):699. doi:10.1186/s12879-018-3635-2.
 132. Debi U, Ravisankar V, Prasad KK, Sinha SK, Sharma AK. Abdominal tuberculosis of the gastrointestinal tract: revisited. *World J Gastroenterol.* 2014;20(40):14831–14840. doi:10.3748/wjg.v20.i40.14831.
 133. Mukewar S, Mukewar S, Ravi R, Prasad A, S Dua K. Colon tuberculosis: endoscopic features and prospective endoscopic follow-up after anti-tuberculosis treatment. *Clin Transl Gastroenterol.* 2012;3:e24. doi:10.1038/ctg.2012.19.
 134. Nagi B, Kochhar R, Bhasin DK, Singh K. Colorectal tuberculosis. *Eur Radiol.* 2003;13(8):1907–1912. doi:10.1007/s00330-002-1409-z.
 135. Almadi MA, Ghosh S, Aljebreen AM. Differentiating intestinal tuberculosis from Crohn's disease: a diagnostic challenge. *Am J Gastroenterol.* 2009;104(4):1003–1012. doi:10.1038/ajg.2008.162.
 136. Jakubowski A, Elwood RK, Enarson DA. Clinical features of abdominal tuberculosis. *J Infect Dis.* Oct 1988;158(4):687–692. doi:10.1093/infdis/158.4.687.
 137. Lee W-K, Van Tonder F, Tartaglia CJ, et al. CT appearances of abdominal tuberculosis. *Clin Radiol.* 2012;67(6):596–604. doi:10.1016/j.crad.2011.11.003.
 138. Balthazar EJ, Gordon R, Hulnick D. Ileocecal tuberculosis: CT and radiologic evaluation. *AJR Am J Roentgenol.* 1990;154(3):499–503. doi:10.2214/ajr.154.3.2106212.
 139. Pereira JM, Madureira AJ, Vieira A, Ramos I. Abdominal tuberculosis: imaging features. *Eur J Radiol.* 2005;55(2):173–180. doi:10.1016/j.ejrad.2005.04.015.
 140. Gupta P, Kumar S, Sharma V, et al. Common and uncommon imaging features of abdominal tuberculosis. *J Med Imaging Radiat Oncol.* 2019;63(3):329–339. doi:10.1111/1754-9485.12874.
 141. Pulimood AB, Ramakrishna BS, Kurian G, et al. Endoscopic mucosal biopsies are useful in distinguishing granulomatous colitis due to Crohn's disease from tuberculosis. *Gut.* 1999;45(4):537–541. doi:10.1136/gut.45.4.537.
 142. Ye Z, Lin Y, Cao Q, He Y, Xue L. Granulomas as the Most Useful Histopathological Feature in Distinguishing between Crohn's Disease and Intestinal Tuberculosis in Endoscopic Biopsy Specimens. *Medicine (Baltimore).* 2015;94(49):e2157. doi:10.1097/MD.0000000000002157.
 143. Pratap Mouli V, Munot K, Ananthakrishnan A, et al. Endoscopic and clinical responses to anti-tubercular therapy can differentiate intestinal tuberculosis from Crohn's disease. *Aliment Pharmacol Ther.* 2017;45(1):27–36. doi:10.1111/apt.13840.
 144. Patton PH, Parker CE, MacDonald JK, Chande N. Anti-tuberculous therapy for maintenance of remission in Crohn's disease. *Cochrane Database Syst Rev.* 2016;7:CD000299. doi:10.1002/14651858.CD000299.pub3.

145. Gupta A, Pratap Mouli V, Mohta S, et al. Anti-tubercular therapy given to differentiate Crohn's disease from intestinal tuberculosis predisposes to stricture formation. *J Crohns Colitis*. 2020;14(11):1611–1618. doi:10.1093/ecco-jcc/jjaa091.
146. Marrie TJ, Hershfield ES. Tuberculosis peritonitis in Manitoba. *Can J Surg*. 1978;21(6):533–536.
147. Tirumani SH, Ojili V, Gunabushanam G, et al. Imaging of tuberculosis of the abdominal viscera: beyond the intestines. *J Clin Imaging Sci*. 2013;3:17. doi:10.4103/2156-7514.111234.
148. Sharma MP, Bhatia V. Abdominal tuberculosis. *Indian J Med Res*. 2004;120(4):305–315.
149. Abdelaal A, Alfkey R, Abdelaziem S, et al. Role of laparoscopic peritoneal biopsy in the diagnosis of peritoneal tuberculosis. A seven-year experience. *Chirurgia (Bucur)*. 2014;109(3):330–334.
150. Yilmaz A, Ece F, Bayramgurler B, Akkaya E, Baran R. The value of Ca 125 in the evaluation of tuberculosis activity. *Respir Med*. 2001;95(8):666–669. doi:10.1053/rmed.2001.1121.
151. Jullien S, Jain S, Ryan H, Ahuja V. Six-month therapy for abdominal tuberculosis. *Cochrane Database Syst Rev*. 2016;11:CD012163. doi:10.1002/14651858.CD012163.pub2.
152. Park SH, Yang SK, Yang DH, et al. Prospective randomized trial of six-month versus nine-month therapy for intestinal tuberculosis. *Antimicrob Agents Chemother*. 2009;53(10):4167–4171. doi:10.1128/aac.00874-09.
153. Leonard MK, Blumberg HM. Musculoskeletal Tuberculosis. *Microbiol Spectr*. 2017;5(2). doi:10.1128/microbiolspec.TNMI7-0046-2017.
154. Sharma A, Chhabra HS, Chabra T, Mahajan R, Batra S, Sangondimath G. Demographics of tuberculosis of spine and factors affecting neurological improvement in patients suffering from tuberculosis of spine: a retrospective analysis of 312 cases. *Spinal Cord*. 2017;55(1):59–63. doi:10.1038/sc.2016.85.
155. Johansen IS, Nielsen SL, Hove M, et al. Characteristics and Clinical Outcome of Bone and Joint Tuberculosis From 1994 to 2011: A Retrospective Register-based Study in Denmark. *Clin Infect Dis*. 2015;61(4):554–562. doi:10.1093/cid/civ326.
156. Broderick C, Hopkins S, Mack DJF, et al. Delays in the diagnosis and treatment of bone and joint tuberculosis in the United Kingdom. *Bone Joint J*. 2018;100-B(1):119–124. doi:10.1302/0301-620x.100b1.Bjj-2017-0357.R1.
157. Singh R, Magu NK, Rohilla RK. Clinicoradiologic Profile of Involvement and Healing in Tuberculosis of the Spine. *Ann Med Health Sci Res*. 2016;6(5):311–327. doi:10.4103/amhsr.amhsr_188_15.
158. Dunn RN, Ben Husien M. Spinal tuberculosis: review of current management. *Bone Joint J*. 2018;100-B(4):425–431. doi:10.1302/0301-620x.100b4.Bjj-2017-1040.R1.
159. Boachie-Adjei O, Squillante RG. Tuberculosis of the spine. *Orthop Clin North Am*. 1996;27(1):95–103.
160. Rezai AR, Lee M, Cooper PR, Errico TJ, Koslow M. Modern management of spinal tuberculosis. *Neurosurgery*. 1995;36(1):87–97. doi:10.1227/00006123-199501000-00011.
161. Bhandari A, Garg RK, Malhotra HS, et al. Outcome assessment in conservatively managed patients with cervical spine tuberculosis. *Spinal Cord*. 2014;52(6):489–493. doi:10.1038/sc.2014.49.
162. Kim JH, Ahn JY, Jeong SJ, et al. Prognostic factors for unfavourable outcomes of patients with spinal tuberculosis in a country with an intermediate tuberculosis burden: a multicentre cohort study. *Bone Joint J*. 2019;101-B(12):1542–1549. doi:10.1302/0301-620x.101b12.Bjj-2019-0558.R1.
163. Kilborn T, Janse van Rensburg P, Candy S. Pediatric and adult spinal tuberculosis: imaging and pathophysiology. *Neuroimaging Clin N Am*. 2015;25(2):209–231. doi:10.1016/j.nic.2015.01.002.
164. Burrill J, Williams CJ, Bain G, Conder G, Hine AL, Misra RR. Tuberculosis: a radiologic review. *Radiographics*. 2007;27(5):1255–1273. doi:10.1148/rg.275065176.
165. Boxer DI, Pratt C, Hine AL, McNicol M. Radiological features during and following treatment of spinal tuberculosis. *Br J Radiol*. 1992;65(774):476–479. doi:10.1259/0007-1285-65-774-476.
166. Joseffer SS, Cooper PR. Modern imaging of spinal tuberculosis. *J Neurosurg Spine*. 2005;2(2):145–150. doi:10.3171/spi.2005.2.2.0145.
167. Jain AK, Sreenivasan R, Mukunth R, Dhammi IK. Tubercular spondylitis in children. *Indian J Orthop*. 2014;48(2):136–144. doi:10.4103/0019-5413.128747.
168. Shikhare SN, Singh DR, Shimpi TR, Peh WC. Tuberculous osteomyelitis and spondylodiscitis. *Semin Musculoskelet Radiol*. 2011;15(5):446–458. doi:10.1055/s-0031-1293491.
169. Gu Y, Wang G, Dong W, et al. Xpert MTB/RIF and GenoType MTBDRplus assays for the rapid diagnosis of bone and joint tuberculosis. *Int J Infect Dis*. 2015;36:27–30. doi:10.1016/j.ijid.2015.05.014.
170. Held M, Laubscher M, Mears S, et al. Diagnostic Accuracy of the Xpert MTB/RIF Assay for Extrapulmonary Tuberculosis in Children With Musculoskeletal Infections. *Pediatr Infect Dis J*. 2016;35(11):1165–1168. doi:10.1097/INF.0000000000001271.
171. Held M, Laubscher M, Zar HJ, Dunn RN. GeneXpert polymerase chain reaction for spinal tuberculosis: an accurate and rapid diagnostic test. *Bone Joint J*. 2014;96-B(10):1366–1369. doi:10.1302/0301-620x.96b10.34048.
172. Shen Y, Yu G, Zhong F, Kong X. Diagnostic accuracy of the Xpert MTB/RIF assay for bone and joint tuberculosis: A meta-analysis. *PLoS One*. 2019;14(8):e0221427. doi:10.1371/journal.pone.0221427.
173. Cao G, Rao J, Cai Y, et al. Analysis of Treatment and Prognosis of 863 Patients with Spinal Tuberculosis in Guizhou Province. *Biomed Res Int*. 2018;2018:3265735. doi:10.1155/2018/3265735.
174. Wei G, Mu J, Wang G, et al. The reliability analysis of Xpert-positive result for smear-negative and culture-negative specimen collected from bone and joint tuberculosis suspects. *J Thorac Dis*. 2016;8(6):1205–1209. doi:10.21037/jtd.2016.04.19.
175. Faroug R, Psyllakis P, Gulati A, Makvana S, Pareek M, Mangwani J. Diagnosis and treatment of tuberculosis of the foot and ankle-A literature review. *Foot (Edinb)*. 2018;37:105–112. doi:10.1016/j.foot.2018.07.005.
176. Barik S, Choudhury AK, Singh V, Bali S. Extra-Spinal Osteoarticular Tuberculosis: A Retrospective Analysis of 103 Cases. *Curr Health Sci J*. 2019;45(2):142–147. doi:10.12865/chsj.45.02.03.
177. Watts HG, Lifeso RM. Tuberculosis of bones and joints. *J Bone Joint Surg Am*. 1996;78(2):288–298. doi:10.2106/00004623-199602000-00019.
178. Hogan JJ, Hurtado RM, Nelson SB. Mycobacterial Musculoskeletal Infections. *Infect Dis Clin North Am*. 2017;31(2):369–382. doi:10.1016/j.idc.2017.01.007.
179. Choi JA, Koh SH, Hong SH, Koh YH, Choi JY, Kang HS. Rheumatoid arthritis and tuberculosis arthritis: differentiating MRI features. *AJR Am J Roentgenol*. 2009;193(5):1347–1353. doi:10.2214/AJR.08.2164.
180. Sawlani V, Chandra T, Mishra RN, Aggarwal A, Jain UK, Gujral RB. MRI features of tuberculosis of peripheral joints. *Clin Radiol*. 2003;58(10):755–762. doi:10.1016/s0009-9260(03)00271-x.
181. Sanghvi DA, Iyer VR, Deshmukh T, Hoskote SS. MRI features of tuberculosis of the knee. *Skeletal Radiol*. 2009;38(3):267–273. doi:10.1007/s00256-008-0617-2.
182. Shu CC, Wang JY, Yu CJ, Lee LN. Mycobacterial arthritis of large joints. *Ann Rheum Dis*. 2009;68(9):1504–1505. doi:10.1136/ard.2008.095554.
183. Donald PR. The chemotherapy of osteo-articular tuberculosis with recommendations for treatment of children. *J Infect*. 2011;62(6):411–439. doi:10.1016/j.jinf.2011.04.239.
184. Controlled trial of short-course regimens of chemotherapy in the ambulatory treatment of spinal tuberculosis. Results at three years of a study in Korea. Twelfth report of the Medical Research Council Working Party on Tuberculosis of the Spine. *J Bone Joint Surg Br*. 1993;75(2):240–248. doi:10.1302/0301-620x.75b2.8444944.
185. A 15-year assessment of controlled trials of the management of tuberculosis of the spine in Korea and Hong Kong. Thirteenth Report of the Medical Research Council Working Party on

- Tuberculosis of the Spine. *J Bone Joint Surg Br.* 1998;80(3):456–462. doi:10.1302/0301-620x.80b3.8544.
186. Parthasarathy R, Sriram K, Santha T, Prabhakar R, Somasundaram PR, Sivasubramanian S. Short-course chemotherapy for tuberculosis of the spine. A comparison between ambulant treatment and radical surgery-ten-year report. *J Bone Joint Surg Br.* 1999;81(3):464–471. doi:10.1302/0301-620x.81b3.9043.
 187. Jain AK, Sreenivasan R, Saini NS, Kumar S, Jain S, Dhammi IK. Magnetic resonance evaluation of tubercular lesion in spine. *Int Orthop.* 2012;36(2):261–269. doi:10.1007/s00264-011-1380-x.
 188. Mittal S, Jain AK, Chakraborti KL, Aggarwal AN, Upreti L, Bhayana H. Evaluation of Healed Status in Tuberculosis of Spine by Fluorodeoxyglucose-positron Emission Tomography/Computed Tomography and Contrast Magnetic Resonance Imaging. *Indian J Orthop.* 2019;53(1):160–168. doi:10.4103/ortho.IJOrtho_224_18.
 189. Jutte PC, Van Loenhout-Rooyackers JH. Routine surgery in addition to chemotherapy for treating spinal tuberculosis. *Cochrane Database Syst Rev.* 2006;(1):Cd004532. doi:10.1002/14651858.CD004532.pub2.
 190. Thwaites G, Fisher M, Hemingway C, Scott G, Solomon T, Innes J. British Infection Society guidelines for the diagnosis and treatment of tuberculosis of the central nervous system in adults and children. *J Infect.* 2009;59(3):167–187. doi:10.1016/j.jinf.2009.06.011.
 191. Leonard JM. Central Nervous System Tuberculosis. *Microbiol Spectr.* 2017;5(2). doi:10.1128/microbiolspec.TNMI7-0044-2017.
 192. Arvanitakis Z, Long RL, Hershfield ES, et al. M. tuberculosis molecular variation in CNS infection: evidence for strain-dependent neurovirulence. *Neurology.* 1998;50(6):1827–1832. doi:10.1212/WNL.50.6.1827.
 193. Hsu PC, Yang CC, Ye JJ, Huang PY, Chiang PC, Lee MH. Prognostic factors of tuberculosis meningitis in adults: a 6-year retrospective study at a tertiary hospital in northern Taiwan. *J Microbiol Immunol Infect.* 2010;43(2):111–118. doi:10.1016/S1684-1182(10)60018-7.
 194. Wilkinson RJ, Rohlwick U, Misra UK, Tuberculous Meningitis International Research Consortium, et al. Tuberculous meningitis. *Nat Rev Neurol.* 2017;13(10):581–598. doi:10.1038/nrneurol.2017.120.
 195. Chiang SS, Khan FA, Milstein MB, et al. Treatment outcomes of childhood tuberculous meningitis: a systematic review and meta-analysis. *Lancet Infect Dis.* Oct 2014;14(10):947–957. doi:10.1016/S1473-3099(14)70852-7.
 196. Thwaites GE, Nguyen DB, Nguyen HD, et al. Dexamethasone for the treatment of tuberculous meningitis in adolescents and adults. *N Engl J Med.* 2004;351(17):1741–1751. doi:10.1056/NEJMoa040573.
 197. Torok ME, Nguyen DB, Tran TH, et al. Dexamethasone and long-term outcome of tuberculous meningitis in Vietnamese adults and adolescents. *PLoS One.* 2011;6(12):e27821. doi:10.1371/journal.pone.0027821.
 198. Erdem H, Ozturk-Engin D, Tireli H, et al. Hamsi scoring in the prediction of unfavorable outcomes from tuberculous meningitis: results of Haydarpasa-II study. *J Neurol.* 2015;262(4):890–898. doi:10.1007/s00415-015-7651-5.
 199. Marais BJ, Heemskerk AD, Marais SS, Tuberculous Meningitis International Research Consortium, et al. Standardized Methods for Enhanced Quality and Comparability of Tuberculous Meningitis Studies. *Clin Infect Dis.* 2017;64(4):501–509. doi:10.1093/cid/ciw757.
 200. Thao LTP, Wolbers M, Heemskerk AD, et al. Dynamic Prediction of Death in Patients With Tuberculous Meningitis Using Time-updated Glasgow Coma Scale and Plasma Sodium Measurements. *Clin Infect Dis.* 2020;70(5):827–834. doi:10.1093/cid/ciz262.
 201. Sanei Taheri M, Karimi MA, Haghighatkhah H, Pourghorban R, Samadian M, Delavar Kasmaei H. Central nervous system tuberculosis: an imaging-focused review of a reemerging disease. *Radiol Res Pract.* 2015;2015:202806. doi:10.1155/2015/202806.
 202. Botha H, Ackerman C, Candy S, Carr JA, Griffith-Richards S, Bateman KJ. Reliability and diagnostic performance of CT imaging criteria in the diagnosis of tuberculosis meningitis. *PLoS One.* 2012;7(6):e38982. doi:10.1371/journal.pone.0038982.
 203. Pienaar M, Andronikou S, van Toorn R. MRI to demonstrate diagnostic features and complications of TBM not seen with CT. *Childs Nerv Syst.* 2009;25(8):941–947. doi:10.1007/s00381-008-0785-3.
 204. Bourgi K, Fiske C, Sterling TR. Tuberculosis Meningitis. *Curr Infect Dis Rep.* 2017;19(11):39. doi:10.1007/s11908-017-0595-4.
 205. Stewart SM. The bacteriological diagnosis of tuberculous meningitis. *J Clin Pathol.* 1953;6(3):241–242. doi:10.1136/jcp.6.3.241.
 206. Kennedy DH, Fallon RJ. Tuberculous meningitis. *JAMA.* 1979;241(3):264–268.
 207. Chakravorty S, Simmons AM, Rowneki M, et al. The New Xpert MTB/RIF Ultra: Improving Detection of Mycobacterium tuberculosis and Resistance to Rifampin in an Assay Suitable for Point-of-Care Testing. *mBio.* 2017;8(4). doi:10.1128/mBio.00812-17.
 208. World Health Organization. *Automated Real-Time Nucleic Acid Amplification Technology for Rapid and Simultaneous Detection of Tuberculosis and Rifampicin Resistance: Xpert MTB/RIF Assay for the Diagnosis of Pulmonary and Extra-Pulmonary TB in Adults and Children. Policy Update.* Geneva: World Health Organization. 2013.
 209. World Health Organization. *WHO meeting report of a technical expert consultation: non-inferiority analysis of Xpert MTF/RIF Ultra compared to Xpert MTB/RIF.* Geneva: World Health Organization. 2017. <https://apps.who.int/iris/handle/10665/254792>. License: CC BY-NC-SA 3.0 IGO. Accessed March 2, 2022.
 210. Kent SJ, Crowe SM, Yung A, Lucas CR, Mijch AM. Tuberculous meningitis: a 30-year review. *Clin Infect Dis.* 1993;17(6):987–994. doi:10.1093/clinids/17.6.987.
 211. Thwaites G, Chau TT, Mai NT, Drobniewski F, McAdam K, Farrar J. Tuberculous meningitis. *J Neurol Neurosurg Psychiatry.* 2000;68(3):289–299. doi:10.1136/jnnp.68.3.289.
 212. Rock RB, Olin M, Baker CA, Molitor TW, Peterson PK. Central nervous system tuberculosis: pathogenesis and clinical aspects. *Clin Microbiol Rev.* 2008;21(2):243–261, table of contents. doi:10.1128/CMR.00042-07.
 213. Nahid P, Dorman SE, Alipanah N, et al. Official American Thoracic Society/Centers for Disease Control and Prevention/Infectious Diseases Society of America Clinical Practice Guidelines: Treatment of Drug-Susceptible Tuberculosis. *Clin Infect Dis.* 2016;63(7):e147–e195. doi:10.1093/cid/ciw376.
 214. National Institute for Health and Care Excellence. Tuberculosis [NICE Guideline No. 33]. November 13, 2020. <https://www.nice.org.uk/guidance/ng33>. Accessed November 13, 2020.
 215. American Academy of Pediatrics. *Red Book: 2018 Report of the Committee on Infectious Diseases.* 31st ed. Itasca, IL: American Academy of Pediatrics; 2018.
 216. World Health Organization. Guidelines for treatment of drug-susceptible tuberculosis and patient care, 2017 update. World Health Organization; 2017. <https://apps.who.int/iris/handle/10665/255052>. Accessed March 2, 2022.
 217. World Health Organization. *Treatment of Tuberculosis: guidelines.* 4th ed. Geneva: World Health Organization; 2010. <https://apps.who.int/iris/handle/10665/44165>. Accessed on March 2, 2022.
 218. Cresswell FV, Te Brake L, Atherton R, et al. Intensified antibiotic treatment of tuberculosis meningitis. *Expert Rev Clin Pharmacol.* 2019;12(3):267–288. doi:10.1080/17512433.2019.1552831.
 219. ClinicalTrials.gov [Internet]. Intensified Tuberculosis Treatment to Reduce the Mortality of Patients With Tuberculosis Meningitis (INTENSE-TBM). National Library of Medicine (US). Updated August 28, 2020. Accessed September 28, 2020. <https://clinicaltrials.gov/ct2/show/NCT04145258>.
 220. Marais S, Cresswell FV, Hamers RL, et al. High dose oral rifampicin to improve survival from adult tuberculosis meningitis: A randomised placebo-controlled double-blinded phase III trial

- (the HARVEST study). *Wellcome Open Research*. 2019;4:190. doi:10.12688/wellcomeopenres.15565.1.
221. ClinicalTrials.gov [Internet]. Optimizing Treatment to Improve TBM Outcomes in Children (TBM-KIDS). National Library of Medicine (US). Sept 28, 2020. Updated Aug 13, 2020. <https://clinicaltrials.gov/ct2/show/NCT02958709>. Accessed September 29, 2020.
 222. Charlie L, Abay SM, Tesfaye A, Mlera RN, Mwangi S, Goiretti M. Safety and efficacy of high-dose rifampicin in the management of tuberculous meningitis: Systematic review and meta-analysis. *Int J Mycobacteriol*. 2021;10(3):312–319. doi:10.4103/ijmy.ijmy_135_21.
 223. Cresswell FV, Meya DB, Kagimu E, et al. High-Dose Oral and Intravenous Rifampicin for the Treatment of Tuberculosis Meningitis in Predominantly Human Immunodeficiency Virus (HIV)-Positive Ugandan Adults: A Phase II Open-Label Randomized Controlled Trial. *Clin Infect Dis*. 2021;73(5):876–884. doi:10.1093/cid/ciab162.
 224. Wasserman S, Davis A, Stek C, et al. Plasma Pharmacokinetics of High-Dose Oral versus Intravenous Rifampicin in Patients with Tuberculous Meningitis: a Randomized Controlled Trial. *Antimicrob Agents Chemother*. 2021;65(8):e0014021. doi:10.1128/AAC.00140-21.
 225. Svensson EM, Dian S, Te Brake L, et al. Model-Based Meta-analysis of Rifampicin Exposure and Mortality in Indonesian Tuberculous Meningitis Trials. *Clin Infect Dis*. 2020;71(8):1817–1823. Nov 5doi:10.1093/cid/ciz1071.
 226. Arbiv. *Thorax*. 2022.
 227. Ellard GA, Humphries MJ, Allen BW. Cerebrospinal fluid drug concentrations and the treatment of tuberculous meningitis. *Am Rev Respir Dis*. 1993;148(3):650–655. doi:10.1164/ajrcm/148.3.650.
 228. Rizvi I, Malhotra HS, Garg RK, Kumar N, Uniyal R, Pandey S. Fluoroquinolones in the management of tuberculous meningitis: Systematic review and meta-analysis. *J Infect*. 2018;77(4):261–275. doi:10.1016/j.jinf.2018.06.009.
 229. Ruslami R, Ganiem AR, Dian S, et al. Intensified regimen containing rifampicin and moxifloxacin for tuberculosis meningitis: an open-label, randomised controlled phase 2 trial. *Lancet Infect Dis*. 2013;13(1):27–35. doi:10.1016/S1473-3099(12)70264-5.
 230. Heemskerk AD, Bang ND, Mai NT, et al. Intensified Antituberculosis Therapy in Adults with Tuberculous Meningitis. *N Engl J Med*. 2016;374(2):124–134. doi:10.1056/NEJMoA1507062.
 231. Heemskerk AD, Nguyen MTH, Dang HTM, et al. Clinical Outcomes of Patients With Drug-Resistant Tuberculosis Meningitis Treated With an Intensified Antituberculosis Regimen. *Clin Infect Dis*. 2017;65(1):20–28. doi:10.1093/cid/cix230.
 232. Sun F, Ruan Q, Wang J, et al. Linezolid manifests a rapid and dramatic therapeutic effect for patients with life-threatening tuberculous meningitis. *Antimicrob Agents Chemother*. 2014;58(10):6297–6301. doi:10.1128/AAC.02784-14.
 233. Li H, Lu J, Liu J, Zhao Y, Ni X, Zhao S. Linezolid is Associated with Improved Early Outcomes of Childhood Tuberculosis Meningitis. *Pediatr Infect Dis J*. 2016;35(6):607–610. doi:10.1097/INF.0000000000001114.
 234. Alvarez-Uria G, Midde M, Pakam R, Naik PK. Initial Antituberculous Regimen with Better Drug Penetration into Cerebrospinal Fluid Reduces Mortality in HIV Infected Patients with Tuberculous Meningitis: Data from an HIV Observational Cohort Study. *Tuberc Res Treat*. 2013;2013:242604. doi:10.1155/2013/242604.
 235. Donald PR, Schoeman JE, Van Zyl LE, De Villiers JN, Pretorius M, Springer P. Intensive short course chemotherapy in the management of tuberculous meningitis. *Int J Tuberc Lung Dis*. 1998;2(9):704–711.
 236. Jullien S, Ryan H, Modi M, Bhatia R. Six months therapy for tuberculous meningitis. *Cochrane Database Syst Rev*. 2016;9:CD012091. doi:10.1002/14651858.CD012091.pub2.
 237. World Health Organization. *Guidelines for treatment of drug-susceptible tuberculosis and patient care, 2017 update*. 2017. Licence: CC BY-NC-SA 3.0 IGO.
 238. Nahid P, Dorman SE, Alipanah N, et al. Official American Thoracic Society/Centers for Disease Control and Prevention/ Infectious Diseases Society of America Clinical Practice Guidelines: Treatment of Drug-Susceptible Tuberculosis. *Clin Infect Dis*. 2016;63(7):e147–e195. doi:10.1093/cid/ciw376.
 239. Prasad K, Singh MB, Ryan H. Corticosteroids for managing tuberculous meningitis. *Cochrane Database Syst Rev*. 2016;4:CD002244. doi:10.1002/14651858.CD002244.pub4.
 240. Donovan J, Thwaites GE, Huynh J. Tuberculosis meningitis: where to from here? *Curr Opin Infect Dis*. 2020;33(3):259–266. doi:10.1097/QCO.0000000000000648.
 241. Schoeman JE, Janse van Rensburg A, Laubscher JA, Springer P. The role of aspirin in childhood tuberculous meningitis. *J Child Neurol*. 2011;26(8):956–962. doi:10.1177/0883073811398132.
 242. Mai NT, Dobbs N, Phu NH, et al. A randomised double blind placebo controlled phase 2 trial of adjunctive aspirin for tuberculous meningitis in HIV-uninfected adults. *eLife*. 2018;7:e33478. doi:10.7554/eLife.33478.
 243. ClinicalTrials.gov [Internet]. Linezolid, Aspirin and Enhanced Dose Rifampicin in HIV-TBM (LASER-TBM). National Library of Medicine (US). Sept 28 2020. Updated August 14, 2020. <https://clinicaltrials.gov/ct2/show/NCT03927313>.
 244. ISRCTN registry [Internet]. SURE: Short intensive treatment for children with tuberculosis meningitis. BMC. September, 28, 2020. Updated Sept 23, 2020. doi:10.1186/ISRCTN40829906.
 245. Rohlwick UK, Kilborn T, Wieselthaler N, Banderker E, Zwane E, Figaji AA. Imaging Features of the Brain, Cerebral Vessels and Spine in Pediatric Tuberculous Meningitis With Associated Hydrocephalus. *Pediatr Infect Dis J*. Oct 2016;35(10):e301–e310. doi:10.1097/INF.0000000000001236.
 246. Marais S, Van Toorn R, Chow FC, Tuberculous Meningitis International Research Consortium, et al. Management of intracranial tuberculous mass lesions: how long should we treat for? *Wellcome Open Res*. 2019;4:158. doi:10.12688/wellcomeopenres.15501.2.
 247. Donovan J, Figaji A, Imran D, Phu NH, Rohlwick U, Thwaites GE. The neurocritical care of tuberculosis meningitis. *Lancet Neurol*. 2019;18(8):771–783. doi:10.1016/S1474-4422(19)30154-1.
 248. Crump JA, Reller LB. Two decades of disseminated tuberculosis at a university medical center: the expanding role of mycobacterial blood culture. *Clin Infect Dis*. 2003;37(8):1037–1043. Oct 15doi:10.1086/378273.
 249. Meira L, Chaves C, Araujo D, et al. Predictors and outcomes of disseminated tuberculosis in an intermediate burden setting. *Pulmonology*. 2019;25(6):320–327. doi:10.1016/j.pulmoe.2018.11.001.
 250. Lee J, Lim JK, Kim EJ, et al. Comparison of clinical manifestations and treatment outcome according to age groups in adult patients with miliary tuberculosis. *J Thorac Dis*. 2018;10(5):2881–2889. doi:10.21037/jtd.2018.04.139.
 251. Mohan A, Sharma SK, Pande JN. Acute respiratory distress syndrome (ARDS) in miliary tuberculosis: a twelve year experience. *Indian J Chest Dis Allied Sci*. 1996;38(3):157–162.
 252. Boushab BM, Basco LK. Miliary tuberculosis and acute respiratory distress syndrome. *J Clin Tuberc Other Mycobact Dis*. 2019;16:100113. doi:10.1016/j.jctube.2019.100113.
 253. Long R, O'Connor R, Palayew M, Hershfield E, Manfreda J. Disseminated tuberculosis with and without a miliary pattern on chest radiograph: a clinical-pathologic-radiologic correlation. *Int J Tuberc Lung Dis*. 1997;1(1):52–58.
 254. Sharma SK, Mohan A, Sharma A. Miliary tuberculosis: A new look at an old foe. *J Clin Tuberc Other Mycobact Dis*. 2016;3:13–27. doi:10.1016/j.jctube.2016.03.003.
 255. Heiden D, Saranchuk P, Keenan JD, et al. Eye examination for early diagnosis of disseminated tuberculosis in patients with AIDS. *Lancet Infect Dis*. 2016;16(4):493–499. doi:10.1016/S1473-3099(15)00269-8.

256. Mehta S. Patterns of ocular inflammation in patients with milary tuberculosis. *F1000Res*. 2017;6:412. doi:10.12688/f1000research.11035.1.
257. Rieder HL, Kelly GD, Bloch AB, Cauthen GM, Snider DE Jr. Tuberculosis diagnosed at death in the United States. *Chest*. 1991;100(3):678–681. doi:10.1378/chest.100.3.678.
258. Optican RJ, Ost A, Ravin CE. High-resolution computed tomography in the diagnosis of milary tuberculosis. *Chest*. 1992;102(3):941–943. doi:10.1378/chest.102.3.941.
259. Lefebvre N, Argemi X, Meyer N, et al. Clinical usefulness of 18F-FDG PET/CT for initial staging and assessment of treatment efficacy in patients with lymph node tuberculosis. *Nucl Med Biol*. 2017;50:17–24. doi:10.1016/j.nucmedbio.2017.04.003.
260. Abbara A, Davidson RN. Etiology and management of genitourinary tuberculosis. *Nat Rev Urol*. 2011;8(12):678–688. doi:10.1038/nrurol.2011.172.
261. Fanning A. Tuberculosis: 6. Extrapulmonary disease. *CMAJ*. 1999;160(11):1597–1603.
262. Sourial MW, Brimo F, Horn R, Andonian S. Genitourinary tuberculosis in North America: A rare clinical entity. *Can Urol Assoc J*. 2015;9(7-8):E484–9. doi:10.5489/cuaj.2643.
263. Cao Y, Fan Y, Chen Y, et al. Gross Hematuria Is More Common in Male and Older Patients with Renal Tuberculosis in China: A Single-Center 15-Year Clinical Experience. *Urol Int*. 2017;99(3):290–296. doi:10.1159/000464472.
264. Eastwood JB, Corbishley CM, Grange JM. Tuberculosis and the kidney. *J Am Soc Nephrol*. 2001;12(6):1307–1314. doi:10.1681/ASN.V1261307.
265. Merchant S, Bharati A, Merchant N. Tuberculosis of the genitourinary system-Urinary tract tuberculosis: Renal tuberculosis-Part II. *Indian J Radiol Imaging*. 2013;23(1):64–77. doi:10.4103/0971-3026.113617.
266. Maclean KA, Becker AK, Chang SD, Harris AC. Extrapulmonary tuberculosis: imaging features beyond the chest. *Can Assoc Radiol J*. 2013;64(4):319–324. doi:10.1016/j.carj.2012.07.002.
267. Wang Y, Wu JB, Qin GC, et al. Computerised tomography and intravenous pyelography in urinary tuberculosis: a retrospective descriptive study. *Int J Tuberc Lung Dis*. 2015;19(12):1441–1447. doi:10.5588/ijtld.14.0888.
268. Webster D, Long R, Shandro C, et al. Fluoroquinolone resistance among renal tuberculosis isolates in Alberta Canada;1990-2003. *International Journal of Tuberculosis and Lung Disease*. 2010;14:217–222.
269. Goldfarb DS. SL. Tuberculosis of the genitourinary tract. 1996. 609–622. Tuberculosis.
270. Grace GA, Devaleenal DB, Natrajan M. Genital tuberculosis in females. *Indian J Med Res*. 2017;145(4):425–436. doi:10.4103/ijmr.IJMR_1550_15.
271. Sharma JB, Sharma E, Sharma S, Dharmendra S. Female genital tuberculosis: Revisited. *Indian J Med Res*. 2018;148(Suppl):S71–s83. Decdoi:10.4103/ijmr.IJMR_648_18.
272. Gupta B, Shree S, Rajaram S, Goel N. Genital tuberculosis: Unusual presentations. *Int J Mycobacteriol*. 2016;5(3):357–359. doi:10.1016/j.ijmyco.2016.06.017.
273. Khanna A, Agrawal A. Markers of genital tuberculosis in infertility. *Singapore Med J*. 2011;52(12):864–867.
274. Shah HU, Sannanjanja B, Baheti AD, Udare AS, Badhe PV. Hysterosalpingography and ultrasonography findings of female genital tuberculosis. *Diagn Interv Radiol*. 2015;21(1):10–15. doi:10.5152/dir.2014.13517.
275. Sharma JB, Roy KK, Pushparaj M, Kumar S, Malhotra N, Mittal S. Laparoscopic findings in female genital tuberculosis. *Arch Gynecol Obstet*. 2008;278(4):359–364. doi:10.1007/s00404-008-0586-7.
276. Kumar P, Shah NP, Singhal A, et al. Association of tuberculous endometritis with infertility and other gynecological complaints of women in India. *J Clin Microbiol*. 2008;46(12):4068–4070. doi:10.1128/jcm.01162-08.
277. Neelam BM, Namita K. Genital tuberculosis and its consequences on subsequent fertility. *J Obstet Gynecol India*. 2005;2005(6):534–537.
278. Flibotte JJ, Lee GE, Buser GL, et al. Infertility, in vitro fertilization and congenital tuberculosis. *J Perinatol*. 2013;33(7):565–568. doi:10.1038/jp.2012.146.
279. Yadav S, Singh P, Hemal A, Kumar R. Genital tuberculosis: current status of diagnosis and management. *Transl Androl Urol*. 2017;6(2):222–233. doi:10.21037/tau.2016.12.04.
280. Aziz EM, Abdelhak K, Hassan FM. Tuberculous prostatitis: mimicking a cancer. *Pan Afr Med J*. 2016;25:130 doi:10.11604/pamj.2016.25.130.7577.
281. Gonzalez OY, Musher DM, Brar I, et al. Spectrum of bacille Calmette-Guérin (BCG) infection after intravesical BCG immunotherapy. *Clin Infect Dis*. 2003;36(2):140–148. doi:10.1086/344908.
282. Kim S, Wise G, Schleel P, Khani F, Herr H. Adverse Effects of Intravesical bacillus Calmette-Guerin Therap. *AUA Update Series*. 2018;37(Lesson 8)
283. Lamm DL. Efficacy and safety of bacille Calmette-Guérin immunotherapy in superficial bladder cancer. *Clin Infect Dis*. 2000;31 Suppl 3:S86–S90. doi:10.1086/314064.
284. Larsen ES, Nordholm AC, Lillebaek T, Holden IK, Johansen IS. The epidemiology of bacille Calmette-Guérin infections after bladder instillation from 2002 through 2017: a nationwide retrospective cohort study. *BJU Int*. 2019;124(6):910–916. doi:10.1111/bju.14793.
285. Pérez-Jacoiste Asín MA, Fernández-Ruiz M, López-Medrano F, et al. Bacillus Calmette-Guérin (BCG) infection following intravesical BCG administration as adjunctive therapy for bladder cancer: incidence, risk factors, and outcome in a single-institution series and review of the literature. *Medicine (Baltimore)*. 2014;93(17):236–254. doi:10.1097/MD.000000000000119.
286. Marques M, Vazquez D, Sousa S, Mesquita G, Duarte M, Ferreira R. Disseminated Bacillus Calmette-Guérin (BCG) infection with pulmonary and renal involvement: A rare complication of BCG immunotherapy. *A Case Report and Narrative Review. Pulmonology*. 2020;26(6):346–352. doi:10.1016/j.pulmoe.2019.10.001.
287. Waked R, Choucair J, Chehata N, Haddad E, Saliba G. Intravesical Bacillus Calmette-Guérin (BCG) treatment's severe complications: A single institution review of incidence, presentation and treatment outcome. *J Clin Tuberc Other Mycobact Dis*. 2020;19:100149. doi:10.1016/j.jctube.2020.100149.
288. Cek M, Lenk S, Naber KG, Members of the Urinary Tract Infection (UTI) Working Group of the European Association of Urology (EAU) Guidelines Office, et al. EAU guidelines for the management of genitourinary tuberculosis. *Eur Urol*. 2005;48(3):353–362. doi:10.1016/j.eururo.2005.03.008.
289. Gow JG. Genitourinary tuberculosis: a 7-year review. *Br J Urol*. 1979;51(4):239–244. doi:10.1111/j.1464-410x.1979.tb04700.x.
290. Migliori GB, Sotgiu G, Rosales-Klintz S, et al. ERS/ECDC Statement: European Union standards for tuberculosis care, 2017 update. *Eur Respir J*. 2018;51(5):1702678. doi:10.1183/13993003.02678-2017.
291. Sharma JB, Singh N, Dharmendra S, et al. Six months versus nine months anti-tuberculous therapy for female genital tuberculosis: a randomized controlled trial. *Eur J Obstet Gynecol Reprod Biol*. 2016;203:264–273. doi:10.1016/j.ejogrb.2016.05.035.
292. Syed FF, Mayosi BM. A modern approach to tuberculous pericarditis. *Prog Cardiovasc Dis*. 2007;50(3):218–236. doi:10.1016/j.pcad.2007.03.002.
293. Cherian G, Habashy AG, Uthaman B, Cherian JM, Salama A, Anim JT. Detection and follow-up of mediastinal lymph node enlargement in tuberculous pericardial effusions using computed tomography. *Am J Med*. 2003;114(4):319–322. doi:10.1016/s0002-9343(02)01521-8.
294. Sun JS, Park KJ, Kang DK. CT findings in patients with pericardial effusion: differentiation of malignant and benign disease. *AJR Am J Roentgenol*. 2010;194(6):W489–94. doi:10.2214/AJR.09.2599.
295. Agner RC, Gallis HA. Pericarditis: differential diagnostic considerations. *Arch Intern Med*. 1979;139(4):407–412. doi:10.1001/archinte.139.4.407.
296. Wiysonge CS, Ntsekhe M, Thabane L, et al. Interventions for treating tuberculous pericarditis. *Cochrane Database Syst Rev*. 2017;9:CD000526. doi:10.1002/14651858.CD000526.pub2.

297. Reuter H, Burgess LJ, Schneider J, Van Vuuren W, Doubell AF. The role of histopathology in establishing the diagnosis of tuberculous pericardial effusions in the presence of HIV. *Histopathology*. 2006;48(3):295–302. doi:10.1111/j.1365-2559.2005.02320.x.
298. Tuon FF, Litvoc MN, Lopes MI. Adenosine deaminase and tuberculous pericarditis—a systematic review with meta-analysis. *Acta Trop*. 2006;99(1):67–74. doi:10.1016/j.actatropica.2006.07.004.
299. Strang JJ, Kakaza HH, Gibson DG, Girling DJ, Nunn AJ, Fox W. Controlled trial of prednisolone as adjuvant in treatment of tuberculosis constrictive pericarditis in Transkei. *Lancet*. 1987;2(8573):1418–1422. doi:10.1016/S0140-6736(87)91127-5.
300. Reuter H, Burgess LJ, Louw VJ, Doubell AF. The management of tuberculosis pericardial effusion: experience in 233 consecutive patients. *Cardiovasc J S Afr*. 2007;18(1):20–25.
301. Mayosi BM, Ntsekhe M, Bosch J, IMPI Trial Investigators, et al. Prednisolone and Mycobacterium indicus pranii in tuberculous pericarditis. *N Engl J Med*. 2014;371(12):1121–1130. doi:10.1056/NEJMoa1407380.
302. Gupta V, Shoughy SS, Mahajan S, et al. Clinics of ocular tuberculosis. *Ocul Immunol Inflamm*. 2015;23(1):14–24. doi:10.3109/09273948.2014.986582.
303. Conant MM, Vrasich CR, Wongsakhaluang JV, et al. Role of the Infectious Disease Consultant in Management of Patients With Tuberculosis-Associated Ocular Inflammation. *Open Forum Infect Dis*. 2016;3(1):ofv195. doi:10.1093/ofid/ofv195.
304. Patel SS, Saraiya NV, Tessler HH, Goldstein DA. Mycobacterial ocular inflammation: delay in diagnosis and other factors impacting morbidity. *JAMA Ophthalmol*. 2013;131(6):752–758. doi:10.1001/jamaophthalmol.2013.71.
305. Lynn WA, Lightman S. The eye in systemic infection. *Lancet*. 2004;364(9443):1439–1450. doi:10.1016/S0140-6736(04)17228-0.
306. Cunningham ET, Jr., Rathinam SR, Albin TA, Chee SP, Zierhut M. Tuberculous uveitis. *Ocul Immunol Inflamm*. 2015;23(1):2–6. doi:10.3109/09273948.2014.1004016.
307. Arora SK, Gupta V, Gupta A, Bamberg P, Kapoor GS, Sehgal S. Diagnostic efficacy of polymerase chain reaction in granulomatous uveitis. *Tuber Lung Dis*. 1999;79(4):229–233. doi:10.1054/tuld.1999.0210.
308. Ang M, Htoon HM, Chee SP. Diagnosis of tuberculous uveitis: clinical application of an interferon-gamma release assay. *Ophthalmology*. 2009;116(7):1391–1396. doi:10.1016/j.ophtha.2009.02.005.
309. Trad S, Bodaghi B, Saadoun D. Update on Immunological Test (Quantiferon-TB Gold) Contribution in the Management of Tuberculosis-Related Ocular Inflammation. *Ocul Immunol Inflamm*. 2018;26(8):1192–1199. doi:10.1080/09273948.2017.1332232.
310. Lee C, Agrawal R, Pavesio C. Ocular Tuberculosis-A Clinical Conundrum. *Ocul Immunol Inflamm*. 2016;24(2):237–242. doi:10.3109/09273948.2014.985387.
311. Figueira L, Fonseca S, Ladeira I, Duarte R. Ocular tuberculosis: Position paper on diagnosis and treatment management. *Rev Port Pneumol* (2006). 2017;23(1):31–38. doi:10.1016/j.rppnen.2016.10.004.
312. Alvarez GG, Roth VR, Hodge W. Ocular tuberculosis: diagnostic and treatment challenges. *Int J Infect Dis*. 2009;13(4):432–435. doi:10.1016/j.ijid.2008.09.018.
313. Gan WL, Jones NP. Serpiginous-like choroiditis as a marker for tuberculosis in a non-endemic area. *Br J Ophthalmol*. 2013;97(5):644–647. doi:10.1136/bjophthalmol-2012-302918.
314. Ang M, Hedayatfar A, Wong W, Chee SP. Duration of anti-tubercular therapy in uveitis associated with latent tuberculosis: a case-control study. *Br J Ophthalmol*. 2012;96(3):332–336. doi:10.1136/bjophthalmol-2011-300209.
315. Bansal R, Gupta A, Gupta V, Dogra MR, Bamberg P, Arora SK. Role of anti-tubercular therapy in uveitis with latent/manifest tuberculosis. *Am J Ophthalmol*. 2008;146(5):772–779. doi:10.1016/j.ajo.2008.06.011.
316. Agrawal R, Gupta B, Gonzalez-Lopez JJ, et al. The role of anti-tubercular therapy in patients with presumed ocular tuberculosis. *Ocul Immunol Inflamm*. 2015;23(1):40–46. doi:10.3109/09273948.2014.986584.
317. Sanghvi C, Bell C, Woodhead M, Hardy C, Jones N. Presumed tuberculous uveitis: diagnosis, management, and outcome. *Eye (Lond)*. 2011;25(4):475–480. doi:10.1038/eye.2010.235.
318. Lou SM, Montgomery PA, Larkin KL, Winthrop K, Zierhut M, Rosenbaum JT, Uveitis Specialists Study Group. Diagnosis and treatment for ocular tuberculosis among uveitis specialists: the international perspective. *Ocul Immunol Inflamm*. 2015;23(1):32–39. doi:10.3109/09273948.2014.994784.
319. Chen SC, Lin MC, Sheu SJ. Incidence and prognostic factor of ethambutol-related optic neuropathy: 10-year experience in southern Taiwan. *Kaohsiung J Med Sci*. 2015;31(7):358–362. doi:10.1016/j.kjms.2015.05.004.
320. van Zyl L, Du Plessis J, Viljoen J. Cutaneous tuberculosis overview and current treatment regimens. *Tuberculosis (Edinb)*. 2015;95(6):629–638. doi:10.1016/j.tube.2014.12.006.
321. Chen Q, Chen W, Hao F. Cutaneous tuberculosis: A great imitator. *Clin Dermatol*. 2019;37(3):192–199. doi:10.1016/j.clindermatol.2019.01.008.
322. Burgin S, Pomeranz MK, Orbuch P, et al. Mycobacteria and the skin. In: Rom WN, Garay S, eds. *Tuberculosis*. Toronto: Little, Brown and Company; 2004:593–608.
323. Delaval L, Goulenok T, Achouh P, et al. New insights on tuberculous aortitis. *J Vasc Surg*. 2017;66(1):209–215. doi:10.1016/j.jvs.2016.11.045.
324. Liu A, Nicol E, Hu Y, Coates A. Tuberculous endocarditis. *Int J Cardiol*. 2013;167(3):640–645. doi:10.1016/j.ijcard.2012.08.009.
325. Hakawi AM, Alrajhi AA. Tuberculosis of the bone marrow: clinico-pathological study of 22 cases from Saudi Arabia. *Int J Tuberc Lung Dis*. 2006;10(9):1041–1044.
326. Long R, Guzman R, Greenberg H, Safneck J, Hershfield E. Tuberculous mycotic aneurysm of the aorta: review of published medical and surgical experience. *Chest*. 1999;115(2):522–531. doi:10.1378/chest.115.2.522.
327. Agarwal R, Gupta D, Aggarwal AN, Behera D, Jindal SK. Experience with ARDS caused by tuberculosis in a respiratory intensive care unit. *Intensive Care Med*. 2005;31(9):1284–1287. doi:10.1007/s00134-005-2721-2.
328. Lerolle N, Laanani M, Galicier L, et al. Factors associated with tuberculosis-associated haemophagocytic syndrome: a multicentre case-control study. *Int J Tuberc Lung Dis*. 2020;24(1):124–130. doi:10.5588/ijtld.19.0856.
329. Girgis NI, Farid Z, Kilpatrick ME, Sultan Y, Mikhail IA. Dexamethasone adjunctive treatment for tuberculous meningitis. *Pediatr Infect Dis J*. 1991;10(3):179–183. doi:10.1097/00006454-199103000-00002.
330. Schoeman JF, Van Zyl LE, Laubscher JA, Donald PR. Effect of corticosteroids on intracranial pressure, computed tomographic findings, and clinical outcome in young children with tuberculous meningitis. *Pediatrics*. 1997;99(2):226–231. doi:10.1542/peds.99.2.226.
331. Dutt AK, Moers D, Stead WW. Short-course chemotherapy for extrapulmonary tuberculosis. Nine years' experience. *Ann Intern Med*. 1986;104(1):7–12. doi:10.7326/0003-4819-104-1-7.
332. Ramachandran S, Clifton IJ, Collyns TA, Watson JP, Pearson SB. The treatment of spinal tuberculosis: a retrospective study. *Int J Tuberc Lung Dis*. 2005;9(5):541–544.
333. Dian S, Yunivita V, Ganiem AR, et al. Double-Blind, Randomized, Placebo-Controlled Phase II Dose-Finding Study To Evaluate High-Dose Rifampin for Tuberculosis Meningitis. *Antimicrob Agents Chemother*. 2018;62(12). doi:10.1128/AAC.01014-18.
334. Yunivita V, Dian S, Ganiem AR, et al. Pharmacokinetics and safety/tolerability of higher oral and intravenous doses of rifampicin in adult tuberculous meningitis patients. *Int J Antimicrob Agents*. 2016;48(4):415–421. doi:10.1016/j.ijantimicag.2016.06.016.
335. Skutil V, Varsa J, Obsitnik M. Six-month chemotherapy for urogenital tuberculosis. *Eur Urol*. 1985;11(3):170–176. doi:10.1159/000472484.