

## ASSIGNMENT 1: [IFT6390]

LÉA RICARD & JOSEPH D. VIVIANO

### 1. PROBABILITIES

$P(D)$  = the probability of having disease.

$P(T)$  = the probability of having a positive test result.

- 1.5% of women in their 40s have breast cancer, therefore  $P(D) = 0.015$ .
- 87% true positive rate, therefore  $P(T|D) = 0.87$ .
- 9.6% false positive rate, therefore  $P(T|\neg D) = 0.096$ .

We want to know:

$$(1.1) \quad P(D|T) = \frac{P(T|D)P(D)}{P(T)}$$

Which means we need to calculate  $P(T)$  which is

$$(1.2) \quad P(T) = P(T|D)P(D) + P(T|\neg D)P(\neg D) = 0.87 \times 0.015 + 0.096 \times (1 - 0.015) \approx 0.1076$$

$$(1.3) \quad P(D|T) = \frac{0.87 \times 0.015}{0.1076} \approx 0.1213$$

### 2. CURSE OF DIMENSIONALITY

(A) Consider a hypercube in dimension  $d$  with side length  $c$ . What is the volume  $V$ ?

In the 2-dimensional case,  $\text{area} = c^2$ . In the 3 dimensional case,  $V = c^3$ . In the  $n$ -dimensional case,  $V = c^d$ .

(B)  $X$  is a random vector of dimension  $d$  ( $x \in d$ ) distributed uniformly within the hypercube (the probability density  $p(x) = 0$  for all  $x$  outside the cube). What is the probability density function  $p(x)$  for  $x$  inside the cube? Indicate which property(ies) of probability densities functions allow you to calculate this result.

For all probability distributions:

$$(2.1) \quad \int_{-\inf}^{\inf} p(x) = 1$$

We know  $p(x) = 0$  for all points outside of the hypercube. Therefore, the probability of being in a particular point inside the hypercube is

$$(2.2) \quad p(x) = \frac{1}{c^d}$$

(C) Consider the outer shell (border) of the hypercube of width 3% of  $c$  (covering the part of the hypercube extending from the faces of the cube and  $0.03c$  inwards). For example, if  $c = 100\text{cm}$ , the border will be  $3\text{cm}$  (left, right, top, etc ...) and will delimit this way a second (inner) hypercube of side  $100 - 3 - 3 = 94\text{cm}$ . If we generate a point  $x$  according to the previously defined probability distribution (by sampling), what is the probability that it falls in the border area? What is the probability that it falls in the smaller hypercube?

Let  $b$  be the amount to remove from the border on one side (i.e., left) of the outer hypercube.

$p(x_{large}) = 1$ . Therefore in the general case the probability we are in the smaller hypercube is:

$$(2.3) \quad p(x_{small}) = (c - 2b)^d / c^d$$

$$(2.4) \quad p(x_{border}) = 1 - p(x_{small})$$

Therefore for the above example:

$$(2.5) \quad p(x_{small}) = (100 - 2 \times 3)^d / 100^d = 94^d / 100^d$$

And as before:

$$(2.6) \quad p(x_{border}) = 1 - p(x_{small})$$

And the probability we are in the border is:

(D) Calculate the above for  $d = 1, 2, 3, 5, 10, 100, 1000$ .

$$(2.7) \quad 1 - 94^1 / 100^1 = 0.06$$

$$(2.8) \quad 94^2 / 100^2 = 0.1163$$

$$(2.9) \quad 94^3 / 100^3 = 0.1694$$

$$(2.10) \quad 94^5/100^5 = 0.2661$$

$$(2.11) \quad 94^{10}/100^{10} = 0.4614$$

$$(2.12) \quad 94^{100}/100^{100} = 0.9980$$

$$(2.13) \quad 94^{1000}/100^{1000} \approx 1$$

(E) When the dimension grows, the probability that  $x$  falls into the narrow border at the edge of the hypercube becomes more likely, which is contrary of our intuitions at lower dimensions.

### 3. PARAMETRIC GAUSSIAN VS PARZEN WINDOW DENSITY ESTIMATION

**3.1. Isotropic Gaussian Distribution.** (A) The named parameters are  $\mu$ , a  $d$ -long vector of means, and  $\Sigma$ , a  $d \times d$  covariance matrix, where  $n$  is the number of data points.

(B)

$$(3.1) \quad \mu = \frac{1}{n} \sum_{i=1}^n x_i.$$

$$(3.2) \quad \Sigma = \frac{1}{n} \sum_{i=1}^n (\mathbf{x}_i - \mu)(\mathbf{x}_i - \mu)^T$$

(C) For the  $\mu$  parameter, the algorithm complexity is in  $\mathcal{O}(n)$ , since it is summing over the vectors  $(x_i)$ .

For the  $\Sigma$  parameter, since the Gaussian density function is isotropic, the algorithm complexity is also in  $\mathcal{O}(n)$ .

(D)

$$(3.3) \quad p(x) = \frac{1}{(2\pi)^{\frac{d}{2}} \sqrt{|\Sigma|}} e^{-\frac{1}{2}(x-\mu)^T \Sigma^{-1} (x-\mu)}$$

(E) The cost of this operation is limited by the find the determinant of  $\Sigma$  and the inversion of  $\Sigma$ . Both of these operations assuming we use the Coppersmith–Winograd algorithm for the inversion and fast matrix multiplication for finding the determinant.

**3.2. Parzen windows with Isotropic Gaussian Kernels.** (A) If  $\sigma$  is fixed by the user, nothing is learned during training (the Gaussians are simply centered on each training data point and then summed to create a density). Basically this algorithm learn to remember the data.

(B)

$$(3.4) \quad p(x) = \frac{1}{n} \sum_{i=1}^n \mathcal{N}_{x_i, \sigma}(x)$$

Expanded becomes:

$$(3.5) \quad p(x) = \frac{1}{n} \sum_{i=1}^n \frac{1}{(2\pi)^{\frac{d}{2}} \sigma^d} e^{-\frac{1}{2} \frac{d(x_{test}, x_{train})^2}{\sigma^2}}$$

(C)  $\mathcal{O}(n)$ , since we have to calculate the distance between  $x_{test}$  and all of the  $n$  data points  $x_{train}$ .

**3.3. Capacity/Expressivity.** (A) The Parzen Gaussian is more expressive, because it can store information for every data point. The capacity of the algorithm grows as we give it more data points, this isn't true for the Gaussian distribution, which averages over all data points, so it has a fixed capacity for a given dimensionality, no matter how many training data the algorithm is shown.

(B) Parzen windows with Isotropic Gaussian Kernels, in the case that we used a large number of training examples with a small  $\sigma$  would result in extreme memorization of the noise in the training data (i.e., overfitting).

(C) Because in parametric Gaussian density estimation,  $\sigma$  is learned from the data, while it is fixed for all data points when using Parzen windows.

**3.4. Empirical Risk.** (A) *Express the equation of a diagonal Gaussian density in  $\mathbb{R}^d$ . Specify what are its parameters and their dimensions.*

$$p(x) = \frac{1}{(2\pi)^{\frac{d}{2}} \sqrt{|\Sigma|}} e^{-\frac{1}{2} (x-\mu)^T \Sigma^{-1} (x-\mu)}$$

– parameters are  $\Sigma()$  and  $\mu()$ .

(B) *Show that the components of a random vector following a diagonal Gaussian distribution are independent random variables.*

(C) *Using  $-\log p(x)$  as the loss, write down the equation corresponding to the empirical risk minimization on the training set  $D$  (in order to learn the parameters).*

*The log-likelihood function is:*

$$(3.6) \quad \log(P|\mu, \Sigma) = -\frac{N}{2} \log|\Sigma| - \frac{1}{2} \sum_{n=1}^N [(X_n - \mu)^T \Sigma^{-1} (X_n - \mu)] + c$$

Where  $c$  is a constant.

(D) Solve this equation analytically in order to obtain the optimal parameters.

First we find :

$$(3.7) \quad \frac{\partial \log p(X|\mu, \Sigma)}{\partial \mu} = \sum_{i=1}^N (X_i - \mu)^T \Sigma^{-1} \mathbf{1} = 0$$

$$(3.8) \quad 0 = N\mu - \sum_{i=1}^N X_i$$

#### 4. SOFTMAX ACTIVATION FUNCTION

UNIVERSITÉ DE MONTRÉAL

Email address: [joseph@viviano.ca](mailto:joseph@viviano.ca)