



MAX-PLANCK-GESELLSCHAFT

Technical University of Munich
Max Planck Institute of Quantum Optics

An atom-photon quantum gate using an atomic clock qubit and a birrefringent cavity

Leart Zuka

Thesis submitted within the M.Sc. Condensed Matter Physics

Supervisor: Prof. Dr. Gerhard Rempe
Second Examiner:

February 22, 2026

Abstract

Single atoms coupled to optical cavities provide a powerful platform for photonic quantum information processing. Minimizing the influence of external factors like magnetic field fluctuations is a key goal in this endeavor. In this work, I investigate a novel gate protocol that employs a single ^{87}Rb atom prepared in a magnetic-field-insensitive clock qubit and coupled to a high-finesse birefringent Fabry–Perot cavity. The cavity’s high birefringence creates two well-separated polarization eigenmodes, enabling polarization-selective reflectivity that depends on the atomic qubit state. This allows for the implementation of a controlled phase (CPHASE) gate between the atomic clock states and our photonic qubits. By rotating the basis of the photonic qubit, one is then able to realize an atom-photon controlled-NOT (CNOT) gate.

Show some results here and talk about how well it works and also how one might be able to use the gate

I present simulations based on input–output theory and master-equation modeling that quantify the conditional reflection amplitudes, gate truth table, and resulting fidelities under realistic experimental imperfections such as finite mode matching, cavity loss channels, and multiphoton contributions. Our results indicate that atom-state-dependent phase shifts on the photonic qubit are achievable in the current system, providing a viable path toward a robust, high-fidelity, cavity-assisted atom–photon quantum gate.

Contents

Abstract	1
1 Introduction	3
2 Theory	4
2.1 Cavity Quantum Electrodynamics	4
2.1.1 Jaynes-Cummings Model	4
2.1.2 Dissipation and Decay Channels	4
2.1.2.1 Atomic Decay Channels	5
2.1.2.2 Photonic Decay Channels	5
2.2 Reflection from a Cavity-Atom System	5
2.3 Conditional Reflection	5
2.3.1 Reflection in CNOT basis	5
A Analytical Derivations	6
A.1 Driving Hamiltonian	6
A.2 Analytical derivation for case of atom in $ 0\rangle$ and photon is in $ h\rangle$	7
A.2.1 Lindblad Formalism	8
A.2.2 Calculation of the Maxwell Bloch Equations (no lossy channels)	9
A.2.2.1 Jaynes Cummings Hamiltonian	9
A.2.3 Cavity mode a	10
A.2.4 Atomic lowering operator σ	11
A.2.5 Pauli-Z operator σ_z	12
Bibliography	14

Chapter 1

Introduction

Bla bla bla quantum gate important for distant computation between two nodes very cool. Cite some stuff from kimble and duan here [1], talk about recent advancements or stuff that was similar(cite reiserer here [2]).

Basically the idea here is to have a good introduction into the topic. Here I can then talk about:

- what makes up quantum computation - why it's so difficult to scale up in a certain way (maybe look at different examples with large amounts of ^{87}Rb atoms and comment on what the challenges there are when it comes to first of all having lots of them, and then keeping them coherent and stuff like that).
- Talk basically a lot about what Duan and Kimble talked about in their introduction of the paper since they were talking a lot about how computing at one node is hard, but splitting it up into smaller nodes and thus making it able to compute stuff.
- Then talk about how one is then able to connect the nodes with the gate approach.
- Explain what makes it unique (aka the birefringence of the cavity and the ability to then use the clock states which makes it insensitive to external magnetic fields)
- Give an outlook over the chapters that are to follow

Chapter 2

Theory

What follows is going to be the introduction and explanation of specific concepts required to understand the key points of this thesis.

2.1 Cavity Quantum Electrodynamics

At the heart of an atom-photon gate lies the interaction between the two components, which in the context of this thesis is achieved by placing an atom in a high-finesse Fabry-Perot cavity. The resulting dynamics then fall in the domain of Cavity Quantum Electrodynamics (CQED). Placing the atom between two highly-reflective mirrors of an optical cavity increases the amount of times a photon that is inside of a cavity bounces back and forth. This effectively increases the absorption cross section by the number of round-trips the photon takes, allowing for a strong light-matter interaction.

The theoretical model which explains this interaction in a fully quantum mechanical picture is called the *Jaynes-Cummings Model* [3] and was named after Edwin Jaynes and Fred Cummings who developed it in 1963.

2.1.1 Jaynes-Cummings Model

- Basic setup is atom with 3 levels $|u\rangle$, $|c\rangle$, $|e\rangle$ and cavity with single mode that is resonant to $|c\rangle \leftrightarrow |e\rangle$, with detuning between resonant transition and cavity, include picture from [4] and cite ofc
- Write down hamiltonian that includes first the basic structure and then explain rotating wave approximation (possibly show derivation in appendix)
- Show that the solutions of the eigenstates have this splitting term leading to the jaynes cummings ladder where at zero detuning (aka resonance) levels are seperated by $2g$

2.1.2 Dissipation and Decay Channels

Now we can include some losses here and there which are important to take into account as they reduce fidelity

Will proly scrap this section and rather talk about the cavities and their birefringence

Should proly just include these two subsubsections in the one before to save space and just include extra stuff that I'm considering like mode matching, extra dump state, and whatnot

2.1.2.1 Atomic Decay Channels

2.1.2.2 Photonic Decay Channels

2.2 Reflection from a Cavity-Atom System

- Different cases for reflection and thus transmission when driving with weak coherent pulse, based on state of the atom and detunings and stuff like that
- system was introduced by gardiner and collet [5] show equation and then say how you can solve for analytical solution for photonic field reflection $r(\omega)$ and then show the equation for the coupled and uncoupled case and maybe already show theoretical plots that compare the two both in reflectivity and phase shift.

2.3 Conditional Reflection

- Go off what Kimble and Duan were saying in [1] and how you get a phase shift which can also be seen in the plot when plotting the phase shift across different detunings and explain where it's coming from
- Say that this can be used to build a controlled phase flip gate when photon is in eigenbasis of the cavity with the atom being the controlling part and the photon being the target part

2.3.1 Reflection in CNOT basis

- Show that by turning the basis and instead of being in the polarization eigenbasis of the cavity, one can build a CNOT gate which is much more interesting for quantum computation by just sending a superposition of the two eigenbases of the cavity.

Appendix A

Analytical Derivations

A.1 Driving Hamiltonian

Since we're mainly going to be working with the *mesolve* function in QuTip to simulate the dynamics of the atom-photon gate it's imperative to know what each and every part does. Next to the Jaynes-Cummings-Hamiltonian the most important part is the one explaining the *photon that gets reflected from the cavity*, most notably the part before it gets reflected aka the *driving Hamiltonian* H_d . For this we're going to just write down once, the dynamics of a coherent driving field which will be the foundation of my photon that shall or shall not enter the cavity.

The photon is going to come from a laser, that is going to be described as a classical oscillating electric field. That light field has an amplitude E_0 and a frequency ω_d :

$$E(t) = E_0 \cos(\omega_d t)$$

Using the euler formula we can rewrite this to:

$$E(t) = \frac{E_0}{2} (e^{i\omega_d t} + e^{-i\omega_d t})$$

The light field operator $\hat{E}$ interacts with the light in our cavity (which get described by our creation and annihilation operators) linearly:

$$\hat{E} \propto (a + a^\dagger)$$

When now describing the *driving Hamiltonian* what we're doing is just overlapping the light field operator with the oscillating electric field:

$$\begin{aligned} H_{drive} &\propto E(t) \cdot (a + a^\dagger) \\ H_{drive} &= (e^{i\omega_d t} + e^{-i\omega_d t}) \cdot (a + a^\dagger) \end{aligned}$$

From this we get four terms where two cancel out as they aren't energy conserving rendering us with the final expression for our driving Hamiltonian:

$$H_{drive} = a^\dagger e^{-i\omega_d t} + a e^{+i\omega_d t}$$

Where the first term adds a photon to our cavity and the other removes one. Sometimes you'll see a minus instead of a plus between the two terms, this is just due to different conventions of the light field, where people chose a sin to describe the oscillating part instead of a cos, but the physics remain the same.

A.2 Analytical derivation for case of atom in $|0\rangle$ and photon is in $|h\rangle$

Our Hamiltonian is as follows:

$$H = \hbar g(|e\rangle \langle 1| a_h + |1\rangle \langle e| a_h^\dagger)$$

We drive the a_h mode of our cavity through $a_h^{in}(t)$, thus giving us the following equation of motion for our a_h cavity mode:

$$\dot{a}_h = -i[a_h, H] - (i\Delta + \frac{\kappa}{2})a_h - \sqrt{\kappa}a_h^{in}(t)$$

Because a_h commutes with H , the commutator is 0 and we get

$$\rightarrow \dot{a}_h = -(i\Delta + \frac{\kappa}{2})a_h - \sqrt{\kappa}a_h^{in}(t)$$

We now drop the subscript h as we'll only be looking at that case.

Multiplying with $e^{i\omega t}$, and integrating we get:

$$\int dt e^{i\omega t} \dot{a}(t) = \underbrace{[e^{i\omega t} a(t)]_{-\infty}^{\infty}}_{=0} - \underbrace{i\omega \int dt e^{i\omega t} a(t)}_{=i\omega a(\omega)}$$

where in the last expression we used that the integral performs a Fourier transformation.

$$\begin{aligned} \Rightarrow -i\omega a(\omega) &= -(i\Delta + \frac{\kappa}{2})a(\omega) - \sqrt{\kappa}a^{in}(\omega) \\ \sqrt{\kappa}a^{in}(\omega) &= -(i\Delta + \frac{\kappa}{2} - i\omega)a(\omega) \\ \rightarrow a(\omega) &= -\frac{\sqrt{\kappa}}{i\Delta + \frac{\kappa}{2} - i\omega}a^{in}(\omega) \end{aligned}$$

Now from our regular input output relation

$$a^{out}(\omega) = a^{in}(\omega) + \kappa a(\omega)$$

We thus get:

$$\begin{aligned} a^{out}(\omega) &= a^{in}(\omega) - \frac{\sqrt{\kappa}}{i\Delta + \frac{\kappa}{2} - i\omega}a^{in}(\omega) \\ a^{out}(\omega) &= (1 - \frac{\kappa}{i\Delta + \frac{\kappa}{2} - i\omega})a^{in}(\omega) \\ a^{out}(\omega) &= (1 - \frac{i\Delta + \frac{\kappa}{2} - i\omega - \kappa}{i\Delta + \frac{\kappa}{2} - i\omega})a^{in}(\omega) \\ a^{out}(\omega) &= (\frac{i\Delta - \frac{\kappa}{2} - i\omega}{i\Delta + \frac{\kappa}{2} - i\omega})a^{in}(\omega) \\ &= \underbrace{\frac{i(\Delta - \omega) - \frac{\kappa}{2}}{i(\Delta - \omega) + \frac{\kappa}{2}}}_{h(\omega)}a^{in}(\omega) \end{aligned}$$

We can then use the convolution theorem in order to solve for $a^{out}(t)$

$$a^{out}(t) = \int_{-\infty}^{\infty} d\tau \underbrace{h(t-\tau)}_{h(t) = \frac{1}{2\pi} \int_{-\infty}^{\infty} d\omega e^{-i\omega t} h(\omega)} a^{in}(\tau)$$

Now using that $h(\omega) = 1 - \frac{\kappa}{i(\Delta - \omega) + \frac{\kappa}{2}}$ we get for $h(t)$ (via partial integration):

$$\begin{aligned} h(t) &= \delta(t) - \kappa e^{-i\delta t} e^{-\frac{\kappa}{2}t} \theta(t) \\ \Rightarrow a^{out}(t) &= a^{in}(t) - \kappa \int_{-\infty}^{\infty} d\tau e^{-i\delta(t-\tau)} e^{-\frac{\kappa}{2}(t-\tau)} a^{in}(\tau) \end{aligned}$$

For slowly varying pulses we can now approximate that $a^{in}(\tau) \approx a^{in}(t)$ thus giving us:

$$\begin{aligned} \Rightarrow a^{out}(t) &\approx a^{in}(t) - \frac{\kappa}{i\Delta + \frac{\kappa}{2}} a^{in}(t) \\ &= \frac{i\Delta - \frac{\kappa}{2}}{i\Delta + \frac{\kappa}{2}} a^{in}(t) \end{aligned}$$

Which for no detuning (so $\Delta = 0$) gives us:

$$a^{out}(t) = -a^{in}(t) \quad (\text{A.1})$$

A.2.1 Lindblad Formalism

We start from the master equation:

$$\dot{\rho} = -\frac{i}{\hbar} [H, \rho] + \sum_j \gamma_j \left(L_j \rho L_j^\dagger - \frac{1}{2} \{L_j^\dagger L_j, \rho\} \right), \quad (\text{A.2})$$

Now, for an operator O ,

$$\frac{d}{dt} \langle O \rangle = \text{Tr} (O \dot{\rho}) \quad (\text{A.3})$$

$$= \text{Tr} \left(O \left[-\frac{i}{\hbar} [H, \rho] + \sum_j \gamma_j \left(L_j \rho L_j^\dagger - \frac{1}{2} \{L_j^\dagger L_j, \rho\} \right) \right] \right). \quad (\text{A.4})$$

The Hamiltonian contribution gives

$$- \text{Tr} \left(O \frac{i}{\hbar} [H, \rho] \right) = -\frac{i}{\hbar} \text{Tr} (OH\rho - O\rho H) = -\frac{i}{\hbar} \text{Tr} ([O, H]\rho) = -\frac{i}{\hbar} \langle [O, H] \rangle. \quad (\text{A.5})$$

For the dissipative terms we obtain

$$\sum_j \gamma_j \text{Tr} \left(OL_j \rho L_j^\dagger - \frac{1}{2} OL_j^\dagger L_j \rho - \frac{1}{2} O \rho L_j^\dagger L_j \right). \quad (\text{A.6})$$

Using cyclicity of the trace:

$$= \sum_j \gamma_j \left(\langle L_j^\dagger O L_j \rangle - \frac{1}{2} \langle \{L_j^\dagger L_j, O\} \rangle \right). \quad (\text{A.7})$$

Therefore the general equation of motion is

$$\frac{d}{dt} \langle O \rangle = -\frac{i}{\hbar} \langle [O, H] \rangle + \sum_j \gamma_j \left(\langle L_j^\dagger O L_j \rangle - \frac{1}{2} \langle \{L_j^\dagger L_j, O\} \rangle \right). \quad (\text{A.8})$$

A.2.2 Calculation of the Maxwell Bloch Equations (no lossy channels)

A.2.2.1 Jaynes Cummings Hamiltonian

Assuming ground state has zero energy our Jaynes-Cummings Hamiltonian is equal to:

$$H_{JC,S} = \underbrace{\omega_c a^\dagger a + \omega_a \sigma^\dagger \sigma}_{=H_{0,S}} + \underbrace{g(\sigma + \sigma^\dagger)(a + a^\dagger)}_{=H_{int,S}} + H_{drive,S} \quad (\text{A.9})$$

where σ is the atomic lowering operator, ω_a the atomic transition frequency between the levels $|1\rangle$ and $|e\rangle$, a the photonic destruction operator, and ω_a the cavity mode resonance frequency.

H_{drive} in this case is the external light field which is driving our system, which in the semiclassical formalism has the following form [6]:

$$H_{drive,S} = i(\epsilon e^{-i\omega_d t} + \epsilon^* e^{i\omega_d t})(a + a^\dagger) \quad (\text{A.10})$$

Here, a again is the photonic destruction operator, ω_d is the frequency of the driving field, and ϵ is the complex amplitude of our incoming pulse.

In general going from the Schrödinger picture to the interaction picture, is achieved by converting our operators into the interaction picture via following equation:

$$i \frac{dA_I}{dt} = [A_I(t), H_{0,S}] \quad (\text{A.11})$$

Furthermore, $H_{0,I} = H_{0,S}$. Equipped with this we can now calculate the expressions for operators in the interaction picture:

$$i \frac{da_I(t)}{dt} = [a_I(t), H_{0,S}] \quad (\text{A.12})$$

$$= [a_I(t), \omega_c a^\dagger a] \quad (\text{A.13})$$

$$= \omega_c a_I(t) \quad (\text{A.14})$$

$$\Rightarrow \frac{da_I(t)}{dt} = -i\omega_c a_I(t) \rightarrow a_I(t) = a e^{-i\omega_c t} \quad (\text{A.15})$$

Where we used that:

$$[a, H_{0,S}] = \omega_c [a, a^\dagger a] = \omega_c (\underbrace{[a, a^\dagger]}_{=1} a + a^\dagger \underbrace{[a, a]}_{=0}) = \omega_c a \quad (\text{A.16})$$

Now we repeat the same thing for the atomic lowering operator σ :

$$i \frac{d\sigma_I(t)}{dt} = [\sigma_I(t), H_{0,S}] \quad (\text{A.17})$$

$$i \frac{d\sigma_I(t)}{dt} = \omega_a \sigma_I(t) \quad (\text{A.18})$$

$$\Rightarrow \frac{d\sigma_I(t)}{dt} = -i\omega_a \sigma_I(t) \rightarrow \sigma_I(t) = \sigma e^{-i\omega_a t} \quad (\text{A.19})$$

Now we can calculate the Hamiltonian for the interaction part and the drive part, in the interaction picture, by just plugging these operators in there, giving us:

$$H_{int,I} = \sigma a e^{-i(\omega_a - \omega_c)t} + \sigma a^\dagger e^{-i(\omega_a - \omega_c)t} + \sigma^\dagger a e^{-i(\omega_c - \omega_a)t} + \sigma^\dagger a^\dagger e^{i(\omega_c + \omega_a)t} \quad (\text{A.20})$$

$$\xrightarrow{RWA} \sigma a^\dagger e^{-i(\omega_a - \omega_c)t} + \sigma^\dagger a e^{-i(\omega_c - \omega_a)t} \quad (\text{A.21})$$

$$H_{drive,I} = i(\epsilon e^{-i\omega_d t} + \epsilon^\dagger e^{i\omega_d t})(a_I(t) + a_I^\dagger(t)) \quad (\text{A.22})$$

$$\xrightarrow{RWA} i(a\epsilon^* e^{-i(\omega_c - \omega_d)t} + a^\dagger \epsilon e^{i(\omega_c - \omega_d)t}) \quad (\text{A.23})$$

Thus giving us the total Hamiltonian in the interaction picture under the rotating wave approximation:

$$H_{JC,I} \stackrel{\text{RWA}}{=} \omega_c a^\dagger a + \omega_a \sigma^\dagger \sigma + g(\sigma a^\dagger e^{-i(\omega_a - \omega_c)t} + \sigma^\dagger a e^{-i(\omega_c - \omega_a)t}) + i(a\epsilon^* e^{-i(\omega_c - \omega_d)t} + a^\dagger \epsilon e^{i(\omega_c - \omega_d)t}) \quad (\text{A.24})$$

If we now transform back into the Schrödinger picture, we get:

$$H_{JC,S} \stackrel{\text{RWA}}{=} \omega_c a^\dagger a + \omega_a \sigma^\dagger \sigma + g(a^\dagger \sigma + a \sigma^\dagger) + i(a\epsilon^* e^{i\omega_d t} + a^\dagger \epsilon e^{-i\omega_d t}) \quad (\text{A.25})$$

If we now change into the rotating frame of the driving field via

$$U(t) = \exp\{i\omega_d t(a^\dagger a + \sigma^\dagger \sigma)\} \quad (\text{A.26})$$

$$H \rightarrow U H U^\dagger + i\dot{U} U^\dagger =: H_{RF} \quad (\text{A.27})$$

we then get the final form of our Hamiltonian:

$$H_{JC,RF} = \underbrace{(\omega_c - \omega_d)}_{=\Delta_c} a^\dagger a + \underbrace{(\omega_a - \omega_d)}_{=\Delta_a} \sigma^\dagger \sigma + g(a^\dagger \sigma + a \sigma^\dagger) + i(a\epsilon^* + a^\dagger \epsilon) \quad (\text{A.28})$$

Which gives rise to 4 scenarios which I'll be covering with my project:

scenario	meaning
$\Delta_a = \Delta_c = 0$	Atom is in coupled state (here: $ 1\rangle$) and we drive with a light field that's resonant to the cavity (here: we drive $ 1\rangle \leftrightarrow e\rangle$ which would be our $ \pi\rangle$ polarized light)
$\Delta_a = 0; \Delta_c \neq 0$	Atom is in coupled state but we drive with light that's non-resonant to cavity (here: $ V\rangle$ polarized light)
$\Delta_a \neq 0; \Delta_c = 0$	Atom is in uncoupled state (here: $ 0\rangle$) and we drive with a light field that's resonant to the cavity
$\Delta_a, \Delta_c \neq 0$	Atom is in uncoupled state and we drive with light that's non-resonant to cavity

In order to improve readability we'll be splitting the full JC Hamiltonian into three parts:

$$H_0 = \Delta_c a^\dagger a + \Delta_a \sigma^\dagger \sigma \quad (\text{A.29})$$

$$H_{int} = g(a^\dagger \sigma + a \sigma^\dagger) \quad (\text{A.30})$$

$$H_{drive} = i(a\epsilon^* + a^\dagger \epsilon) \quad (\text{A.31})$$

with following losses:

$$\gamma_1 = \gamma, \quad L_1 = \sigma, \quad (\text{A.32})$$

$$\gamma_2 = \kappa, \quad L_2 = a. \quad (\text{A.33})$$

A.2.3 Cavity mode a

For H_0 :

$$[a, H_0] = \Delta_c [a, a^\dagger a] + \Delta_c a \underbrace{[a, \sigma^\dagger \sigma]}_{=0} \quad (\text{A.34})$$

$$= \Delta_c (\underbrace{[a, a^\dagger]}_{=1} a + a^\dagger \underbrace{[a, a]}_{=0}) \quad (\text{A.35})$$

$$= \Delta_c a \quad (\text{A.36})$$

$$\Rightarrow -i\langle [a, H_0] \rangle = -i\Delta_c \langle a \rangle \quad (\text{A.37})$$

For H_{int} :

$$[a, H_{int}] = g([a, a^\dagger \sigma] + [a, a \sigma^\dagger]) \quad (\text{A.38})$$

$$= g(a a^\dagger \sigma - a^\dagger a \sigma + \cancel{a a \sigma^\dagger} - \cancel{a a \sigma^\dagger}) \quad (\text{A.39})$$

$$\rightarrow \langle a a^\dagger \sigma \rangle = \langle n | a a^\dagger \sigma | n \rangle = (n+1) \langle \sigma \rangle \quad (\text{A.40})$$

$$\rightarrow \langle a^\dagger a \sigma \rangle = \langle n | a^\dagger a \sigma | n \rangle = n \langle \sigma \rangle \quad (\text{A.41})$$

$$\Rightarrow -i\langle [a, H_{int}] \rangle = -ig\langle \sigma \rangle \quad (\text{A.42})$$

For H_{drive} :

$$[a, H_{drive}] = [a, i(a\epsilon^* + a^\dagger\epsilon)] \quad (\text{A.43})$$

$$= i([a, a\epsilon^*] + [a, a^\dagger\epsilon]) \quad (\text{A.44})$$

$$= i(0 + \epsilon) \quad (\text{A.45})$$

$$= i\epsilon \quad (\text{A.46})$$

$$\Rightarrow -i\langle [a, H_{drive}] \rangle = i\langle \epsilon \rangle \stackrel{\epsilon \in \mathbb{C}}{=} i\epsilon \quad (\text{A.47})$$

If we now include the fact that:

$$i\epsilon \hat{=} \sqrt{\kappa_{oc}} a^{in}(t) \quad (\text{A.48})$$

we are then left with:

$$\Rightarrow -i\langle [a, H_{drive}] \rangle = -\sqrt{\kappa_{oc}} a^{in}(t) \quad (\text{A.49})$$

For rest of the Lindbladian:

$$D(a) = a^\dagger aa - \frac{1}{2}\{a^\dagger a, a\} \quad (\text{A.50})$$

$$= a^\dagger aa - \frac{1}{2}(a^\dagger aa - aa^\dagger a) \quad (\text{A.51})$$

$$= \frac{1}{2}a^\dagger aa - \frac{1}{2}aa^\dagger a \quad (\text{A.52})$$

$$= \frac{1}{2}(a^\dagger aa - aa^\dagger a) \quad (\text{A.53})$$

$$= \frac{1}{2}([a^\dagger, a]a) \quad (\text{A.54})$$

$$= \frac{1}{2}(-a) \quad (\text{A.55})$$

$$= -\frac{1}{2}a \quad (\text{A.56})$$

$$\Rightarrow \kappa\langle D(a) \rangle = -\frac{\kappa}{2}\langle a \rangle \quad (\text{A.57})$$

So for $\langle a \rangle(t)$ we get:

$$\begin{aligned} \frac{d\langle a \rangle}{dt} &= -i\Delta_c\langle a \rangle - ig\langle \sigma \rangle - \sqrt{\kappa_{oc}} a_{in}(t) - \frac{\kappa}{2}\langle a \rangle \\ &= -(i\Delta_c + \frac{\kappa}{2})\langle a \rangle - ig\langle \sigma \rangle - \sqrt{\kappa_{oc}} a_{in}(t) \end{aligned} \quad (\text{A.58})$$

A.2.4 Atomic lowering operator σ

For H_0 :

$$[\sigma, H_0] = \Delta_a \underbrace{[\sigma, \sigma^\dagger \sigma]} \quad (\text{A.59})$$

$$= \underbrace{[\sigma, \sigma^\dagger]}_{=1} \sigma + \sigma^\dagger \underbrace{[\sigma, \sigma]}_{=0}$$

$$= \Delta_a \mathbb{1}\sigma \quad (\text{A.60})$$

$$\Rightarrow -i\langle [\sigma, H_0] \rangle = -i\Delta_a\langle \sigma \rangle \quad (\text{A.61})$$

For H_{int} :

$$[\sigma, H_{int}] = g([\sigma, a^\dagger \sigma] + [\sigma, a \sigma^\dagger]) \quad (\text{A.62})$$

$$= g(\cancel{a^\dagger \sigma \sigma} - \cancel{a^\dagger \sigma \sigma} + a \sigma \sigma^\dagger - a \sigma^\dagger \sigma) \quad (\text{A.63})$$

$$= g(a \sigma \sigma^\dagger - a \sigma^\dagger \sigma) \quad (\text{A.64})$$

Now we can consider that:

$$\sigma^\dagger \sigma = |e\rangle \langle 1| \langle e| = |e\rangle \langle e| \quad (\text{A.65})$$

$$\sigma \sigma^\dagger = |1\rangle \langle e| \langle 1| = |1\rangle \langle 1| \quad (\text{A.66})$$

$$[\sigma, H_{int}] = g(a \sigma \sigma^\dagger - a \sigma^\dagger \sigma) \quad (\text{A.67})$$

$$= ga(|1\rangle \langle 1| - |e\rangle \langle e|) \quad (\text{A.68})$$

$$= ga \sigma_z \quad (\text{A.69})$$

$$\Rightarrow -i\langle [\sigma, H_{int}] \rangle = -ig\langle a \sigma_z \rangle \quad (\text{A.70})$$

No interaction with H_{drive} .

For rest of Lindbladian:

$$D(\sigma) = \sigma^\dagger \underbrace{\sigma \sigma}_{=0} - \frac{1}{2} \{ \sigma^\dagger \sigma, \sigma \} \quad (\text{A.71})$$

$$= -\frac{1}{2} \underbrace{\sigma^\dagger \sigma \sigma}_{=0} - \frac{1}{2} \sigma \sigma^\dagger \sigma \quad (\text{A.72})$$

$$= -\frac{1}{2} \sigma \quad (\text{A.73})$$

$$\gamma \langle D(\sigma) \rangle = -\frac{\gamma}{2} \langle \sigma \rangle \quad (\text{A.74})$$

So for $\langle \sigma \rangle(t)$ we get:

$$\begin{aligned} \frac{d\langle \sigma \rangle}{dt} &= -i\Delta_a \langle \sigma \rangle - ig\langle a \sigma_z \rangle - \frac{\gamma}{2} \langle \sigma \rangle \\ &= -(i\Delta_a + \frac{\gamma}{2}) \langle \sigma \rangle - ig\langle a \sigma_z \rangle \end{aligned} \quad (\text{A.75})$$

A.2.5 Pauli-Z operator σ_z

For H_0 :

$$[\sigma_z, H_0] = \Delta_a [\sigma_z, \sigma^\dagger \sigma] \quad (\text{A.76})$$

$$= \Delta_a [\sigma_z, |e\rangle \langle e|] \quad (\text{A.77})$$

$$= \Delta_a (\sigma_z |e\rangle \langle e| - |e\rangle \langle e| \sigma_z) \quad (\text{A.78})$$

$$= \Delta_a (-|e\rangle \langle e| + |e\rangle \langle e|) \quad (\text{A.79})$$

$$= 0 \quad (\text{A.80})$$

For H_{int} :

$$[\sigma_z, H_{int}] = g([\sigma_z, a^\dagger \sigma] + [\sigma_z, a \sigma^\dagger]) \quad (\text{A.81})$$

$$= g(a^\dagger \sigma_z \sigma - a^\dagger \sigma \sigma_z + a \sigma_z \sigma^\dagger - a \sigma^\dagger \sigma_z) \quad (\text{A.82})$$

$$= g(a^\dagger \sigma + a^\dagger \sigma - a \sigma^\dagger - a \sigma^\dagger) \quad (\text{A.83})$$

$$= 2g(a^\dagger \sigma - a \sigma^\dagger) \quad (\text{A.84})$$

$$\Rightarrow -i\langle[\sigma_z, H_{int}]\rangle = -2ig(\langle a^\dagger \sigma \rangle - \langle a \sigma^\dagger \rangle) \quad (\text{A.85})$$

No interaction with H_{drive} . For rest of Lindbladian:

$$D(\sigma_z) = \sigma^\dagger \sigma_z \sigma - \frac{1}{2} \{ \sigma^\dagger \sigma, \sigma \} \quad (\text{A.86})$$

$$= |e\rangle \langle e| - \frac{1}{2} \underbrace{\sigma^\dagger \sigma \sigma_z}_{=-|e\rangle \langle e|} - \frac{1}{2} \underbrace{\sigma_z \sigma^\dagger \sigma}_{=-|e\rangle \langle e|} \quad (\text{A.87})$$

$$= |e\rangle \langle e| + |e\rangle \langle e| \quad (\text{A.88})$$

$$= 2|e\rangle \langle e| \quad (\text{A.89})$$

$$= \mathbb{1} - \sigma_z \quad (\text{A.90})$$

$$\Rightarrow \gamma \langle D(\sigma_z) \rangle = \gamma(1 - \langle \sigma_z \rangle) \quad (\text{A.91})$$

So for $\langle \sigma_z \rangle(t)$ we get:

$$\frac{d\langle \sigma_z \rangle(t)}{dt} = -2ig(\langle a^\dagger \sigma \rangle - \langle a \sigma^\dagger \rangle) + \gamma(1 - \langle \sigma_z \rangle) \quad (\text{A.92})$$

The final Maxwell Bloch Equations then take the form of:

$$\frac{d\langle a \rangle}{dt} = -(i\Delta_c + \frac{\kappa}{2})\langle a \rangle - ig\langle \sigma \rangle - \sqrt{\kappa_{oc}} a_{in}(t) \quad (\text{A.93})$$

$$\frac{d\langle \sigma \rangle}{dt} = -(i\Delta_a + \frac{\gamma}{2})\langle \sigma \rangle - ig\langle a \sigma_z \rangle \quad (\text{A.94})$$

$$\frac{d\langle \sigma_z \rangle}{dt} = -2ig(\langle a^\dagger \sigma \rangle - \langle a \sigma^\dagger \rangle) + \gamma(1 - \langle \sigma_z \rangle) \quad (\text{A.95})$$

Bibliography

- [1] L.-M. Duan and H. J. Kimble. “Scalable Photonic Quantum Computation through Cavity-Assisted Interactions”. In: *Phys. Rev. Lett.* 92 (12 Mar. 2004), p. 127902. DOI: 10.1103/PhysRevLett.92.127902. URL: <https://link.aps.org/doi/10.1103/PhysRevLett.92.127902>.
- [2] Andreas Reiserer et al. “A quantum gate between a flying optical photon and a single trapped atom”. In: *Nature* 508 (7495 Apr. 2014), pp. 237–240. ISSN: 14764687. DOI: 10.1038/nature13177. URL: <https://www.nature.com/articles/nature13177>.
- [3] E.T. Jaynes and F.W. Cummings. “Comparison of quantum and semiclassical radiation theories with application to the beam maser”. In: *Proceedings of the IEEE* 51.1 (1963), pp. 89–109. DOI: 10.1109/PROC.1963.1664.
- [4] Andreas Reiserer and Gerhard Rempe. “Cavity-based quantum networks with single atoms and optical photons”. In: *Rev. Mod. Phys.* 87 (4 Dec. 2015), pp. 1379–1418. DOI: 10.1103/RevModPhys.87.1379. URL: <https://link.aps.org/doi/10.1103/RevModPhys.87.1379>.
- [5] C. W. Gardiner and M. J. Collett. “Input and output in damped quantum systems: Quantum stochastic differential equations and the master equation”. In: *Phys. Rev. A* 31 (6 June 1985), pp. 3761–3774. DOI: 10.1103/PhysRevA.31.3761. URL: <https://link.aps.org/doi/10.1103/PhysRevA.31.3761>.
- [6] D. F. Walls and G. J. Milburn. *Quantum Optics*. 2nd. Berlin, Heidelberg: Springer, 2008. ISBN: 978-3-540-28574-8. DOI: 10.1007/978-3-540-28574-8.

Selbständigkeitserklärung

Ich versichere hiermit, dass ich die von mir eingereichte Abschlussarbeit selbstständig verfasst und keine anderen als die angegebenen Quellen und Hilfsmittel benutzt habe.

gezeichnet,
Leart Zuka, München den TBD.