

References

- Ding, Y., Fu, Y., Lee, J. C., & Hooper, D. C. (2012). Staphylococcus aureus NorD, a putative efflux pump coregulated with the Opp1 oligopeptide permease, contributes selectively to fitness in vivo. *Journal of Bacteriology*, 194(23), 6586–6593. <https://doi.org/10.1128/JB.01414-12>
- Fraunholz, M., Bernhardt, J., Schuldes, J., Daniel, R., Hecker, M., & Sinha, B. (2013). Complete Genome Sequence of Staphylococcus aureus 6850, a Highly Cytotoxic and Clinically Virulent Methicillin-Sensitive Strain with Distant Relatedness to Prototype Strains. *Genome Announcements*, 1(5), e00775-13. <https://doi.org/10.1128/genomeA.00775-13>
- Hiron, A., Borezée-Durant, E., Piard, J.-C., & Juillard, V. (2007). Only one of four oligopeptide transport systems mediates nitrogen nutrition in Staphylococcus aureus. *Journal of Bacteriology*, 189(14), 5119–5129. <https://doi.org/10.1128/JB.00274-07>
- Lehman, M. K., Nuxoll, A. S., Yamada, K. J., Kielian, T., Carson, S. D., & Fey, P. D. (2019). Protease-Mediated Growth of Staphylococcus aureus on Host Proteins Is opp3 Dependent. *mBio*, 10(2), e02553-18. <https://doi.org/10.1128/mBio.02553-18>
- Nguyen, A. T., & Oglesby-Sherrouse, A. G. (2016). Interactions between Pseudomonas aeruginosa and Staphylococcus aureus during co-cultivations and polymicrobial infections. *Applied Microbiology and Biotechnology*, 100(14), 6141–6148. <https://doi.org/10.1007/s00253-016-7596-3>
- Niggli, S., Schwyter, L., Poveda, L., Grossmann, J., & Kümmerli, R. (2023). Rapid and strain-specific resistance evolution of Staphylococcus aureus against inhibitory molecules secreted by Pseudomonas aeruginosa. *mBio*, 14(5), e03153-22. <https://doi.org/10.1128/mbio.03153-22>
- Queck, S. Y., Jameson-Lee, M., Villaruz, A. E., Bach, T.-H. L., Khan, B. A., Sturdevant, D. E., Ricklefs, S. M., Li, M., & Otto, M. (2008). RNAIII-Independent Target Gene Control by the agr Quorum-Sensing System: Insight into the Evolution of Virulence Regulation in

Staphylococcus aureus. *Molecular Cell*, 32(1), 150–158.

<https://doi.org/10.1016/j.molcel.2008.08.005>

Remy, L., Carrière, M., Derré-Bobillot, A., Martini, C., Sanguinetti, M., & Borezée-Durant, E. (2013). The *Staphylococcus aureus* Opp1 ABC transporter imports nickel and cobalt in zinc-depleted conditions and contributes to virulence. *Molecular Microbiology*, 87(4), 730–743. <https://doi.org/10.1111/mmi.12126>

Schoenfelder, S. M. K., Marincola, G., Geiger, T., Goerke, C., Wolz, C., & Ziebuhr, W. (2013). Methionine Biosynthesis in *Staphylococcus aureus* Is Tightly Controlled by a Hierarchical Network Involving an Initiator tRNA-Specific T-box Riboswitch. *PLoS Pathogens*, 9(9), e1003606. <https://doi.org/10.1371/journal.ppat.1003606>

Schwyster, L., Niggli, S., Zajac, N., Poveda, L., Wolski, W. E., Panse, C., Schlapbach, R., Kümmerli, R., & Grossmann, J. (2025). A combination of phenotypic responses and genetic adaptations enables *Staphylococcus aureus* to withstand inhibitory molecules secreted by *Pseudomonas aeruginosa*. *mSystems*, 10(8), e00262-25. <https://doi.org/10.1128/msystems.00262-25>

Sun, F., Liang, H., Kong, X., Xie, S., Cho, H., Deng, X., Ji, Q., Zhang, H., Alvarez, S., Hicks, L. M., Bae, T., Luo, C., Jiang, H., & He, C. (2012). Quorum-sensing *agr* mediates bacterial oxidation response via an intramolecular disulfide redox switch in the response regulator AgrA. *Proceedings of the National Academy of Sciences*, 109(23), 9095–9100. <https://doi.org/10.1073/pnas.1200603109>

Szklarczyk, D., Kirsch, R., Koutrouli, M., Nastou, K., Mehryary, F., Hachilif, R., Gable, A. L., Fang, T., Doncheva, N. T., Pyysalo, S., Bork, P., Jensen, L. J., & von Mering, C. (2023). The STRING database in 2023: Protein-protein association networks and functional enrichment analyses for any sequenced genome of interest. *Nucleic Acids Research*, 51(D1), D638–D646. <https://doi.org/10.1093/nar/gkac1000>