

ROB311 Quiz 2

Hanhee Lee

March 13, 2025

Contents

1	Bayesian Networks	2
1.1	Junction	2
1.1.1	Causal Chain	2
1.1.2	Common Cause	2
1.1.3	Common Effect	3
2	Dependence Separation	4
2.1	Independence	4
2.1.1	Blocked Undirected Path	4
2.1.2	Blocked Junction	4
2.2	Consequence of Dependence Separation	4
2.3	Canonical Problems	6
2.3.1	Undirected Path Blocked?	6
2.3.2	Independence	6

Probabilistic Inference Problems

1 Bayesian Networks

Definition: Vertices represent random variables and edges represent dependencies between variables.

1.1 Junction

Definition: A **junction** \mathcal{J} consists of three vertices, X_1 , X_2 , and X_3 , connected by two edges, e_1 and e_2 :



Figure 1

- X_1 and X_2 are not independent, X_2 and X_3 are not independent, but when is X_1 and X_3 independent?

1.1.1 Causal Chain

Definition: A causal chain is a junction \mathcal{J} s.t.



Figure 2

- X_1 and X_3 are not independent (unconditionally), but are independent given X_2 .

Notes:

- **Analogy:** Given X_2 , X_1 and X_3 are independent. Why? X_2 's door closes when you know X_2 , so X_1 and X_3 are independent.
- **Distinction b/w Causal and Dependence:** X_1 and X_2 are dependent. However, from a causal perspective, X_1 is influencing X_2 (i.e. $X_1 \rightarrow X_2$).

1.1.2 Common Cause

Definition: A common cause is a junction \mathcal{J} s.t.




Figure 3

- X_1 and X_3 are not independent (unconditionally), but are independent given X_2 .

Notes:

- **Analogy:** Given X_2 , X_1 and X_3 are independent. Why? Consider the following example:
 - Let X_2 represent whether a person smokes or not, X_1 represent whether they have yellow teeth, X_3 represent whether they have lung cancer.
- Without knowing X_2 , observing X_1 provides information about X_3 because yellow teeth are associated with smoking, which in turn increases the likelihood of lung cancer.
- If X_2 is known, then knowing whether a person has yellow teeth provides no additional information about whether they have lung cancer beyond what is already known from smoking status.

1.1.3 Common Effect

Definition: A common effect is a junction \mathcal{J} s.t.




Figure 4

- X_1 and X_3 are independent (unconditionally), but are not independent given X_2 or any of X_2 's descendants.

Notes:

- **Analogy:** Consider the following example:
 - Let X_2 represent whether the grass is wet, X_1 represent whether it rained, X_3 represent whether the sprinkler was on.
- Without knowing whether the grass is wet (X_2), the occurrence of rain (X_1) and the sprinkler being on (X_3) are independent events. The rain may occur regardless of the sprinkler, and vice versa.
- However, once we observe that the grass is wet (X_2), the two events become dependent:
 - If we learn that the sprinkler was not on, then the wet grass must have been caused by rain.
 - If we learn that it did not rain, then the wet grass must have been caused by the sprinkler.

2 Dependence Separation

2.1 Independence

Theorem: Any two variables, X_1 and X_2 , in a Bayesian network, $\mathcal{B} = (\mathcal{V}, \mathcal{E})$, are independent given $\mathcal{K} \subseteq \mathcal{V}$ if every undirected path is blocked.

2.1.1 Blocked Undirected Path

Definition: An undirected path,

$$p = \langle (X_1, e_1, X_2), \dots, (X_{|p|-1}, e_{|p|-1}, X_{|p|}) \rangle,$$




Figure 5

is **blocked** given $\mathcal{K} \subseteq \mathcal{V}$ if any of its junctions,

$$\mathcal{J}^{(n)} = \{(X_{n-1}, X_n, X_{n+1}), (e_{n-1}, e_n)\},$$

is blocked given \mathcal{K} .

2.1.2 Blocked Junction

Definition: $\mathcal{J} = (\{X_1, X_2, X_3\}, \{e_1, e_2\})$ is **blocked** given $\mathcal{K} \subseteq \mathcal{V}$ if X_1 and X_3 are independent given \mathcal{K} .

2.2 Consequence of Dependence Separation

Theorem: For any variable, $X \in \mathcal{V}$, it can be shown that X is independent of X 's non-descendants, $\mathcal{V} \setminus \text{des}(X)$, given X 's parents, $\text{pts}(X)$.

Notes:




Figure 6

- Given X 's parent, apply junction rules to determine that X is independent of its non-descendants.
- $\mathcal{J} = \{(Z_1, \text{pt}(X), X), (e_1, e)\}$ shows that Z_1 and X are independent given $\text{pt}(X)$ (causal chain).
- $\mathcal{J} = \{(Z_2, \text{pt}(X), X), (e_2, e)\}$ shows that Z_2 and X are independent given $\text{pt}(X)$ (common cause).
- Given $\text{ch}(X)$'s parent, apply junction rules to determine that $\text{ch}(X)$ is independent of its non-descendants.
- $\mathcal{J} = \{\text{pt}(X), X, \text{ch}(X)\}, (e, e')\}$ shows that $\text{pt}(X)$ and $\text{ch}(X)$ are independent given X (causal chain).

- Given Z_4 's parent, apply junction rules to determine that Z_4 is independent of its non-descendants.
- $\mathcal{J} = \{X, \text{ch}(X), Z_4, (e', e_4)\}$ shows that X and Z_4 are independent given $\text{ch}(X)$ (causal chain).
- CHECK THIS OVER AGAIN WITH THE PROFESSOR.

2.3 Canonical Problems

2.3.1 Undirected Path Blocked?

Process:

1. **Given:** Undirected path p and \mathcal{K}
2. Check if any of the junctions on the undirected path are blocked given \mathcal{K} .
 - i.e. Check if X_1 and X_3 of the junction are independent given \mathcal{K} .

2.3.2 Independence

Process:

1. Given a Bayesian network w/ 2 variables to find independence.
2. Find all undirected paths between the 2 variables in the Bayesian network.
3. Identify a set of variables, \mathcal{K} , that block at least one junction in all undirected paths.
 - Test a junction by seeing junction given relationships.
4. If all undirected paths are blocked, then the 2 variables are independent given \mathcal{K} .

Warning:

- Be careful of common effect, in which it is blocked by default.
- Be careful of decedents of a common effect (i.e. outward arrows from a common effect) as given it may not be blocked.
- Cyclic paths are not blocked by default.

Example:

1. **Given:** Bayesian network.




Figure 7

2. **Problem:** A and E are
 - independent if $\mathcal{K} =$
 - not necessarily independent for $\mathcal{K} =$
3. **Soln:**
 - (a) **Undirected Paths:**
 - $A \rightarrow G \rightarrow H \rightarrow B \rightarrow E$
 - $A \rightarrow G \rightarrow D \rightarrow B \rightarrow E$
 - $A \rightarrow C \rightarrow F \rightarrow B \rightarrow E$
 - $A \rightarrow C \rightarrow D \rightarrow B \rightarrow E$

Example: Independent:

\mathcal{K}
$\{G, C\}$
<ul style="list-style-type: none"> • $A \iff G \iff H \iff B \iff E$ is blocked given \mathcal{K} since $\mathcal{J} = \{(A, G, H), (e_1, e_2)\}$ is blocked given G since A, H independent given G (causal chain) • $A \iff G \iff D \iff B \iff E$ is blocked given \mathcal{K} since $\mathcal{J} = \{(A, G, D), (e_1, e_2)\}$ is blocked given G since A, D independent given G (causal chain) • $A \iff C \iff F \iff B \iff E$ is blocked given \mathcal{K} since $\mathcal{J} = \{(A, C, F), (e_1, e_2)\}$ is blocked given C since A, F independent given C (causal chain) • $A \iff C \iff D \iff B \iff E$ is blocked given \mathcal{K} since $\mathcal{J} = \{(A, C, D), (e_1, e_2)\}$ is blocked given C since A, D independent given C (causal chain)
$\{D, F\}$
<ul style="list-style-type: none"> • $A \iff G \iff H \iff B \iff E$ is blocked given \mathcal{K} since $\mathcal{J} = \{(G, H, B), (e_1, e_2)\}$ is blocked NOT given H since G, B independent NOT given H (common effect) • $A \iff G \iff D \iff B \iff E$ is blocked given \mathcal{K} since $\mathcal{J} = \{(G, D, B), (e_1, e_2)\}$ is blocked given D since G, B independent given D (causal chain) • $A \iff C \iff F \iff B \iff E$ is blocked given \mathcal{K} since $\mathcal{J} = \{(C, F, B), (e_1, e_2)\}$ is blocked given F since C, B independent given F (causal chain) • $A \iff C \iff D \iff B \iff E$ is blocked given \mathcal{K} since $\mathcal{J} = \{(C, D, B), (e_1, e_2)\}$ is blocked given D since C, B independent given D (causal chain)

Not Necessarily Independent:

\mathcal{K}
$\{H, D, F\}$
<ul style="list-style-type: none"> • $A \iff G \iff H \iff B \iff E$ is unblocked given \mathcal{K} since $\mathcal{J} = \{(G, H, B), (e_1, e_2)\}$ is unblocked given H since G, B not independent given H (common effect) • $A \iff G \iff D \iff B \iff E$ is blocked given \mathcal{K} since $\mathcal{J} = \{(G, D, B), (e_1, e_2)\}$ is blocked given D (causal chain) since G, B independent given D (causal chain) • $A \iff C \iff F \iff B \iff E$ is blocked given \mathcal{K} since $\mathcal{J} = \{(C, F, B), (e_1, e_2)\}$ is blocked given F since C, B independent given F (causal chain) • $A \iff C \iff D \iff B \iff E$ is blocked given \mathcal{K} since $\mathcal{J} = \{(C, D, B), (e_1, e_2)\}$ is blocked given D since C, B independent given D (causal chain)

Example: Determine all subsets of $\{B, C, D, F, G, H\}$ for which A and E are independent.




Figure 8

1. **Undirected Paths:**

- $A \rightarrow G \rightarrow H \rightarrow B \rightarrow E$
- $A \rightarrow G \rightarrow D \rightarrow B \rightarrow E$
- $A \rightarrow C \rightarrow F \rightarrow B \rightarrow E$
- $A \rightarrow C \rightarrow D \rightarrow B \rightarrow E$

\mathcal{K}

$\{B\}$ (Any subset that contains B will be independent)

- $AGHBE$ is b given \mathcal{K} since $\mathcal{J} = \{(H, B, E), (e_1, e_2)\}$ is b since H, E indep. given B (causal chain)
 - $AGDBE$ is b given \mathcal{K} since $\mathcal{J} = \{(D, B, E), (e_1, e_2)\}$ is b since D, E indep. given B (causal chain)
 - $ACFBE$ is b given \mathcal{K} since $\mathcal{J} = \{(F, B, E), (e_1, e_2)\}$ is b since F, E indep. given B (causal chain)
 - $ACDBE$ is b given \mathcal{K} since $\mathcal{J} = \{(D, B, E), (e_1, e_2)\}$ is b since D, E indep. given B (causal chain)
-

$\{C\}$ (Not independent)

- $AGDBE$ is ub given \mathcal{K} since $\forall \mathcal{J}$ on p , all are ub.
-

$\{D\}$ (Not independent)

- $ACFBE$ is ub given \mathcal{K} since $\forall \mathcal{J}$ on p , all are ub.
-

$\{F\}$ (Not independent)

- $AGDBE$ is ub given \mathcal{K} since $\forall \mathcal{J}$ on p , all are ub.
-

$\{G\}$ (Not independent)

- $ACFBE$ is ub given \mathcal{K} since $\forall \mathcal{J}$ on p , all are ub.
-

$\{H\}$ (Not independent)

- $ACFBE$ is ub given \mathcal{K} since $\forall \mathcal{J}$ on p , all are ub.
-

Example:

 \mathcal{K}

 $\{C, D\}$ (Any subset that contains C and D except H will be independent)

- $AGHBE$ is b given \mathcal{K} since $\mathcal{J} = \{(G, H, B), (e_1, e_2)\}$ is b since G, B indep. not given H (causal effect)
 - $AGDBE$ is b given \mathcal{K} since $\mathcal{J} = \{(G, D, B), (e_1, e_2)\}$ is b since G, B indep. given D (causal chain)
 - $ACFBE$ is b given \mathcal{K} since $\mathcal{J} = \{(A, C, F), (e_1, e_2)\}$ is b since A, F indep. given C (causal chain)
 - $ACDBE$ is b given \mathcal{K} since $\mathcal{J} = \{(A, C, D), (e_1, e_2)\}$ is b since A, D indep. given C (causal chain)
-

 \dots

Example: Any set with

$$B \vee [(G \vee (D \wedge (\neg H))) \wedge (C \vee F)]$$

Example:

1. **Given:**




Figure 9

2. **Problem 1:** Is it guaranteed that $A \perp C$?
3. **Solution 1:** True b/c all undirected paths are blocked.
 - (a) ADEFC is b since $\mathcal{J} = \{(D, E, F), (e_1, e_2)\}$ is b since D, F indep. NOT given E (common effect)
 - (b) ADGHEFC is b since $\mathcal{J} = \{(G, H, E), (e_1, e_2)\}$ is b since G, E indep. NOT given H (common effect)
 - (c) ADGHIEFC is b since $\mathcal{J} = \{(H, I, E), (e_1, e_2)\}$ is b since H, E indep. NOT given I (common effect)
4. **Problem 2:** Is it guaranteed that $B \perp C \mid I$?
5. **Solution 2:** False b/c BEFC is ub.
 - (a) BEFC is ub since $\mathcal{J} = \{(B, E, F), (e_1, e_2)\}$ is ub since B, F NOT indep. given E 's descendent, I (common effect)
6. **Problem 3:** Is it guaranteed that $D \perp I \mid \{E, G\}$?
7. **Solution 3:** True b/c all undirected paths are blocked.
 - (a) DEI is b since $\mathcal{J} = \{(D, E, I), (e_1, e_2)\}$ is b since D, I indep. given E (causal chain)
 - (b) DEHI is b since $\mathcal{J} = \{(D, E, H), (e_1, e_2)\}$ is b since D, H indep. given E (causal chain)
 - (c) DGHI is b since $\mathcal{J} = \{(D, G, H), (e_1, e_2)\}$ is b since D, H indep. given G (causal chain)
 - (d) DGHEI is b since $\mathcal{J} = \{(D, G, H), (e_1, e_2)\}$ is b since D, H indep. given G (causal chain)
8. **Problem 4:** Is it guaranteed that $C \perp H \mid G$?
9. **Solution 4:** False b/c CFEH is ub.
 - (a) CFEH is ub since $\mathcal{J} = \{(C, F, E), (e_1, e_2)\}$ is ub since C, E NOT indep. given G (causal chain)
10. **Problem 5:** Suppose each variable is binary. What is the size of the domain of the joint distribution b/w the variables?
11. **Solution 5:**
 - (a) Since 9 variables, each with 2 values, the size of the domain of the joint distribution is $2^9 = 512$.
12. **Problem 6:** Suppose each variable is binary. What is the min # of values that actually need to be stored to represent the joint distribution entirely based on the Bayesian network? Use the fact that probability distributions are normalized.
13. **Solution 6:** $1+1+1+2+8+2+2+4+4 = 25$ values need to be stored.
 - (a) $P(A), P(B), P(C)$ has 1 value each
 - Since $P(\#)$ can represent 2 values, i.e. $P(0) = 1 - P(1)$, so only need to store 1 value.
 - (b) $P(D \mid A), P(F \mid C), P(G \mid D)$ has 2 values each
 - Same idea, can take the complement of the other value for 4 values.
 - (c) $P(H \mid G, E), P(I \mid E, H)$ has 4 values each
 - Same idea, can take the complement of the other value for 8 values.
 - (d) $P(E \mid D, B, F)$ has 8 values
 - Same idea, can take the complement of the other value for 16 values.