

ECE368 Cheatsheet

Hanhee Lee

January 15, 2025

Contents

1	L1: Probability Review	2
1.1	Sample Space	2
1.2	Probability Definitions	2
1.3	Axioms of Probability	2
1.4	Conditional Probability	3
1.4.1	Consequences of Conditional Probability	3
1.4.2	Independence	3
1.4.3	Importance of Labelling	3
2	L2: Probability Review	4
2.1	Total Probability	4
2.2	Bayes' Rule	4
2.2.1	Posteriori Probability, Priori Probability (Prior), Likelihood	4
2.2.2	Interpretation of Bayes' Rule	4
2.3	Random Variables	5
2.4	Distribution of RV	5
2.4.1	Cumulative Distribution Function (CDF) of RV	5
2.4.2	Discrete RV Probability Mass Function (PMF)	5
2.4.3	Continuous RV Probability Density Function (PDF)	5
2.4.4	Conditional PMF/PDF	6
2.5	Expected Values	6
3	L3: Probability Review	8
3.1	2 RVs	8
3.2	Joint PMF/PDF	8
3.3	Expectations	8
3.3.1	Correlation	8
3.3.2	Covariance	9
3.3.3	Correlation Coefficient	9
3.4	Marginal PMF/PDF	9
3.5	Conditional PMF/PDF	9
3.6	Bayes' Rule	10
3.7	Independent vs. Uncorrelated vs. Orthogonal	10
3.8	Conditional Expectation	10
4	L4: Estimation of Sample Mean	13
4.1	Parameter Estimation:	13
4.2	Estimator:	13
4.2.1	Estimation Error:	13
4.2.2	Unbiased	13
4.2.3	Consistent	13
4.3	Sample Mean & Law of Large Numbers	13
4.3.1	Digression for Sum of RVs (not necessarily independent or identically distributed)	14
4.3.2	Unbiased (i.i.d.)	14
4.3.3	Consistent (i.i.d.)	15

4.3.4	Weak Law of Large Numbers	15
4.3.5	Confidence Interval: Finding n	15
5	L5: Sample Mean and Maximum Likelihood Estimation	17
5.1	Maximum Likelihood Estimation	17
5.1.1	Log-Likelihood	17
6	L6: Maximum Likelihood and Laplace	18

Summary: On second thoughts, the lecture notes he posts are good, so I think I'll just do the cheatsheet.

W1 (LG-IPPR 1.1, 1.2; Murphy 2.1 – 2.3)

1 L1: Probability Review

Summary:

FAQ:

- How to study? Practice, practice.
- What textbooks? Use 2024 version of Murphy, Leon Garcia as main reference, Bishop, 4th textbook is intro.
- How is HW graded? Effort, and tutorials are used to explain soln.

1.1 Sample Space

Motivation: If you have 4 sheep and a flea, the probability that starting from sheep 1, the flea will jump to sheep 4 in 10 steps is 0.2.

- Ambiguous as there are 2 different interpretations for the sample space (i.e. space of probability is not clear):
 - Set of sheep
 - Set of number of steps

1.2 Probability Definitions

Definition:

- **Random Experiment:** An outcome (realization) for each run.
- **Sample Space Ω :** Set of all possible outcomes.
- **Events:** (measurable) subsets of Ω .
- **Probability of Event A :** $P[A] \equiv P[\text{'outcome is in } A\text{'}]$.

Example: Roll Fair Die

- $\Omega = \{1, 2, 3, 4, 5, 6\}$.
- $P[\text{'even number'}] = \frac{1}{2}$.

1.3 Axioms of Probability

Definition:

1. $P[A] \geq 0$ for all $A \in \Omega$.
2. $P[\Omega] = 1$.
3. If $A \cap B = \emptyset$, then $P[A \cup B] = P[A] + P[B]$ for all $A, B \in \Omega$.



Figure 1: 3rd Axiom

1.4 Conditional Probability

Definition:

$$P[A|B] = \frac{P[A \cap B]}{P[B]} \quad (1)$$

- $|$: Given event (data/obs.).




Figure 2: Conditional Probability

Notes:

- Changing sample space to B .
- Conditional probability satisfy the 3 axioms (i.e. are probabilities), can be viewed as probability measure on new sample space B .

1.4.1 Consequences of Conditional Probability

Definition:

$$P[A \cap B] = P[A|B]P[B] = P[B|A]P[A] \quad (2)$$

1.4.2 Independence

Definition: A and B are independent iff

$$P[A \cap B] = P[A]P[B] \iff P[A|B] = P[A] \iff P[B|A] = P[B] \quad (3)$$

1.4.3 Importance of Labelling

Example: Toss 2 Fair Coins

1. **Given:** Given that one of the coins is heads, what is the probability that the other coin is tails?
2. **Wrong Solution:** $\frac{1}{2}$ since $\{HH, HT, TH, TT\}$, so $P[T|H] = \frac{1}{2}$, which assumes that the coins are distinguishable (i.e. coin #1 is heads)
3. **Correct Solution:** $\frac{2}{3}$ since $\{HH, HT, TH\}$ as we didn't specify which coin was heads, so $P[T|H] = \frac{2}{3}$, which assumes that the coins are indistinguishable.

2 L2: Probability Review

2.1 Total Probability

Definition: If H_1, \dots, H_n form a partition of Ω , then

$$P[A] = \sum_{i=1}^n P[A|H_i]P[H_i] \quad (4)$$




Figure 3: Total Probability

2.2 Bayes' Rule

Definition:

$$P[H_k|A] = \frac{P[H_k \cap A]}{P[A]} = \frac{P[A|H_k]P[H_k]}{\sum_{i=1}^n P[A|H_i]P[H_i]} \quad (5)$$

2.2.1 Posteriori Probability, Priori Probability (Prior), Likelihood

Definition:

- **Posteriori:** $P[H_k|A]$.
- **Priori:** $P[H_k]$.
- **Likelihood:** $P[A|H_k]$.

Example: Suppose a lie detector is 95% accurate, i.e. $P[\text{'out=truth'}|\text{'in=truth'}] = 0.95$ and $P[\text{'out=lie'}|\text{'in=lie'}] = 0.95$. It says that Mr. Ernst is lying. What is the probability Mr. Ernst is actually lying.

- **Observation:** $A = \text{'out=lie'}$.
- **Hypothesis:** $H_0 = \text{'in=lie'}$ and $H_1 = \text{'in=truth'}$.
- **Solution:**
$$P[H_0|A] = \frac{P[A|H_0]P[H_0]}{P[A|H_0]P[H_0] + P[A|H_1]P[H_1]} = \frac{0.95 \times P[H_0]}{0.95 \times P[H_0] + 0.05 \times (1 - P[H_0])}.$$
- $H_0 = 0.01$: i.e. 1% of the population are liars, then
$$P[H_0|A] = \frac{0.95 \times 0.01}{0.95 \times 0.01 + 0.05 \times 0.99} = 0.16.$$

Warning: Need to know priori probability.

2.2.2 Interpretation of Bayes' Rule

Notes: Taking one component of the total probability and normalizing it by the sum of all components.

2.3 Random Variables

Motivation: Coin Toss Mapping of each outcome to a real number

- $w \in \Omega$ is the outcome of a coin toss, and X is the RV, so $H \rightarrow 0$ and $T \rightarrow 1$.



Figure 4: Random Variables

- Mapping is deterministic function. RV is not random or variable.

Definition: Mapping from Ω to \mathbb{R} .

2.4 Distribution of RV

2.4.1 Cumulative Distribution Function (CDF) of RV

Definition:

$$F_X(x) \equiv P[X \leq x] \quad (6)$$

2.4.2 Discrete RV Probability Mass Function (PMF)

Definition:

$$P_X(x_j) \equiv P[X = x_j] \quad j = 1, 2, 3, \dots \quad (7)$$

Example: Binomial RV w/ (n, p)

$$P_X(k) = \binom{n}{k} p^k (1-p)^{n-k} \quad (8)$$




Figure 5: Binomial RV

2.4.3 Continuous RV Probability Density Function (PDF)

Definition:

$$f_X(x) \equiv \frac{d}{dx} F_X(x) \quad (9)$$

$$P[x < X < x + dx] = f_X(x)dx \quad (10)$$

Example: Gaussian RV w/ (μ, σ^2)

$$f_X(x) = \frac{1}{\sqrt{2\pi}\sigma} e^{-\frac{(x-\mu)^2}{2\sigma^2}} \quad (11)$$




Figure 6: Gaussian RV

- $P[X \in A] = \int_A f_X(x)dx.$

Notes: Discrete RV has pdf w/ δ functions.

2.4.4 Conditional PMF/PDF

Definition:

$$P_X(x|A) \quad (12)$$

$$f_X(x|A) \quad (13)$$

Example: Continuous

$$f(x|X > a) = \begin{cases} \frac{f_X(x)}{P[X > a]} & \text{if } x > a \\ 0 & \text{otherwise} \end{cases} \quad (14)$$

Example: Geometric RV Geometric RV X w/ success probability p

$$P_X(k) = (1-p)^{k-1}p \quad (15)$$

- **Memoryless Property:** $P_X[k|X > m] = \frac{p(1-p)^{k-1}}{(1-p)^m} = p(1-p)^{k-m-1}.$
 - So it only cares about the additional trials (i.e. same as resetting after m trials).

2.5 Expected Values

Definition:

$$E[X] = \int_{-\infty}^{\infty} x f_X(x) dx \stackrel{\text{If int. values}}{=} \sum_{k=-\infty}^{\infty} k f_X(k) \quad (16)$$

$$E[h(X)] = \int_{-\infty}^{\infty} h(x) f_X(x) dx \stackrel{\text{If int. values}}{=} \sum_{k=-\infty}^{\infty} h(k) f_X(k) \quad (17)$$

$$\text{Var}[X] = E[(X - E[X])^2] = E[X^2] - E[X]^2 \quad (18)$$

$$E[X|A] = \int_{-\infty}^{\infty} x f_X(x|A) dx \quad (19)$$

Example: Lottery Ticket (Geometric RV)

1. **Given:** Buying one lottery ticket per week
 - Each ticket has $10^{-7} = p$ chance of winning the jackpot.
 - X = '# of weeks to win jackpot'.
2. **Problem:** What is the expected number of weeks to win the jackpot?
3. **Solution:** $E[X] = \sum_{k=1}^{\infty} k(1-p)^{k-1}p = \dots = \frac{1}{p} = 10^7$ weeks.
4. **Extension (Memoryless Property):** If I have already played for 999999 weeks, what is the expected number of weeks to win the jackpot? $E[X - 999999 | X > 999999] = E[X] = 10^7$ weeks.

3 L3: Probability Review

3.1 2 RVs

Notes: RVs are neither random nor a variable.

$$\underline{Z} = (X, Y)$$




Figure 7: Mapping of RVs

3.2 Joint PMF/PDF

Definition:

$$P_{X,Y}(x, y) = P[X = x, Y = y] \quad (20)$$

$$f_{X,Y}(x, y) = \frac{\partial^2}{\partial x \partial y} F_{X,Y}(x, y) \quad (21)$$

$$P[(X, Y) \in A] = \int \int_{(x,y) \in A} f_{X,Y}(x, y) dx dy \quad (22)$$

Example: Jointly Gaussian RVs X and Y with $(\mu_1, \mu_2, \sigma_1^2, \sigma_2^2, \rho)$

$$f_{X,Y}(x, y) = \frac{1}{2\pi\sigma_1\sigma_2\sqrt{1-\rho^2}} \exp \left\{ -\frac{1}{2(1-\rho^2)} \left[\left(\frac{x-\mu_1}{\sigma_1} \right)^2 - 2\rho \left(\frac{x-\mu_1}{\sigma_1} \right) \left(\frac{y-\mu_2}{\sigma_2} \right) + \left(\frac{y-\mu_2}{\sigma_2} \right)^2 \right] \right\}$$

3.3 Expectations

Definition:

$$E[g(X, Y)] = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} g(x, y) f_{X,Y}(x, y) dx dy$$

Notes:

- $g(X, Y)$ is also an RV, but inside the integral or sum, you use x and y as dummy variables to vary through the values of the RVs.

3.3.1 Correlation

Definition:

$$E[XY] \quad (23)$$

3.3.2 Covariance

Definition:

$$\text{Cov}[X, Y] = E[(X - \mu_X)(Y - \mu_Y)] = E[XY] - \mu_X \mu_Y = E[XY] - E[X]E[Y] \quad (24)$$

Notes:

- Mean shifted to 0.

3.3.3 Correlation Coefficient

Definition:

$$\rho_{X,Y} = E \left[\left(\frac{X - \mu_X}{\sigma_X} \right) \left(\frac{Y - \mu_Y}{\sigma_Y} \right) \right] = \frac{\text{Cov}[X, Y]}{\sigma_X \sigma_Y} \quad (25)$$

- $|\rho_{X,Y}| \leq 1$

Notes:

- Mean shifted to 0 and normalized by the standard deviation.

3.4 Marginal PMF/PDF

Definition:

$$P_X(x) = \sum_{j=1}^{\infty} P_{X,Y}(x, y_j), \quad P_Y(y) = \sum_{j=1}^{\infty} P_{X,Y}(x_j, y) \quad (26)$$

$$f_X(x) = \int_{-\infty}^{\infty} f_{X,Y}(x, y) dy, \quad f_Y(y) = \int_{-\infty}^{\infty} f_{X,Y}(x, y) dx \quad (27)$$

Notes:

- Total probability theorem is being used here.

Example: Jointly Gaussian X and Y :

$$\begin{aligned} f_X(x) &= \int_{-\infty}^{\infty} f_{X,Y}(x, y) dy \\ &= \dots \quad (\text{completing the square}) \\ &= \frac{1}{\sqrt{2\pi}\sigma_1} e^{-\frac{(x-\mu_1)^2}{2\sigma_1^2}}, \quad \text{marginally Gaussian} \end{aligned}$$

- Gaussian RVs has a property that the PDF of a single variable is equal to the marginal Gaussian of two variables.

3.5 Conditional PMF/PDF

Definition:

$$P_{X|Y}(x|y) \triangleq P[X = x|Y = y] = \frac{P_{X,Y}(x, y)}{P_Y(y)} \quad (28)$$

$$f_{X|Y}(x|y) \triangleq \frac{f_{X,Y}(x, y)}{f_Y(y)} \quad (29)$$

3.6 Bayes' Rule

Definition:

$$P_{Y|X}(x|y) = \frac{P_{X,Y}(x,y)}{P_X(x)} = \frac{P_{X|Y}(x|y)P_Y(y)}{\sum_{j=1}^{\infty} P_{X,Y}(x,y_j)P_Y(y_j)} \quad (30)$$

$$f_{Y|X}(y|x) = \frac{f_{X,Y}(x,y)}{f_X(x)} = \frac{f_{X|Y}(x|y)f_Y(y)}{\int_{-\infty}^{\infty} f_{X|Y}(x|y')f_Y(y') dy'} \quad (31)$$

3.7 Independent vs. Uncorrelated vs. Orthogonal

Definition:

1. Independent:

$$f_{X|Y}(x|y) = f_X(x) \quad \forall y \Leftrightarrow f_{X,Y}(x,y) = f_X(x)f_Y(y) \quad (32)$$

2. Uncorrelated:

$$\text{Cov}[X,Y] = 0 \quad \Leftrightarrow \quad \rho_{X,Y} = 0 \quad (33)$$

3. Orthogonal:

$$E[XY] = 0 \quad (34)$$

Theorem: If independent, then uncorrelated.

Derivation:

$$\begin{aligned} \text{Independent} &\implies E[XY] = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} xy f_{X,Y}(x,y) dx dy \\ &= \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} xy f_X(x) f_Y(y) dx dy \\ &= \left(\int_{-\infty}^{\infty} x f_X(x) dx \right) \left(\int_{-\infty}^{\infty} y f_Y(y) dy \right) \\ &\implies E[XY] = E[X]E[Y] \\ &\implies \text{Cov}[X,Y] = 0, \quad \text{uncorrelated} \\ &\neq \text{in general.} \end{aligned}$$

Example: Jointly Gaussian RVs X and Y : If uncorrelated, i.e. $\rho_{X,Y} = 0$, then X and Y are independent.

$$\begin{aligned} f_{X,Y}(x,y) &= \frac{1}{2\pi\sigma_1\sigma_2} \exp \left\{ -\frac{1}{2} \left[\left(\frac{x-\mu_1}{\sigma_1} \right)^2 + \left(\frac{y-\mu_2}{\sigma_2} \right)^2 \right] \right\} \\ &= \frac{1}{\sqrt{2\pi}\sigma_1} e^{-\frac{(x-\mu_1)^2}{2\sigma_1^2}} \cdot \frac{1}{\sqrt{2\pi}\sigma_2} e^{-\frac{(y-\mu_2)^2}{2\sigma_2^2}} \\ &= f_X(x)f_Y(y) \quad \text{independent} \end{aligned}$$

3.8 Conditional Expectation

Definition:

$$E[Y] = E[E[Y|X]] \quad (35)$$

$$E[h(Y)] = E[E[h(Y)|X]] \quad (36)$$

Notes:

- $E[E[Y|X]]$ is w.r.t. X .
- $E[Y|X]$ is w.r.t. Y .

Derivation:



$$\begin{aligned}
 E[Y] &= \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} y f_{X,Y}(x,y) dx dy \\
 &= \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} y f_{Y|X}(y|x) f_X(x) dx dy \\
 &= \int_{-\infty}^{\infty} \left(\int_{-\infty}^{\infty} y f_{Y|X}(y|x) dy \right) f_X(x) dx \\
 &= \int_{-\infty}^{\infty} E[Y|X=x] f_X(x) dx \quad (\text{using the total probability theorem}) \\
 &= \int_{-\infty}^{\infty} g(x) f_X(x) dx \\
 &= E[g(X)] \\
 &= E[E[Y|X]].
 \end{aligned}$$

Example:

1. **Given:** An unknown voltage. $X \sim \text{Uniform}(0, 1)$. Measurement from a (bad) voltmeter: $Y \sim \text{Uniform}(0, X)$.

$$\begin{aligned}
 f_X(x) &= \begin{cases} 1, & 0 < x < 1 \\ 0, & \text{otherwise} \end{cases} \\
 f_{Y|X}(y|x) &= \begin{cases} \frac{1}{x}, & 0 < y < x \\ 0, & \text{otherwise} \end{cases}
 \end{aligned}$$

- **Note:** Area under PDF is 1.

Figure 8: Uniform Distribution of X Figure 9: Uniform Distribution of Y

2. Expected Value (Average Reading of Bad Voltmeter):

$$\begin{aligned}
 E[Y] &= E[E[Y|X]] \\
 &= E\left[\frac{X}{2}\right] \quad \text{Since in the middle of 0 and x} \\
 &= \frac{1}{2} \cdot E[X] \\
 &= \frac{1}{2} \cdot \frac{1}{2} = \frac{1}{4} \quad \text{Since } E[X] \text{ (i.e. mean) is 0.5}
 \end{aligned}$$

3. The Long Way:

$$\begin{aligned}
 f_Y(y) &= \int_{-\infty}^{\infty} f_{Y|X}(y|x) f_X(x) dx \\
 &= \int_y^1 f_{Y|X}(y|x) f_X(x) dx \\
 &= \int_y^1 \frac{1}{x} \cdot 1 dx \\
 &= -\ln y. \\
 E[Y] &= \int_0^1 y \cdot (-\ln y) dy = \dots = \frac{1}{4}
 \end{aligned}$$

4. **Question:** Suppose $Y = \frac{1}{8}$. What is "best" given X ? This will be the question for the rest of the course.

4 L4: Estimation of Sample Mean

4.1 Parameter Estimation:

Motivation: The readout of a sensor is $X = \theta + N$ volts




Figure 10:

- There is some noise N in the sensor, so we want to estimate the true value of θ (unknown parameter to be estimated)
 - Mean and/or variance of X .

4.2 Estimator:

Definition: Perform n independent and identically distributed (i.i.d.) measurements/observations of X_i : X_1, X_2, \dots, X_n .

$$\hat{\Theta} = g(X_1, X_2, \dots, X_n) = g(\mathbf{X}) \quad (37)$$

4.2.1 Estimation Error:

Definition:

$$\hat{\Theta}(\underline{X}) - \theta \quad (38)$$

4.2.2 Unbiased

Definition: The estimator $\hat{\Theta}$ is unbiased if

$$\mathbb{E}[\hat{\Theta}(\underline{X})] = \theta \quad (39)$$

- **Asymptotically Unbiased:** $\lim_{n \rightarrow \infty} \mathbb{E}[\hat{\Theta}(\underline{X})] = \theta$ (big data)

4.2.3 Consistent

Definition: The estimator $\hat{\Theta}$ is consistent if $\hat{\Theta} \rightarrow \theta$ as $n \rightarrow \infty$, in probability, i.e., $\forall \epsilon > 0$,

$$\lim_{n \rightarrow \infty} P(|\hat{\Theta}(\underline{X}) - \theta| > \epsilon) \rightarrow 0 \quad (40)$$

as $n \rightarrow \infty$.

4.3 Sample Mean & Law of Large Numbers

Definition: Given a sequence of i.i.d. random variables (RVs), X_1, X_2, \dots, X_n , w/ unknown mean μ , estimate μ . Let $S_n = X_1 + X_2 + \dots + X_n$. The *sample mean* is

$$M_n = \frac{1}{n} S_n$$

- How good is M_n as an estimator of μ ?
 - Use unbiased and consistent to evaluate M_n .

Example: Previous voltage measurement, e.g.,

$$X_i = \mu + N_i$$

where μ is the true value and N_i is the noise.

If we assume N_i are i.i.d. with zero mean,

$$E[X_i] = E[\mu + N_i] = E[\mu] + E[N_i] = \mu + 0 = \mu, \quad \forall i$$

4.3.1 Digression for Sum of RVs (not necessarily independent or identically distributed)

Derivation:

$$\begin{aligned} E[S_n] &= E[X_1 + \cdots + X_n] \\ &= E[X_1] + \cdots + E[X_n] \end{aligned}$$

Derivation:

$$\begin{aligned} \text{Var}[S_n] &= E[(S_n - E[S_n])^2] \\ &= E\left[\left(\sum_{i=1}^n X_i - E[X_i]\right)^2\right] \\ &= E\left[\sum_{i=1}^n \sum_{j=1}^n (X_i - E[X_i])(X_j - E[X_j])\right] \\ &= \sum_{i=1}^n \sum_{j=1}^n E[(X_i - E[X_i])(X_j - E[X_j])] \\ &= \sum_{i=1}^n \sum_{j=1}^n \text{Cov}[X_i, X_j] \\ &= \sum_{i=1}^n \text{Var}[X_i] + \sum_{i \neq j} \text{Cov}[X_i, X_j] \end{aligned}$$

4.3.2 Unbiased (i.i.d.)

Derivation:

$$\begin{aligned} E[M_n] &= E\left[\frac{1}{n} S_n\right] \\ &= \frac{1}{n} (E[X_1] + \cdots + E[X_n]) \\ &= \frac{1}{n} (n\mu) \quad \text{since } X_i \text{ are i.i.d.} \\ &= \mu \Rightarrow \text{Unbiased!} \end{aligned}$$

4.3.3 Consistent (i.i.d.)

Derivation:

$$\begin{aligned}
 \text{Var}[M_n] &= \text{Var}\left[\frac{1}{n}S_n\right] \\
 &= \frac{1}{n^2}\text{Var}[S_n] \quad \text{taking out constant requires squaring} \\
 &= \frac{1}{n^2}\left(\sum_{i=1}^n \text{Var}[X_i] + \sum_{i \neq j} \text{Cov}[X_i, X_j]\right) \\
 &= \frac{1}{n^2}(n\sigma^2) \quad \sigma^2 \triangleq \text{Var}[X_i] \text{ and } X_i \text{ are i.i.d.} \\
 &= \frac{\sigma^2}{n} \rightarrow 0 \text{ as } n \rightarrow \infty.
 \end{aligned}$$

- This means that there is no variance in the sample mean as n approaches infinity, so it converges to the true mean.

Recall the Chebyshev Inequality:

$$P[|X - E[X]| \geq \epsilon] \leq \frac{\text{Var}[X]}{\epsilon^2}, \quad \forall \epsilon > 0.$$

Substitute in M_n :

$$\begin{aligned}
 P[|M_n - E[M_n]| \geq \epsilon] &\leq \frac{\text{Var}[M_n]}{\epsilon^2} \\
 P[|M_n - \mu| \geq \epsilon] &\leq \frac{\sigma^2}{n\epsilon^2} \\
 \Rightarrow P[|M_n - \mu| < \epsilon] &\geq 1 - \frac{\sigma^2}{n\epsilon^2} \rightarrow 1 \text{ as } n \rightarrow \infty \text{ then it is consistent}
 \end{aligned}$$

4.3.4 Weak Law of Large Numbers

Definition: Even if σ is infinite, then $\forall \epsilon > 0$,

$$\lim_{n \rightarrow \infty} P[|M_n - \mu| < \epsilon] = 1$$

4.3.5 Confidence Interval: Finding n

Example: Measure an unknown voltage θ for n times and obtain independent measurements:

$$X_i = \theta + N_i,$$

where N_i are i.i.d. random variables with mean 0 and variance 1.

- We want to determine how many measurements n are sufficient so that

$$P(|M_n - \theta| < 0.1) \geq 0.95,$$

where 0.1 is the desired precision and 0.95 is the confidence level.

- The sample mean is given by:

$$M_n = \frac{1}{n} \sum_{i=1}^n X_i = \theta + \frac{1}{n} \sum_{i=1}^n N_i.$$

- The variance of the sample mean is:

$$\sigma^2 = \text{Var}[M_n] = \text{Var}[N_i] = 1.$$

- Using Chebyshev's inequality:

$$1 - \frac{\sigma^2}{n\epsilon^2} \geq 0.95,$$

where $\epsilon = 0.1$ (precision).

- Solving for n :

$$1 - \frac{1}{n(0.1)^2} \geq 0.95,$$

$$\frac{1}{n(0.1)^2} \leq 0.05,$$

$$n \geq 2000.$$

Thus, at least 2000 measurements are needed to achieve the desired precision and confidence level.

5 L5: Sample Mean and Maximum Likelihood Estimation

5.1 Maximum Likelihood Estimation

Motivation: Choose parameter θ that is most likely to generate the observation x_1, x_2, \dots, x_n .




Figure 11:

Definition:

$$\hat{\Theta} = \arg \max_{\theta} P_{\underline{X}}(\underline{x}|\theta), \text{ discrete } X. \quad (41)$$

$$\hat{\Theta} = \arg \max_{\theta} f_{\underline{X}}(\underline{x}|\theta), \text{ continuous } X. \quad (42)$$

5.1.1 Log-Likelihood

Definition:

$$\hat{\theta} = \arg \max_{\theta} \sum_{i=1}^n \log P_X(x_i|\theta) \quad (43)$$

$$\hat{\theta} = \arg \max_{\theta} \sum_{i=1}^n \log f_X(x_i|\theta). \quad (44)$$

Example:

Notes:

6 L6: Maximum Likelihood and Laplace