Table 1: Five-DGP comparison. Each cell reports validity (%, ↑) / joint MMD (↓) / NN coverage (%, ↑), averaged over three seeds.

Method	Discrete	Linear	Cubic	Interaction	2D Bridge
CD	18.33 / 0.0896 / 44.45	83.22 / 0.0025 / 78.52	78.12 / 0.0062 / 74.35	84.98 / **0.0030** / 67.97	80.65 / 0.0072 / 73.08
Compositional CD	**94.37** / 0.0747 / 76.73	38.54 / 0.0139 / 31.25	31.74 / 0.0368 / 27.67	6.84 / 0.0717 / 6.05	20.93 / 0.1054 / 20.04
Additive mean†	0.00 / 0.1937 / 0.00	**100.00** / 0.0088 / 0.55	0.00 / 0.1711 / 0.00	**100.00** / 0.0169 / 0.65	0.00 / 0.3416 / 0.00
Additive + residual†	2.80 / 0.2020 / 10.60	7.81 / 0.1407 / 6.17	0.00 / 0.2037 / 0.00	3.11 / 0.1431 / 1.76	0.18 / 0.3266 / 0.10
Fixed composition + residual†	3.45 / 0.2344 / 12.47	0.03 / 0.2327 / 0.00	0.02 / 0.2014 / 0.00	0.05 / 0.3248 / 0.00	0.00 / 0.3798 / 0.00
Rep4Ex-inspired + residual†	1.70 / 0.2076 / 5.52	3.68 / 0.2490 / 3.06	0.23 / 0.2732 / 0.02	1.04 / 0.2488 / 0.75	0.03 / 0.3648 / 0.03
CG: sparse factorized ($s = 1$)	83.57 / **0.0489** / 89.53	5.40 / 0.0776 / 1.84	8.92 / 0.1280 / 4.07	2.03 / 0.1874 / 0.76	4.07 / 0.2045 / 3.52
CG: dense factorized ($s = 1$)	88.45 / 0.0661 / 85.35	7.67 / 0.0692 / 2.86	3.45 / 0.1266 / 0.65	1.32 / 0.1780 / 0.73	1.69 / 0.2084 / 1.79
CG: dense joint ($s = 1$)	1.40 / 0.4381 / 3.93	15.35 / 0.0359 / 6.74	13.22 / 0.0745 / 5.49	13.13 / 0.0339 / 4.26	6.18 / 0.1360 / 4.82
SEDGE: no weighting	7.33 / 0.2352 / 22.30	8.98 / 0.4006 / 20.82	8.97 / 0.3650 / 17.61	8.53 / 0.4809 / 14.01	2.46 / 0.6294 / 4.10
SEDGE-LGD ($s = 0.5$)	45.28 / 0.0703 / 74.82	44.11 / 0.1343 / 64.81	31.92 / 0.2048 / 49.61	49.25 / 0.1531 / 53.34	23.67 / 0.3420 / 35.40
SEDGE-LGD ($s = 1$)	77.55 / 0.0653 / **89.88**	78.24 / 0.0104 / 79.17	60.17 / 0.0711 / 64.49	78.96 / 0.0263 / 62.94	66.37 / 0.1153 / 66.60
SEDGE-LGD ($s = 2$)	69.52 / 0.0815 / 72.63	77.07 / 0.0302 / 62.42	71.89 / 0.0302 / 50.91	70.36 / 0.0457 / 42.33	76.25 / 0.0516 / 56.77
SEDGE-LWS	88.40 / 0.0624 / 88.92	91.98 / **0.0018** / **87.37**	**91.26** / **0.0024** / **84.29**	92.97 / 0.0039 / **79.12**	**87.39** / **0.0016** / **78.86**

Bold identifies the best displayed method within each DGP, separately for each metric; ties are bolded. No statistical-significance claim is implied. Validity requires both specifications and root support. Joint MMD compares (X, Z) distributions; NN coverage measures the fraction of reference points covered at the original fixed-radius criterion. †Paper-inspired adaptations, not faithful reproductions. The additive mean is a deterministic control, not a posterior sampler. All three previously completed LGD scales are shown; CG rows fix $s = 1$. The expanded LGD sweep is not included. The discrete reference has 5% conditional exclusion and is an explicitly relaxed Assumption-3 control; The four continuous settings pass the working 1% criterion. Results are an exploratory selection; all-method, per-seed data are retained.