Table 1: Five-DGP comparison. Each cell reports validity (%, ↑) / joint MMD (↓) / NN coverage (%, ↑), averaged over three seeds.

Method	Weighted Discrete	Linear	Cubic	Smoothstep	2D Bridge
CD	13.55 / 0.1467 / 40.00	83.22 / 0.0025 / 78.52	78.12 / 0.0062 / 74.35	85.55 / 0.0034 / 70.49	80.65 / 0.0072 / 73.08
Compositional CD	75.50 / 0.1065 / 84.78	38.54 / 0.0139 / 31.25	31.74 / 0.0368 / 27.67	21.21 / 0.0497 / 16.06	20.93 / 0.1054 / 20.04
Additive mean†	0.00 / 0.1174 / 0.00	**100.00** / 0.0088 / 0.55	0.00 / 0.1711 / 0.00	0.00 / 0.0716 / 0.00	0.00 / 0.3416 / 0.00
Additive + residual†	0.18 / 0.2322 / 0.65	7.81 / 0.1407 / 6.17	0.00 / 0.2037 / 0.00	1.35 / 0.1523 / 0.86	0.18 / 0.3266 / 0.10
Fixed composition + residual†	0.13 / 0.2353 / 0.50	0.03 / 0.2327 / 0.00	0.02 / 0.2014 / 0.00	0.07 / 0.2548 / 0.00	0.00 / 0.3798 / 0.00
Rep4Ex-inspired + residual†	0.85 / 0.2714 / 2.87	3.68 / 0.2490 / 3.06	0.23 / 0.2732 / 0.02	0.39 / 0.2621 / 0.02	0.03 / 0.3648 / 0.03
CG: sparse factorized ($s = 1$)	46.93 / 0.1027 / 74.85	5.40 / 0.0776 / 1.84	8.92 / 0.1280 / 4.07	3.43 / 0.1096 / 0.75	4.07 / 0.2045 / 3.52
CG: dense factorized ($s = 1$)	55.43 / 0.1073 / 75.58	7.67 / 0.0692 / 2.86	3.45 / 0.1266 / 0.65	2.86 / 0.1073 / 0.80	1.69 / 0.2084 / 1.79
CG: dense joint ($s = 1$)	1.47 / 0.5295 / 4.27	15.35 / 0.0359 / 6.74	13.22 / 0.0745 / 5.49	14.75 / 0.0526 / 4.57	6.18 / 0.1360 / 4.82
CG: sparse factorized ($s = 2$)	76.13 / 0.0530 / 61.22	24.82 / 0.0161 / 7.42	35.74 / 0.0183 / 13.85	19.48 / 0.0383 / 4.87	36.04 / 0.0758 / 30.99
CG: dense factorized ($s = 2$)	**82.85** / 0.0633 / 56.77	29.61 / 0.0116 / 9.83	26.16 / 0.0194 / 2.67	17.01 / 0.0423 / 1.14	5.70 / 0.0752 / 4.64
CG: dense joint ($s = 2$)	0.45 / 0.5508 / 1.42	18.20 / 0.0111 / 9.18	34.49 / 0.0143 / 13.77	25.72 / 0.0241 / 6.35	14.19 / 0.0394 / 9.52
SEDGE: no weighting	12.00 / 0.2198 / 29.95	8.98 / 0.4006 / 20.82	8.97 / 0.3650 / 17.61	8.92 / 0.4248 / 17.12	2.46 / 0.6294 / 4.10
SEDGE-LGD ($s = 0.5$)	38.07 / 0.1525 / 70.00	44.11 / 0.1343 / 64.81	31.92 / 0.2048 / 49.61	47.90 / 0.1267 / 60.01	23.67 / 0.3420 / 35.40
SEDGE-LGD ($s = 1$)	56.80 / 0.0996 / 81.08	78.24 / 0.0104 / 79.17	60.17 / 0.0711 / 64.49	79.57 / 0.0147 / 70.05	66.37 / 0.1153 / 66.60
SEDGE-LGD ($s = 2$)	39.98 / 0.0272 / 57.12	77.07 / 0.0302 / 62.42	71.89 / 0.0302 / 50.91	79.46 / 0.0328 / 55.27	76.25 / 0.0516 / 56.77
SEDGE-LWS	77.55 / **0.0225** / **89.97**	91.98 / **0.0018** / **87.37**	**91.26** / **0.0024** / **84.29**	**91.55** / **0.0030** / **82.23**	**87.39** / **0.0016** / **78.86**

Bold identifies the best displayed method within each DGP, separately for each metric; ties are bolded. No statistical-significance claim is implied. Validity requires both specifications and root support. Joint MMD compares (X, Z) distributions; NN coverage measures the fraction of reference points covered at the original fixed-radius criterion. †Paper-inspired adaptations, not faithful reproductions. The additive mean is a deterministic control, not a posterior sampler. All three previously completed LGD scales are shown; CG scales 1 and 2 are shown. The expanded LGD sweep is not included. Weighted Discrete has conditional exclusion 2.5% and 5%, and is an explicitly relaxed Assumption-3 control. The four continuous settings pass the working 1% criterion. Results are an exploratory selection; all-method, per-seed data are retained.