Table 1: Five-DGP comparison. Each cell reports validity (%, ↑) / joint MMD (↓) / NN coverage (%, ↑), averaged over three seeds.

Method	Weighted Discrete	Linear	Cubic	Smoothstep	2D Bridge
CD	13.55 / 0.1467 / 40.00	83.22 / **0.0025** / **78.52**	78.12 / 0.0062 / 74.35	**85.55** / 0.0034 / 70.49	80.65 / 0.0072 / 73.08
Compositional CD	**75.50** / 0.1065 / **84.78**	38.54 / **0.0139** / 31.25	31.74 / 0.0368 / 27.67	21.21 / 0.0497 / 16.06	20.93 / 0.1054 / 20.04
Additive mean†	0.00 / 0.1174 / 0.00	**100.00** / **0.0088** / **0.55**	0.00 / 0.1711 / 0.00	0.00 / 0.0716 / 0.00	0.00 / 0.3416 / 0.00
Additive + residual†	0.18 / 0.2322 / 0.65	**7.81** / **0.1407** / **6.17**	0.00 / 0.2037 / 0.00	1.35 / 0.1523 / 0.86	0.18 / 0.3266 / 0.10
Fixed composition + residual†	**0.13** / 0.2353 / **0.50**	0.03 / 0.2327 / 0.00	0.02 / **0.2014** / 0.00	0.07 / 0.2548 / 0.00	0.00 / 0.3798 / 0.00
Rep4Ex-inspired + residual†	0.85 / 0.2714 / 2.87	**3.68** / **0.2490** / **3.06**	0.23 / 0.2732 / 0.02	0.39 / 0.2621 / 0.02	0.03 / 0.3648 / 0.03
CG: sparse factorized ($s = 1$)	**46.93** / 0.1027 / **74.85**	5.40 / **0.0776** / 1.84	8.92 / 0.1280 / 4.07	3.43 / 0.1096 / 0.75	4.07 / 0.2045 / 3.52
CG: dense factorized ($s = 1$)	**55.43** / 0.1073 / **75.58**	7.67 / **0.0692** / 2.86	3.45 / 0.1266 / 0.65	2.86 / 0.1073 / 0.80	1.69 / 0.2084 / 1.79
CG: dense joint ($s = 1$)	1.47 / 0.5295 / 4.27	**15.35** / **0.0359** / **6.74**	13.22 / 0.0745 / 5.49	14.75 / 0.0526 / 4.57	6.18 / 0.1360 / 4.82
CG: sparse factorized ($s = 2$)	**76.13** / 0.0530 / **61.22**	24.82 / **0.0161** / 7.42	35.74 / 0.0183 / 13.85	19.48 / 0.0383 / 4.87	36.04 / 0.0758 / 30.99
CG: dense factorized ($s = 2$)	**82.85** / 0.0633 / **56.77**	29.61 / **0.0116** / 9.83	26.16 / 0.0194 / 2.67	17.01 / 0.0423 / 1.14	5.70 / 0.0752 / 4.64
CG: dense joint ($s = 2$)	0.45 / 0.5508 / 1.42	18.20 / **0.0111** / 9.18	**34.49** / 0.0143 / **13.77**	25.72 / 0.0241 / 6.35	14.19 / 0.0394 / 9.52
SEDGE: no weighting	**12.00** / **0.2198** / **29.95**	8.98 / 0.4006 / 20.82	8.97 / 0.3650 / 17.61	8.92 / 0.4248 / 17.12	2.46 / 0.6294 / 4.10
SEDGE-LGD ($s = 0.5$)	38.07 / 0.1525 / **70.00**	44.11 / 0.1343 / 64.81	31.92 / 0.2048 / 49.61	**47.90** / **0.1267** / 60.01	23.67 / 0.3420 / 35.40
SEDGE-LGD ($s = 1$)	56.80 / 0.0996 / **81.08**	78.24 / **0.0104** / 79.17	60.17 / 0.0711 / 64.49	**79.57** / 0.0147 / 70.05	66.37 / 0.1153 / 66.60
SEDGE-LGD ($s = 2$)	39.98 / **0.0272** / 57.12	77.07 / 0.0302 / **62.42**	71.89 / 0.0302 / 50.91	**79.46** / 0.0328 / 55.27	76.25 / 0.0516 / 56.77
SEDGE-LWS	77.55 / 0.0225 / **89.97**	**91.98** / 0.0018 / 87.37	91.26 / 0.0024 / 84.29	91.55 / 0.0030 / 82.23	87.39 / **0.0016** / 78.86

Bold identifies each method's best DGP separately for each metric (requested row-wise convention). This does not rank methods; MMD magnitudes across distinct DGPs need not be comparable. Validity requires both specifications and root support. Joint MMD compares (X, Z) distributions; NN coverage measures the fraction of reference points covered at the original fixed-radius criterion. †Paper-inspired adaptations, not faithful reproductions. The additive mean is a deterministic control, not a posterior sampler. All three previously completed LGD scales are shown; CG scales 1 and 2 are shown. The expanded LGD sweep is not included. Weighted Discrete has conditional exclusion 2.5% and 5%, and is an explicitly relaxed Assumption-3 control. The four continuous settings pass the working 1% criterion. Results are an exploratory selection; all-method, per-seed data are retained.