YONSEI UNIVERSITY

EverM Oral & Maxillofacial Surgery Clinic

Sang Hyun Song, DDS, Ph.D

Certificated board of Oral and Maxillofacial Surgery, Yonsei University Dental Hospital
Physician-Scientist(Ph.D), Department of Oral Pathology, Yonsei University College of Dentistry
Visiting Scholar, miRcore, Ann Arbor, Michigan

Severance

Work flow for Protrusion Surgery (ASO)

•**Standard:** Weekly post-op checks **until 1 month** → **wire removal at 1month** → refer to **Bangkok for orthodontics**

•**Current practice : EverM –Omakase Smile Dental Clinic only**

1. **EverM (Korea): Wafer fabrication(2-days before)** + Surgery + **1-week post-op check**
2. **Return to Bangkok:** after the **1-week check**
3. **Omakase (Bangkok): Weeks 2–4 checks + wire removal at 1 month + start/continue orthodontic treatment**

Work flow for Contouring Surgery

•**Current practice:**

1. **EverM (Korea): Health check(2-days before)** + Surgery + **1-week post-op check**
2. **Return to Bangkok:** after the **1-week check**

* *For patients who have undergone zygoma reduction surgery, it is best to check at the 6-month follow-up whether the zygomatic plates are broken.*

Etymology & Meaning

Ortho : Correct, Gnatho : Jaw, Odonto : Tooth

Orthognathic Surgery : Surgical correction of jaw (and sometimes facial) skeletal deformities to improve function (bite, chewing, speech, breathing) and facial balance

Orthodontic treatment : Treatment to correct the alignment of the teeth and the relationship of the jaws (occlusion), typically using braces, clear aligners, and other orthodontic appliances, to improve function and aesthetics.

Functional Orthognathic Surgery

: 2-Jaw Surgery or 1-Jaw Surgery with occlusion change (maloccluison -> normal occlusion)
: *Orthodontic treatment is necessary*

Method

1. Classic method (Total Period 2Y ~3Y)
   1) Pre-surgical orthodontic tx (6m ~1.5y)
   2) Orthognathic Surgery (Recovery 1m~1.5m)
   3) Post-surgical orthodontic tx (6m~1.5y)

2. Current trends (Total Period 1Y~1.5Y) -> Coordination with the orthodontic specialist is necessary
   1) Surgery (Recovery 1m~1.5m)
   2) Post-surgical orthodontic tx (6m~1.5y)

Functional Orthognathic Surgery

Surgical method for mandible



IVRO	SSRO
การตรึงขากรรไกร: ประมาณ 2 สัปดาห์ (แบบแข็ง)	การตรึงขากรรไกร: ประมาณ 1 สัปดาห์ (แบบนุ่ม)
ความเสี่ยงต่อเส้นประสาท: ต่ำกว่า SSRO	ความเสี่ยงต่อเส้นประสาท: สูงกว่า IVRO
การตรึงกระดูก: ไม่มี	การตรึงกระดูก: มี
ระยะเวลาทำกายภาพบำบัด: 4–5 สัปดาห์ (ช่วยฟื้นฟูข้อต่อขากรรไกรได้)	ระยะเวลาทำกายภาพบำบัด: 2–3 สัปดาห์

IVRO was developed by professors at Yonsei University.

Work flow for Functional Orthognathic Surgery(SSRO)

•**Current practice:**

1. **EverM (Korea): Health check(2-days before)** + Surgery + **1-week post-op check (Physical Therapy start)**
2. **Return to Bangkok:** after the **1-week check**
3. **EverM (Korea): Weeks 3–4 checks + wafer & screw removal**
4. **Omakase (Bangkok): Start Orthodontics**

Work flow for Functional Orthognathic Surgery(IVRO)

•**Current practice:**

1. **EverM (Korea): Health check(2-days before)** + Surgery + **2-week post-op check (Physical Therapy start)**
2. **Return to Bangkok:** after the **2-week check**
3. **EverM (Korea): Weeks 5–6checks + wafer & screw removal**
4. **Omakase (Bangkok): Start Orthodontics**

Esthetic Orthognathic Surgery

: 2-Jaw Surgery or 1-Jaw Surgery without occlusion change
: *Orthodontic treatment is not necessary*

For Facial Feminization Surgery(FFS), Asymmetry Correction, Facial shortening, Facial Lengthening etc.

Orthognathic Surgery is necessary for Facial Feminization.

Work flow for Esthetic Orthognathic Surgery(SSRO)

•**Current practice:**

1. **EverM (Korea): Health check(2-days before)** + Surgery + **1-week post-op check (Physical Therapy start)**
2. **Return to Bangkok:** after the **1-week check**
3. **EverM (Korea): Weeks 3 (wafer removal) + Weeks 4 (screw removal)**

- Orthodontic treatment is not necessary for esthetic orthognathic surgery, patient is needed to ***set the original occlusion for 1 week after wafer removal.***

- After esthetic orthognathic surgery***,*** some patients may still need a **lip lift (philtrum reduction surgery)**, because even if the facial height is reduced, the philtrum/upper lip soft tissue does not always follow the underlying skeletal movement 100%.

Work flow for Esthetic Orthognathic Surgery(IVRO)

•**Current practice:**

1. **EverM (Korea): Health check(2-days before)** + Surgery + **2-week post-op check (Physical Therapy start)**
2. **Return to Bangkok:** after the **2-week check**
3. **EverM (Korea): Weeks 5 (wafer removal) + Weeks 6 (screw removal)**

- Orthodontic treatment is not necessary for esthetic orthognathic surgery, patient is needed to ***set the original occlusion for 1 week after wafer removal.***

- After esthetic orthognathic surgery***,*** some patients may still need a **lip lift (philtrum reduction surgery)**, because even if the facial height is reduced, the philtrum/upper lip soft tissue does not always follow the underlying skeletal movement 100%.

Postoperative Precautions Following Orthognathic Surgery(SSRO)

Timepoint	Key points	Interpreter / Guardian
Pre-op (2 days before surgery)	**Wafer fabrication** and **informed consent** must be completed.	**Interpreter required**
Post-op Day 1	**Keep the airway tube in place** (do **not** remove due to discomfort; breathing may worse n). High anxiety; possible emergencies such as **syncope** or **hyperventilation**.	**Guardian & Interpreter required**
Post-op Day 2	Generally stabilizing, but **swelling may still trigger urgent events**.	As needed
Post-op Day 3 (Discharge)	**Intermaxillary fixation (IMF : Jaw fixation)** and **mandatory instructions**. If IMF cannot be properly established, **discharge/return travel may be delayed**.	**Interpreter required**
Post-op Week 1	Start **physical therapy (jaw movement exercises)**. Poor understanding/compliance may lead to **prolonged recovery, complications, or revision surgery**.	**Interpreter required**
Post-op Week 3	The wafer was removed at 3 weeks postoperatively. If orthodontic treatment is initiated, the screws will be removed at the same time. As CT comparison is performed at this visit, an interpreter is required.	**Interpreter required**
Post-op Week 4	At 4 weeks postoperatively, screw removal is performed for patients who are not undergoing orthodontic treatment. As CT comparison is performed at this visit, an interpreter is required.	**Interpreter required**

Postoperative Precautions Following Orthognathic Surgery(IVRO)

Timepoint	Key points	Interpreter / Guardian
Pre-op (2 days before surgery)	**Wafer fabrication** and **informed consent** must be completed.	**Interpreter required**
Post-op Day 1	**Keep the airway tube in place** (do **not** remove due to discomfort; breathing may worse n). High anxiety; possible emergencies such as **syncope** or **hyperventilation**.	**Guardian & Interpreter required**
Post-op Day 2	Generally stabilizing, but **swelling may still trigger urgent events**.	As needed
Post-op Day 3 (Discharge)	**Intermaxillary fixation (IMF : Jaw fixation)** and **mandatory instructions**. If IMF cannot be properly established, **discharge/return travel may be delayed**.	**Interpreter required**
Post-op Week 2	Start **physical therapy (jaw movement exercises)**. Poor understanding/compliance may lead to **prolonged recovery, complications, or revision surgery**.	**Interpreter required**
Post-op Week 5	The wafer was removed at 5 weeks postoperatively. If orthodontic treatment is initiated, the screws will be removed at the same time. As CT comparison is performed at this visit, an interpreter is required.	**Interpreter required**
Post-op Week 6	At 6 weeks postoperatively, screw removal is performed for patients who are not undergoing orthodontic treatment. As CT comparison is performed at this visit, an interpreter is required.	**Interpreter required**

YONSEI UNIVERSITY
COLLEGE OF DENTISTRY