

# Truth Tables, Tautologies, and Logical Equivalences

Mathematicians normally use a **two-valued logic**: Every statement is either **True** or **False**. This is called the **Law of the Excluded Middle**.

A statement in sentential logic is built from simple statements using the logical connectives  $\sim$ ,  $\wedge$ ,  $\vee$ ,  $\rightarrow$ , and  $\leftrightarrow$ . The truth or falsity of a statement built with these connective depends on the truth or falsity of its components.

For example, the compound statement  $P \rightarrow (Q \vee \sim R)$  is built using the logical connectives  $\rightarrow$ ,  $\vee$ , and  $\sim$ . The truth or falsity of  $P \rightarrow (Q \vee \sim R)$  depends on the truth or falsity of P, Q, and R.

A **truth table** shows how the truth or falsity of a compound statement depends on the truth or falsity of the simple statements from which it's constructed. So we'll start by looking at truth tables for the five logical connectives.

Here's the table for negation:

P	$\sim P$
T	F
F	T

This table is easy to understand. If P is *true*, its negation  $\sim P$  is *false*. If P is *false*, then  $\sim P$  is *true*.

$P \wedge Q$  should be *true* when both P and Q are *true*, and *false* otherwise:

P	Q	$P \wedge Q$
T	T	T
T	F	F
F	T	F
F	F	F

$P \vee Q$  is *true* if either P is *true* or Q is *true* (or both --- remember that we're using "or" in the inclusive sense). It's only *false* if both P and Q are *false*.

P	Q	$P \vee Q$
T	T	T
T	F	T
F	T	T
F	F	F

Here's the table for logical implication:

$P$	$Q$	$P \rightarrow Q$
T	T	T
T	F	F
F	T	T
F	F	T

To understand why this table is the way it is, consider the following example:

"If you get an A, then I'll give you a dollar."

The statement will be *true* if I keep my promise and *false* if I don't.

Suppose it's *true* that you get an A and it's *true* that I give you a dollar. Since I kept my promise, the implication is *true*. This corresponds to the first line in the table.

Suppose it's *true* that you get an A but it's *false* that I give you a dollar. Since I *didn't* keep my promise, the implication is *false*. This corresponds to the second line in the table.

What if it's false that you get an A? Whether or not I give you a dollar, I haven't broken my promise. Thus, the implication can't be false, so (since this is a two-valued logic) it must be true. This explains the last two lines of the table.

$P \leftrightarrow Q$  means that P and Q are **equivalent**. So the double implication is *true* if P and Q are both *true* or if P and Q are both *false*; otherwise, the double implication is false.

$P$	$Q$	$P \leftrightarrow Q$
T	T	T
T	F	F
F	T	F
F	F	T

You should remember --- or be able to construct --- the truth tables for the logical connectives. You'll use these tables to construct tables for more complicated sentences. It's easier to demonstrate what to do than to describe it in words, so you'll see the procedure worked out in the examples.

**Remark.** (a) When you're constructing a truth table, you have to consider all possible assignments of True (T) and False (F) to the component statements. For example, suppose the component statements are P, Q, and R. Each of these statements can be either true or false, so there are  $2^3 = 8$  possibilities.

When you're listing the possibilities, you should assign truth values to the component statements in a systematic way to avoid duplication or omission. The easiest approach is to use **lexicographic ordering**. Thus, for a compound statement with three components P, Q, and R, I would list the possibilities this way: