

Solving for Scale

*Spatial dimension, three fundamental quantities, and a progression
from back-substitution to the null space*

Lee McCulloch-James

Department of Natural & Unnatural Sciences, Dwight School London

July 23, 2026

Abstract

A unit, looked at properly, is a set of coordinates. Once we agree to track only the base dimensions of mass, length, time and temperature, every physical quantity is fixed by the powers to which those dimensions are raised, so that a velocity sits at $(0, 1, -1, 0)$ and an energy at $(1, 2, -2, 0)$, each a vector in the same space; and building one quantity as a product of powers of others is, in this language, the taking of a linear combination of such vectors. To ask which combination of given constants carries a desired dimension is therefore to solve a system of linear equations, and the character of that system, whether it has one solution, too many, or too few, mirrors the physics with uncommon faithfulness. This essay follows that single thread through three fundamental quantities of space, the Planck length with its area and volume, the mean density of the universe, and the entropy of a black hole, using each to introduce one further rung of a ladder that climbs from the back-substitution a pupil meets first, through Gaussian elimination, to the point at which the mathematics declines to finish the problem and a piece of physics must complete it.

1 Dimension as a coordinate

The constants we shall need carry the dimensions

$$[c] = \text{L T}^{-1}, \quad [h] = \text{M L}^2 \text{T}^{-1}, \quad [G] = \text{M}^{-1} \text{L}^3 \text{T}^{-2}, \quad [k_B] = \text{M L}^2 \text{T}^{-2} \Theta^{-1},$$

together with the Hubble constant $[H_0] = \text{T}^{-1}$ and a plain area $[A] = \text{L}^2$. Reading each as a column of powers turns the search for a formula into a matrix equation. If we seek a product $Q = \prod_i X_i^{p_i}$ with a prescribed dimension, then collecting the exponent of each base dimension gives one linear equation per dimension, and the whole problem becomes

$$\mathbf{D} \mathbf{p} = \mathbf{t},$$

where the columns of $\mathbf{D}$ are the dimension vectors of the chosen quantities, $\mathbf{p}$ is the vector of unknown exponents, and $\mathbf{t}$ is the target dimension. The spatial part of this picture is especially clean, since the lengths, areas and volumes a student cares about lie along a single axis: asking for L^n rather than L replaces $\mathbf{t}$ by $n \mathbf{t}$, and because the map is linear the exponents simply scale by n . Whatever recipe builds a fundamental length builds its area and volume by doubling and tripling the same powers, so the three spatial dimensions are one solution seen at three magnifications.

What remains is to solve $\mathbf{D} \mathbf{p} = \mathbf{t}$, and a linear system can behave in exactly three ways. It can have a single solution, it can be overdetermined yet consistent, and it can be underdetermined, leaving a family of solutions to be narrowed by some condition from outside. The three quantities below were chosen because each lands squarely in one of these cases.

2 One solution: a fundamental length

The Planck length is the unique combination of c , h and G with the dimension of length, written $\ell_P = c^\alpha h^\beta G^\gamma$ up to a pure number. Matching the dimensions $\mathbf{M}, \mathbf{L}, \mathbf{T}$ gives the system

$$\begin{aligned}\mathbf{M} : \quad & \beta - \gamma = 0, \\ \mathbf{L} : \quad & \alpha + 2\beta + 3\gamma = 1, \\ \mathbf{T} : \quad & -\alpha - \beta - 2\gamma = 0.\end{aligned}$$

Back-substitution. The first method a pupil owns is to read a relation off one equation and feed it forward. The mass line gives $\beta = \gamma$ at once; the time line then reads $\alpha = -\beta - 2\gamma = -3\beta$; and the length line becomes $-3\beta + 2\beta + 3\beta = 2\beta = 1$, so that $\beta = \frac{1}{2}$, whence $\gamma = \frac{1}{2}$ and $\alpha = -\frac{3}{2}$. It is quick here because the substitutions happen to fall in a convenient order, and that is also its weakness: the method offers no guarantee that such an order exists, and with more constants the bookkeeping soon outgrows the page.

Gaussian elimination, and the pivot. The substitution just done succeeded because the equations could be untangled in a lucky order, and the purpose of elimination is to manufacture that order on demand. We first set the system out as an *augmented matrix*, a row for each equation and a column for each unknown, with the right-hand sides written after a vertical bar, and we work on whole rows at a time. To *pivot* on an entry is then to use the row it lies in to remove that unknown from every other row, subtracting from each the multiple of the pivot row that cancels the entry standing in the pivot's column; the entry put to this use is the pivot, and forming that multiple means dividing by it, so a pivot must be a number one may divide by. Sweeping down the columns in this way empties everything below the leading diagonal and leaves each row holding one unknown fewer than the row above it, which is the staircase that back-substitution needs.

We begin with the first column, meaning to pivot on the top-left entry and clear the column beneath it; but that entry is a zero,

$$\left[\begin{array}{ccc|c} \boxed{0} & 1 & -1 & 0 \\ 1 & 2 & 3 & 1 \\ -1 & -1 & -2 & 0 \end{array} \right],$$

and clearing an entry below it would ask us to divide by that zero, which is barred. The repair is to exchange the first two rows, lifting a usable entry into the corner,

$$\left[\begin{array}{ccc|c} \boxed{1} & 2 & 3 & 1 \\ 0 & 1 & -1 & 0 \\ -1 & -1 & -2 & 0 \end{array} \right].$$

The 1 in the corner is now the pivot; adding its row to the third cancels the -1 beneath it and clears the column,

$$\left[\begin{array}{ccc|c} 1 & 2 & 3 & 1 \\ 0 & \boxed{1} & -1 & 0 \\ 0 & 1 & 1 & 1 \end{array} \right],$$

the 1 it exposes in the second column is the next pivot, and subtracting its row from the third clears that column in turn,

$$\left[\begin{array}{ccc|c} 1 & 2 & 3 & 1 \\ 0 & 1 & -1 & 0 \\ 0 & 0 & \boxed{2} & 1 \end{array} \right],$$

leaving the 2 in the third column as the last pivot.

The system is now a staircase, and back-substitution climbs it from the bottom, dividing by one pivot at each rung to release one unknown. The last pivot gives $2\gamma = 1$, so $\gamma = \frac{1}{2}$; the middle pivot gives $\beta - \gamma = 0$, so $\beta = \frac{1}{2}$; the first gives $\alpha + 2\beta + 3\gamma = 1$, so $\alpha = -\frac{3}{2}$. The same three pivots that cleared the columns going down are the three numbers we divide by coming back up, and because every column carried one, every unknown is released in turn and the answer is a single point,

$$\ell_P = \sqrt{\frac{hG}{c^3}} \approx 1.6 \times 10^{-35} \text{ m}.$$

The spatial dimensions for free. Because the target enters only through the right-hand column, asking instead for an area or a volume multiplies that column by two or three and, by linearity, the exponents with it. The Planck area and volume therefore require no fresh calculation,

$$\ell_P^2 = \frac{hG}{c^3} \approx 2.6 \times 10^{-70} \text{ m}^2, \quad \ell_P^3 = \left(\frac{hG}{c^3}\right)^{3/2} \approx 4.2 \times 10^{-105} \text{ m}^3,$$

and the exponents $-35, -70, -105$ advance in step because each spatial dimension is the same length vector taken once, twice or three times over.

3 Too many equations, and consistent: the density of the cosmos

The mean density of the universe is a quantity of dimension ML^{-3} , the negative cube that any density wears, and on the largest scale it is set by the Hubble constant and the strength of gravity alone, $\rho = H_0^p G^q$. Now there are three base dimensions to satisfy and only two exponents to satisfy them with, so the system is taller than it is wide:

$$\left[\begin{array}{c|cc|c} & H_0 & G & [\rho] \\ \hline \text{M} & 0 & -1 & 1 \\ \text{L} & 0 & 3 & -3 \\ \text{T} & -1 & -2 & 0 \end{array} \right] \longrightarrow \left[\begin{array}{cc|c} 1 & 2 & 0 \\ 0 & -1 & 1 \\ 0 & 0 & 0 \end{array} \right].$$

The elimination places a pivot in each of the two columns and then meets a third row with no column left to pivot in; that row collapses to zeros, which is what a redundant equation looks like once the work is done, the third equation having added nothing the first two had not already fixed. Its collapse is the certificate that the chosen constants can indeed reach the target. The two pivots determine the two unknowns, $q = -1$ and $p = 2$, so that

$$\rho = \kappa \frac{H_0^2}{G}, \quad \rho_c = \frac{3H_0^2}{8\pi G} \approx 9 \times 10^{-27} \text{ kg m}^{-3},$$

amounting to some five or six hydrogen atoms in every cubic metre of space. Had that last row instead kept a nonzero constant on the right, it would read $0 = \text{something}$, an impossibility, and the elimination would have reported that no product of H_0 and G carries the density's dimension; a row left without a pivot is redundant when its constant vanishes and contradictory when it does not, and the method declares which with equal candour.

4 Too few equations: a black hole's entropy

The entropy of a black hole brings in temperature, and with it a fourth base dimension and a fifth constant. Seeking $S = k_B^a A^b c^d h^e G^f$ with the dimension $\text{ML}^2 \text{T}^{-2} \Theta^{-1}$ of entropy, the

system has four equations, one for each base dimension, in five unknowns. Set out as an augmented matrix, with the five constants heading the columns and the four base dimensions labelling the rows, it reads

$$\left[\begin{array}{c|ccccc|c} & k_B & A & c & h & G & [S] \\ \hline \text{M} & 1 & 0 & 0 & 1 & -1 & 1 \\ \text{L} & 2 & 2 & 1 & 2 & 3 & 2 \\ \text{T} & -2 & 0 & -1 & -1 & -2 & -2 \\ \text{\Theta} & -1 & 0 & 0 & 0 & 0 & -1 \end{array} \right],$$

where each row imposes the homogeneity of one base dimension and the entry past the bar is the power that dimension carries in entropy itself, the target the rest must reproduce.

The temperature row carries only k_B , so it is an equation in the single unknown a , the easiest pivot the problem offers, handing over $a = 1$ before anything is cleared; this is the kind of pivot, sitting in a nearly empty row, that isolates its unknown at a stroke. Reducing the remaining rows to echelon form gives

$$\left[\begin{array}{ccccc|c} 1 & 0 & 0 & 1 & -1 & 1 \\ 0 & 2 & 1 & 0 & 5 & 0 \\ 0 & 0 & -1 & 1 & -4 & 0 \\ 0 & 0 & 0 & 1 & -1 & 0 \end{array} \right],$$

and a pivot now falls in each of the first four columns, those of k_B , A , c and h , while the fifth column, the one belonging to G , receives none. An unknown whose column carries no pivot is free, so the exponent f of G may be chosen at will, and the rows above fix the rest in terms of it, read from the bottom up: the last row gives $e = f$, the third $d = -3f$, the second $b = -f$, and the first $a = 1$, this final value holding whatever f is taken to be. That single loose exponent is where the freedom lives, and it is tidiest to follow it through the area exponent, setting $k = b = -f$, in which the whole family of dimensionally valid answers reads

$$(a, b, d, e, f) = (1, k, 3k, -k, -k) = (1, 0, 0, 0, 0) + k(0, 1, 3, -1, -1).$$

The vector added by k is the heart of the matter. As a product it reads $A c^3 / h G$, and that group is dimensionless, being precisely the horizon area measured in Planck areas, A / ℓ_P^2 ; adding any multiple of it therefore changes no dimension at all. Every member of the family carries the dimension of entropy exactly, so the units cannot prefer one over another, and the freedom is an honest report that dimensional reasoning has said all it can, rather than a defect of the calculation.

What closes the gap is physics. Bekenstein argued that the entropy must be proportional to the horizon area itself, the first power and no other, and that single demand sets $b = 1$, hence $k = 1$, collapsing the family to

$$S = \kappa \frac{k_B A c^3}{h G}, \quad \kappa = \frac{\pi}{2},$$

the only dimensionless number, κ , still standing beyond the reach of any argument about units. The progression has run its course: where the length was determined and the density overdetermined, the entropy is underdetermined, and the null direction along which it is free is the count of Planck tiles that a horizon of area A would take to cover, which is the very quantity the entropy turns out to measure.

5 Where the free parameter leads

The three cases are now told, and it would be natural to stop. There is a further step, though, that costs nothing and repays a great deal, because the free direction found in the entropy problem is not a peculiarity of black holes; it is a quantity in its own right, and following it turns the exercise into something a class can carry outward.

Recall the null direction, the combination that could be added freely without disturbing any dimension,

$$\frac{A c^3}{\hbar G} = \frac{A}{\ell_P^2},$$

the horizon area counted in Planck areas. Nothing in the derivation required A to belong to a black hole. Take instead the area of the cosmological horizon, the sphere of radius $R_H = c/H_0$ that light has had time to cross, and the same dimensionless group becomes

$$N = \frac{R_H^2}{\ell_P^2} = \frac{c^5}{\hbar G H_0^2} \approx 7 \times 10^{121}, \quad (1)$$

the number of Planck tiles it would take to cover the observable universe. The reciprocal of this number, $\Pi_0 = \hbar G H_0^2 / c^5 \approx 1.4 \times 10^{-122}$, is worth a moment's attention, because it is the unique dimensionless combination that the four constants c , $\hbar$, G and H_0 admit: their dimensional matrix has rank three and four columns, so by the same rank-and-nullity argument used throughout this essay the nullity is one, and Π_0 spans it.

One number, several disguises. A dimensionless quantity is indifferent to the units it is computed in, so Π_0 can be moved between bases without being removed. Measuring in Planck units, where $c = \hbar = G = 1$, it appears as $H_0^2 \sim 10^{-122}$, the statement that the universe expands slowly on the Planck clock. Measuring instead in a cosmological basis where $c = \hbar = H_0 = 1$, the same number appears as $G \sim 10^{-122}$, the statement that gravity is weak on cosmological scales. Through the de Sitter relation $H_0^2 = \Lambda c^2 / 3$ it appears again as $\Lambda \ell_P^2 \sim 10^{-122}$, the small cosmological constant. These are one fact wearing three costumes, and a class that has followed the elimination can see why: a change of basis relocates a dimensionless number among the constants, and can never dissolve it.

The pedagogical payoff. The entropy problem was left with a free parameter and a physical law brought in from outside to fix it. The free parameter has now been identified as the tile count N , and the same count reappears as the capacity of the cosmological horizon: the Bekenstein and holographic bounds on the entropy of a Hubble sphere both evaluate to $\pi N k_B \approx 2 \times 10^{122} k_B$. So the under-determined case is the richest of the three, rather than the weakest. Where the Planck length gave a unique answer and the density a consistency check, the entropy gave a direction, and that direction is the largest pure number in physics.

System	Behaviour	What it yields
Planck length	square, full rank	a unique scale
Cosmic density	overdetermined, consistent	a redundancy check
Black-hole entropy	underdetermined	a free direction, N

For a class, the sequence has a satisfying shape: solve a system, find it short of one equation, and discover that the missing equation is where the physics lives; then find that the freedom itself, followed outward, is a number describing the universe.

6 The ladder, and a word to mathematics

Read in sequence the three quantities are a tour of what a linear system can do, and the methods they call for are the rungs by which a student learns to handle such systems: back-substitution while the equations stay small and obliging, Gaussian elimination once order can no longer be trusted, and the reading of the pivots, into rank where a column claims one and into freedom where a column does not, when the answer is a family rather than a point.

The appeal to colleagues in mathematics is therefore an inviting one, since physics here is asking for nothing remedial; it offers instead a setting in which simultaneous equations, elimination and the geometry of solution spaces arrive already carrying the weight of fundamental scales, and in which the difference between a unique solution, a redundant equation and a free parameter is made visible on objects as compelling as the smallest length, the emptiness of space, and the entropy of a black hole.

Taught alongside one another, the two subjects let a pupil watch a piece of pure linear algebra reach its limit and a piece of physics step in to finish the work, which is as faithful a picture of how the science is actually done as any first course can hope to give.

Colophon. Drafted with the assistance of a large language model, Anthropic's Claude, used in the manner one might cite a computer algebra system, as an instrument working under the author's direction and to the author's argument.