

Landmark-based Registration



Aalto-yliopisto
Aalto-universitetet
Aalto University

Medical Image Analysis

Koen Van Leemput

Image registration

Combine information contained in different scans

Images need to be spatially aligned!





Image registration



Combine information contained in different scans

Images need to be spatially aligned!






Example: PET/MR

Before registration...





Example: PET/MR

... after registration






Example: CT/MR

Before registration...



Example: CT/MR

... after registration




Example: longitudinal scans

Patient with multiple sclerosis (MS)




Example: longitudinal scans

Patient with multiple sclerosis (MS)




Example: population studies





**average of 152
individuals**


International
Consortium for Brain
Mapping



Example: population studies




Example: population studies



**average of 152
individuals**

International
Consortium for Brain
Mapping




Aalto-yliopisto
Aalto-universitetet
Aalto University


Coordinate systems

For each image, there are two coordinate systems:

- ✓ Voxel coordinates $\mathbf{v} = (v_1, v_2, v_3)^T$ (integer indices)
- ✓ World coordinates $\mathbf{x} = (x_1, x_2, x_3)^T$ (in mm)



Coordinate systems




Conversion: $\mathbf{x} = \mathbf{A}\mathbf{v} + \mathbf{t}$

$$\mathbf{A} = \begin{pmatrix} -0.8923 & -0.0802 & -0.3732 \\ -0.0850 & 0.8921 & 0.3528 \\ -0.0612 & -0.0696 & 4.9512 \end{pmatrix}$$


$$\mathbf{t} = \begin{pmatrix} 129.2834 \\ -98.7363 \\ -27.6911 \end{pmatrix}$$

World coordinates = convention

Important to know which convention is used!



Right – Anterior – Superior
(RAS)



Left – Posterior – Superior
(LPS)



Aalto-yliopisto
Aalto-universitetet
Aalto University


Homogeneous coordinates

Vectors are augmented with a 1 at the end


✓ **Idea:** Rewrite $\mathbf{x} = \mathbf{A}\mathbf{v} + \mathbf{t}$, i.e.,
$$\begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = \begin{pmatrix} a_{1,1} & a_{1,2} & a_{1,3} \\ a_{2,1} & a_{2,2} & a_{2,3} \\ a_{3,1} & a_{3,2} & a_{3,3} \end{pmatrix} \begin{pmatrix} v_1 \\ v_2 \\ v_3 \end{pmatrix} + \begin{pmatrix} t_1 \\ t_2 \\ t_3 \end{pmatrix}$$

as:
$$\begin{pmatrix} x_1 \\ x_2 \\ x_3 \\ 1 \end{pmatrix} = \underbrace{\begin{pmatrix} a_{1,1} & a_{1,2} & a_{1,3} & t_1 \\ a_{2,1} & a_{2,2} & a_{2,3} & t_2 \\ a_{3,1} & a_{3,2} & a_{3,3} & t_3 \\ 0 & 0 & 0 & 1 \end{pmatrix}}_{\mathbf{M}} \begin{pmatrix} v_1 \\ v_2 \\ v_3 \\ 1 \end{pmatrix}$$

✓ **Benefit:** map voxel indices using only matrix multiplications




Spatial transformations



“fixed” image


$$\mathbf{x} = (x_1, \dots, x_D)^T$$



“moving” image

$$\mathbf{y} = (y_1, \dots, y_D)^T$$


Spatial transformations



"fixed" image

$$\mathbf{x} = (x_1, \dots, x_D)^T$$


$$\mathbf{y}(\mathbf{x}, \mathbf{w}) = \begin{pmatrix} y_1(\mathbf{x}, \mathbf{w}) \\ \vdots \\ y_D(\mathbf{x}, \mathbf{w}) \end{pmatrix}$$



"moving" image

$$\mathbf{y} = (y_1, \dots, y_D)^T$$


Spatial transformations



"fixed" image

$$\mathbf{x} = (x_1, \dots, x_D)^T$$

$$\mathbf{y}(\mathbf{x}, \mathbf{w}) = \begin{pmatrix} y_1(\mathbf{x}, \mathbf{w}) \\ \vdots \\ y_D(\mathbf{x}, \mathbf{w}) \end{pmatrix}$$



"moving" image




$$\mathbf{y} = (y_1, \dots, y_D)^T$$




$$y_d(\mathbf{x}, \mathbf{w})$$

controls how points \mathbf{x} in the fixed image
move along the d -th direction in the moving image
as the parameters \mathbf{w} are varied

Spatial transformations





Spatial transformations



Affine transformation

$$\mathbf{y}(\mathbf{x}, \mathbf{w}) = \mathbf{A}\mathbf{x} + \mathbf{t}$$



$$\mathbf{A} = \begin{pmatrix} a_{1,1} & a_{1,2} \\ a_{2,1} & a_{2,2} \end{pmatrix} \quad \text{and} \quad \mathbf{t} = \begin{pmatrix} t_1 \\ t_2 \end{pmatrix}$$



Affine transformation

$$\mathbf{y}(\mathbf{x}, \mathbf{w}) = \mathbf{A}\mathbf{x} + \mathbf{t}$$

$$\mathbf{A} = \begin{pmatrix} a_{1,1} & a_{1,2} \\ a_{2,1} & a_{2,2} \end{pmatrix} \quad \text{and} \quad \mathbf{t} = \begin{pmatrix} t_1 \\ t_2 \end{pmatrix}$$



$$y_d(\mathbf{x}, \mathbf{w})$$

controls how points \mathbf{x} in the fixed image move along the d -th direction in the moving image as the parameters \mathbf{w} are varied



$$y_d(\mathbf{x}, \mathbf{w}_d) = t_d + a_{d,1}x_1 + \dots + a_{d,D}x_D$$

$$\mathbf{w}_d = (t_d, a_{d,1}, \dots, a_{d,D})^T$$

$$\mathbf{w} = (\mathbf{w}_1^T, \dots, \mathbf{w}_D^T)^T$$

Affine transformation



$$\mathbf{y}(\mathbf{x}, \mathbf{w}) = \mathbf{A}\mathbf{x} + \mathbf{t}$$



$$\mathbf{A} = \begin{pmatrix} 1.0 & 0.0 \\ 0.0 & 1.0 \end{pmatrix}, \mathbf{t} = \begin{pmatrix} 23 \\ 0 \end{pmatrix}$$

Affine transformation



$$\mathbf{y}(\mathbf{x}, \mathbf{w}) = \mathbf{A}\mathbf{x} + \mathbf{t}$$



$$\mathbf{A} = \begin{pmatrix} 1.4 & 0.0 \\ 0.0 & 1.0 \end{pmatrix}, \mathbf{t} = \begin{pmatrix} 23 \\ 0 \end{pmatrix}$$

Affine transformation



$$\mathbf{y}(\mathbf{x}, \mathbf{w}) = \mathbf{A}\mathbf{x} + \mathbf{t}$$



$$\mathbf{A} = \begin{pmatrix} 1.4 & 0.5 \\ 0.0 & 1.0 \end{pmatrix}, \quad \mathbf{t} = \begin{pmatrix} 23 \\ 0 \end{pmatrix}$$

Affine transformation



$$\mathbf{y}(\mathbf{x}, \mathbf{w}) = \mathbf{A}\mathbf{x} + \mathbf{t}$$



$$\mathbf{A} = \begin{pmatrix} 1.0 & 0.0 \\ 0.0 & 1.0 \end{pmatrix}, \mathbf{t} = \begin{pmatrix} 23 \\ 6 \end{pmatrix}$$

Affine transformation



$$\mathbf{y}(\mathbf{x}, \mathbf{w}) = \mathbf{A}\mathbf{x} + \mathbf{t}$$



$$\mathbf{A} = \begin{pmatrix} 1.4 & 0.5 \\ 0.2 & 0.9 \end{pmatrix}, \mathbf{t} = \begin{pmatrix} 23 \\ 6 \end{pmatrix}$$


Rigid transformation

$$\mathbf{y}(\mathbf{x}, \mathbf{w}) = \mathbf{R}\mathbf{x} + \mathbf{t}, \quad \mathbf{R}^T \mathbf{R} = \mathbf{I} \quad \text{and} \quad \det(\mathbf{R}) = 1$$



Rigid transformation

Task: why do $\mathbf{R}^T \mathbf{R} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$
and $\det(\mathbf{R}) = 1$ impose a rotation?




Rigid transformation

$$\|\mathbf{r}_1\| = \sqrt{\mathbf{r}_1^T \mathbf{r}_1} = 1$$

Task: why do $\mathbf{R}^T \mathbf{R} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$
and $\det(\mathbf{R}) = 1$ impose a rotation?

$$\|\mathbf{r}_2\| = \sqrt{\mathbf{r}_2^T \mathbf{r}_2} = 1$$


$$\mathbf{r}_1^T \mathbf{r}_2 = 0$$



Rigid transformation

Task: why do $\mathbf{R}^T \mathbf{R} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$
and $\det(\mathbf{R}) = 1$ impose a rotation?


?



Rigid transformation


Task: why do $\mathbf{R}^T \mathbf{R} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$
and $\det(\mathbf{R}) = 1$ impose a rotation?

?



Landmark-based registration

- ✓ Manually annotate N corresponding points in two images:



- ✓ Register the images by minimizing the distance between matching point pairs:

$$E(\mathbf{w}) = \sum_{n=1}^N \|\mathbf{y}_n - \mathbf{y}(\mathbf{x}_n, \mathbf{w})\|^2$$

Landmark-based registration

Applied to affine registration: $E(\mathbf{w}) = \sum_{n=1}^N \|\mathbf{y}_n - \mathbf{A}\mathbf{x}_n - \mathbf{t}\|^2$

Task 1: if $\mathbf{A} = \mathbf{I}$, what is \mathbf{t} ?

Hint: remember that $\|\mathbf{a} - \mathbf{b}\|^2 = \sum_{d=1}^D (a_d - b_d)^2$

Task 2: in the general case, what are \mathbf{A} and \mathbf{t} ?

Landmark-based registration

Applied to affine registration: $E(\mathbf{w}) = \sum_{n=1}^N \|\mathbf{y}_n - \mathbf{A}\mathbf{x}_n - \mathbf{t}\|^2 = \sum_{n=1}^N \sum_d^D (y_{n,d} - t_d - \sum_{d'=1}^D x_{n,d'} a_{d,d'})^2$


Task 1: if $\mathbf{A} = \mathbf{I}$, what is \mathbf{t} ?

Hint: remember that $\|\mathbf{a} - \mathbf{b}\|^2 = \sum_{d=1}^D (a_d - b_d)^2$

Task 2: in the general case, what are \mathbf{A} and \mathbf{t} ?

Landmark-based registration

Applied to affine registration: $E(\mathbf{w}) = \sum_{n=1}^N \|\mathbf{y}_n - \mathbf{A}\mathbf{x}_n - \mathbf{t}\|^2 = \sum_{n=1}^N \sum_d^D (y_{n,d} - t_d - \sum_{d'=1}^D x_{n,d'} a_{d,d'})^2$



Task 1: if $\mathbf{A} = \mathbf{I}$, what is \mathbf{t} ?

Hint: remember that $\|\mathbf{a} - \mathbf{b}\|^2 = \sum_{d=1}^D (a_d - b_d)^2$

Task 2: in the general case, what are \mathbf{A} and \mathbf{t} ?

Landmark-based registration

Applied to affine registration: $E(\mathbf{w}) = \sum_{n=1}^N \|\mathbf{y}_n - \mathbf{A}\mathbf{x}_n - \mathbf{t}\|^2$

Task 1: if $\mathbf{A} = \mathbf{I}$, what is \mathbf{t} ?

Hint: remember that $\|\mathbf{a} - \mathbf{b}\|^2 = \sum_{d=1}^D (a_d - b_d)^2$

Task 2: in the general case, what are \mathbf{A} and \mathbf{t} ?

$$\begin{aligned} &= \sum_{n=1}^N \sum_d^D (y_{n,d} - t_d - \sum_{d'=1}^D x_{n,d'} a_{d,d'})^2 \\ &= \sum_d^D \sum_{n=1}^N (y_{n,d} - t_d - \sum_{d'=1}^D x_{n,d'} a_{d,d'})^2 \end{aligned}$$

Landmark-based registration

Applied to affine registration: $E(\mathbf{w}) = \sum_{n=1}^N \|\mathbf{y}_n - \mathbf{A}\mathbf{x}_n - \mathbf{t}\|^2 = \sum_{n=1}^N \sum_d^D (y_{n,d} - t_d - \sum_{d'=1}^D x_{n,d'} a_{d,d'})^2$

Task 1: if $\mathbf{A} = \mathbf{I}$, what is \mathbf{t} ?

Hint: remember that $\|\mathbf{a} - \mathbf{b}\|^2 = \sum_{d=1}^D (a_d - b_d)^2$


Task 2: in the general case, what are \mathbf{A} and \mathbf{t} ?

$$= \sum_{n=1}^N \sum_d^D (y_{n,d} - t_d - \sum_{d'=1}^D x_{n,d'} a_{d,d'})^2$$
$$= \sum_d^D \sum_{n=1}^N (y_{n,d} - t_d - \sum_{d'=1}^D x_{n,d'} a_{d,d'})^2$$


$$\begin{pmatrix} t_d \\ a_{d,1} \\ \vdots \\ a_{d,D} \end{pmatrix} = (\mathbf{X}^T \mathbf{X})^{-1} \mathbf{X}^T \begin{pmatrix} y_{1,d} \\ \vdots \\ y_{N,d} \end{pmatrix}$$

where $\mathbf{X} = \begin{pmatrix} 1 & x_{1,1} & \cdots & x_{1,D} \\ 1 & x_{2,1} & \cdots & x_{2,D} \\ \vdots & \vdots & \ddots & \vdots \\ 1 & x_{N,1} & \cdots & x_{N,D} \end{pmatrix}$


Landmark-based registration



fixed image



moving image




fixed + moving

Before registration




Aalto-yliopisto
Aalto-universitetet
Aalto University


Landmark-based registration



fixed image



moving image



fixed + moving



After registration



Aalto-yliopisto
Aalto-universitetet
Aalto University

Landmark-based registration

Applied to rigid registration: $E(\mathbf{w}) = \sum_{n=1}^N \|\mathbf{y}_n - \mathbf{R}\mathbf{x}_n - \mathbf{t}\|^2$



- ✓ Constraints $\mathbf{R}^T \mathbf{R} = \mathbf{I}$ and $\det(\mathbf{R}) = 1$ make the math much more complicated!
- ✓ Solution:

$$\mathbf{R} = \mathbf{V}\mathbf{U}^T, \quad \sum_{n=1}^N \tilde{\mathbf{x}}_n \tilde{\mathbf{y}}_n^T = \mathbf{U}\mathbf{\Sigma}\mathbf{V}^T, \quad \mathbf{U}^T \mathbf{U} = \mathbf{I}, \quad \mathbf{V}^T \mathbf{V} = \mathbf{I}$$

$$\mathbf{t} = \bar{\mathbf{y}} - \mathbf{R}\bar{\mathbf{x}},$$

$$\text{where } \tilde{\mathbf{x}}_n = \mathbf{x}_n - \bar{\mathbf{x}} \text{ and } \tilde{\mathbf{y}}_n = \mathbf{y}_n - \bar{\mathbf{y}}$$

$$\text{with } \bar{\mathbf{x}} = \frac{1}{N} \sum_{n=1}^N \mathbf{x}_n \text{ and } \bar{\mathbf{y}} = \frac{1}{N} \sum_{n=1}^N \mathbf{y}_n$$

(“flip” a column of \mathbf{R} if $\det(\mathbf{R}) = -1$)