



[Mol Cell](#). Author manuscript; available in PMC 2016 May 21.

PMCID: PMC4441954

Published in final edited form as:

NIHMSID: NIHMS689107

[Mol Cell. 2015 May 21; 58\(4\): 598–609.](#)

PMID: 26000845

doi: [10.1016/j.molcel.2015.05.005](https://doi.org/10.1016/j.molcel.2015.05.005)

Advances and Applications of Single Cell Sequencing Technologies

[Yong Wang¹](#) and [Nicholas E. Navin^{1,2,*}](#)

¹Department of Genetics, University of Texas, MD Anderson Cancer Center

²Department of Bioinformatics and Computational Biology, University of Texas, MD Anderson Cancer Center

*Corresponding Author: Nicholas E. Navin, Ph.D. nnavin@mdanderson.org

[Copyright notice](#)

[Publisher's Disclaimer](#)

Abstract

Single cell sequencing (SCS) has emerged as a powerful new set of technologies for studying rare cells and delineating complex populations. Over the past 5 years, SCS methods for DNA and RNA have had a broad impact on many diverse fields of biology, including microbiology, neurobiology, development, tissue mosaicism, immunology and cancer research. In this review, we will discuss SCS technologies and applications, as well as translational applications in the clinic.

Introduction

The fundamental unit of an organism is a single cell. *Homo sapiens* are composed of approximately 3.72×10^{13} single cells that live harmoniously in tissues among their neighbors ([Bianconi et al., 2013](#)). However in diseases such as cancer, the greed and avarice of a single cell can lead to the downfall of the entire organism. Despite the complexity of tissues, most genomic studies to date have focused on analyzing bulk tissue samples, which are composed of millions of cells. In these averaged datasets, it is difficult to resolve cell-to-cell variations and identify rare cells that may play an important role in disease progression. The recent development of single-cell sequencing (SCS) methods has led to a paradigm shift in the field of genomics, away from bulk tissue analysis, and towards detailed and comprehensive studies of individual cells.

Our fascination with single cells dates back to the invention of the first microscopes in the 1660's, which allowed early microscopists to observe single prokaryotic cells moving around in droplets of water. Subsequent work by early pathologists, such as Rudolf Virchow, in the late 1850's established the link between abnormalities in single cells and human diseases. In the late 1900's the development of cell staining techniques and cytological methods galvanized the field, enabling scientists to directly visualize genetic differences on chromosomes in single cells. However, many cytogenetic and immunostaining methods were limited to measuring targeted genes and proteins. In the 1990's quantitative microarray technologies were developed for measuring genome-wide DNA and RNA information, but required too much input material for single cell analysis. Although PCR methods had

been developed, they were only capable of amplifying small targeted regions of the genome. To overcome this limitation whole-transcriptome-amplification (WTA) ([Van Gelder et al., 1990](#)) and whole-genome-amplification (WGA) ([Dean et al., 2002](#); [Telenius et al., 1992](#)) methods were developed to amplify genome-wide DNA and RNA. Another important milestone occurred in 2005 with the development of the first next-generation sequencing (NGS) technologies, which enabled genome-wide sequencing of DNA and RNA ([Mardis, 2011](#)).

The culmination of these technologies led to the invention of the first genomewide single-cell DNA ([Navin et al., 2011](#)) and RNA ([Tang et al., 2009](#)) sequencing methods for mammalian cells. These initial studies (and work by other groups) led to the establishment of a new field of biology: single cell sequencing (SCS). The field has shown tremendous growth over the last 5 years ([Figure 1A](#)) and impacted many diverse areas of biological research ([Figure 1B–C](#), [Supplementary Table 1](#)). In this review, we will discuss the advances and limitations of SCS technologies, and the myriad of applications that they have had in biological research and medicine.

[Open in a separate window](#)

Figure 1

Timeline of Milestones in Single Cell Sequencing

(A) Timeline of SCS Milestones (B) Histogram of the growing number of publications in SCS over the past 5 years (C) Prevalence of publications categorized by fields.

Single Cell Isolation Methods

In order to sequence a single cell, it must first be captured. While the methods for isolating single cells from abundant populations have been well-established, the isolation of rare single cells (<1%) remains a formidable technical challenge. To isolate a single cell randomly from an abundant population, several approaches can be employed: mouth pipetting, serial dilution, robotic micromanipulation, flow-assisted cell sorting (FACS) and microfluidic platforms ([Table 1](#)). Many of these approaches require cells or nuclei in suspension, and therefore cannot preserve their spatial context in tissues. This limitation can be overcome using Laser-capture-microdissection (LCM), which can also be used to isolate rare cells. In contrast, the isolation of rare single cells (< 1%) is far more challenging. Many commercial platforms have been developed for isolating circulating tumor cells (CTCs), which occur at very low frequencies (1 in 1 million) in the blood of cancer patients ([Cristofanilli et al., 2004](#)). The CellSearch system is an FDA approved clinical system that uses magnets with ferrofluid nanoparticles conjugated to antibodies for EpCAM and CD45 to isolate CTCs ([Yu et al., 2011](#)). Nagrath et al. developed another method that uses a nanopost microchip technology with EpCAM antibodies ([Nagrath et al., 2007](#)). The Magsweeper (Illumina Inc.) is a technology that involves dipping a rotating magnet with EpCAM antibodies to isolate CTCs ([Powell et al., 2012](#)). Other methods are more generally applicable to rare cell populations. The DEP-Array system (Silicon Biosciences) uses a microchip with dielectrophoretic cages to navigate individual cells by charge ([Altomare et al., 2003](#)). The CellCeptor (Automated Lab Solutions) uses robotic micromanipulation capillary system to identify single cells for isolation ([Choi et al., 2010](#)). An alternative approach uses nanofilters to isolate

rare cells by size exclusion rather than surface markers (Adams et al., 2014). The advantages and limitations of these methods are summarized in [Table 1](#) and have been reviewed in detail in other review articles (Navin and Hicks, 2011; Navin, 2014; Shapiro et al., 2013).

Table 1**Single-Cell Isolation Methods for Abundant and Rare Populations**[Open in a separate window](#)

This table summarizes the advantages and disadvantages of single-cell isolation methods from abundant populations and rare populations.

Single Cell DNA Sequencing Methods

The development of DNA SCS methods has proven to be more challenging than RNA. This is due to the fact that a single cell contains only 2 copies of each DNA molecule, but thousands of copies of most RNA molecules. The limited amount of input material for WGA results in a number of technical errors, including: coverage non-uniformity, allelic dropout (ADO) events, false-positive (FP) errors and false-negative (FN) errors ([Table 2](#)). The most salient technical errors occur during the initial rounds of amplification and are subsequently propagated by all daughter molecules. FP errors accumulate at random sites due to the infidelity of the WGA polymerase and lead to single nucleotide errors (Dean et al., 2002; Lasken, 2007). However, by far the greatest source of WGA error comes from allelic dropout events at 10–50% of the mutation sites (Fiegler et al., 2007; Hou et al., 2012; Lasken, 2007; Talseth-Palmer et al., 2008; Zong et al., 2012).

Table 2**Technical Errors Associated with Single-Cell Sequencing**[Open in a separate window](#)

This table lists the common technical errors that arise during WGA and WTA in single-cell sequencing experiments.

Importantly, WGA is not a single technique, but encompasses a wide variety of experimental methods. The most common WGA methods are degenerative-oligonucleotide-PCR (DOP-PCR) and multiple-displacement-amplification (MDA) ([Figure 2A–B](#)). DOP-PCR generates low physical coverage (~10%) of a single cell genome, but accurately retains copy number levels during WGA. In the first SCS method developed for genomic DNA, DOP-PCR was combined with flow-sorting of nuclei and NGS to generate high-resolution copy number profiles from single mammalian cells (Baslan et al., 2012; Navin et al., 2011). However, the low physical coverage of DOP-PCR makes it a poor approach

for measuring mutations at base-pair resolution. MDA using either the Phi29 or Bst polymerases has been widely reported to achieve high physical coverage (>90%) from a single cell genome or exome (Hou et al., 2012; Xu et al., 2012; Yong Wang, 2014; Zong et al., 2012) (Figure 2B). However, MDA generates non-uniform coverage and causes distortions in read depth making it a poor method to measure DNA copy number (Navin, 2014). Phi29 is the ideal polymerase for MDA reactions, since it has a very low FP error rate (1e-7) compared to Bst (1e-5), which does not have proofreading activity (Dean et al., 2002; Lasken, 2013). Another DNA SCS method is multipleannealing- and-looping-based-amplification-cycles (MALBAC), which uses the Bst polymerase to form circular DNA fragments followed by adapter ligation PCR (Zong et al., 2012). This method can obtain both copy number information and single nucleotide variants (SNVs), but generates extremely high FP error rates, making it more suitable for copy number profiling. Another method, called NUC-SEQ or single nucleus exome sequencing (SNES) takes advantage of G2/M nuclei which have duplicated the amount of genomic DNA in a single cell (12 picograms) prior to MDA, which reduces many technical error rates during single cell sequencing of exomes and genomes (Yong Wang, 2014)(Leung et al. 2015).

[Open in a separate window](#)

Figure 2

WGA and WTA Methods for Single Cell Sequencing

(A–B) Whole-genome-amplification methods. (C–D) Whole-transcriptome-amplification methods. (A) Degenerative-Oligonucleotide-Primer PCR (B) Multiple-displacementamplification. (C) oligo dT-Anchor Approach (D) Template switching

After WGA the amplified DNA is used to construct libraries for NGS. While several sequencing platforms are available, Illumina has emerged as the primary tools in most studies due to low cost per base, high-throughput and low error rates. To further save costs and increase throughput, single cell libraries are often barcoded and pooled together for multiplexed sequencing. In many studies, the barcoded libraries are used for targeted capture (exome or gene panels) to sequence only regions of interest and obtain higher coverage depth in these areas. For a more comprehensive review of single cell DNA sequencing methods please refer to the following articles (Blainey and Quake, 2014; de Bourcy et al., 2014; Navin and Hicks, 2011; Navin, 2014; Van Loo and Voet, 2014).

Single Cell RNA Sequencing Methods

The development of single cell RNA sequencing methods has shown considerable progress over the past 5 years. To sequence a single cell transcriptome, the RNA must first be amplified by whole-transcriptome-amplification (WTA). This step is necessary because a typical mammalian cell contains only 10 picograms of total RNA and 0.1 picograms of mRNA. Initial WTA methods utilized the T7 RNA polymerase to perform linear amplification of cDNA by *in vitro* transcription (IVT) (Van Gelder et al., 1990). These methods were further developed using oligo d(T) primers conjugated to adapter sequences for reverse transcription and selective amplification of polyadenylated mRNA by PCR (Tang et al., 2009) (Figure 2C). However, these WTA methods were plagued by strong 3' mRNA bias. To mitigate this bias, a WTA method called SMART-Seq was developed that amplifies only full-length mRNA transcripts using an Moloney Murine Leukemia Virus (MMLV) reverse transcriptase

([Ramskold et al., 2012](#)). MMLV has both template-switching and terminal transferase activity, which results in the addition of nontemplated cytosine residues to the 5' end of the cDNA. By adding a poly-G template with an adapter sequence, MMLV can switch templates and transcribe the other strand, leading to full-length cDNA transcripts that are amplified by PCR ([Figure 2D](#)). Another technical artifact of single-cell RNA sequencing is amplification bias, in which mRNAs levels are distorted during WTA. To reduce these errors, a recent method developed unique molecular indexes (UMIs) to label the original pool of RNA molecules prior to WTA ([Islam et al., 2014](#)). After WTA, the resulting cDNA libraries are barcoded and pooled for multiplexed NGS. For a more detailed discussion of RNA SCS methods please refer to the following review articles ([Macaulay and Voet, 2014](#); [Saliba et al., 2014](#); [Sandberg, 2014](#)).

Single Cell Epigenomic Sequencing Methods

Epigenomic profiling of single cells remains to be one of the greatest technical challenges in the field. The problem is that standard epigenomic sequencing methods require a pool of DNA that is split into two separate fractions for treatment with bisulfide or methylation restriction enzymes prior to sequencing. The other technical barrier is that epigenetic DNA modifications cannot be amplified with DNA polymerases. Despite these technical hurdles, two recent studies have made initial progress. The Hi-C approach was recently adapted for single cell profiling at megabase resolution to identify physical chromatin interactions in single cells ([Nagano et al., 2013](#)). In another study, reduced single cell representation bisulfite sequencing (scRRBS) was developed to measure cytosine methylation modifications at 1.5 million CpG sites in a single cell, which is equivalent to about 10% of the genome ([Guo et al., 2013](#)). While these studies are clearly pioneering, they were also challenged by limited coverage (2.5% and 10%), low resolution and many technical errors. However, future refinement of these technologies are likely to lead to accurate whole-genome epigenomic profiling of single cells.

Distinguishing Technical Errors from Biological Variations

At the current state of technology, most SCS methods introduce extensive technical errors and variability into datasets. Unfortunately, naïve users often interpret these technical errors as extensive biological heterogeneity. To eliminate these errors and confirm that a mutation or transcript is truly heterogeneous in a population of cells, orthogonal validation using a targeted approach is critical. To validate heterogeneous DNA variants or mutations, targeted validation can be performed using deep-sequencing with molecular barcodes ([Kennedy et al., 2014](#); [Schmitt et al., 2012](#); [Wang et al., 2014](#)) or digital droplet PCR (Bio-Rad, Raindance Inc.). To validate heterogeneous RNA expression changes, targeted validation can be performed with single cell RT-qPCR or with digital droplet PCR. Unfortunately, many published studies to date have incorrectly reported extensive biological heterogeneity without orthogonal validation, which is more likely to be explained by technical errors.

In summary, there has been tremendous progress in the development of single cell DNA and RNA sequencing methods. However, all SCS methods generate technical errors during WGA and WTA and thus orthogonal validation using a targeted approach is critical at the current state of technology. We now turn to a detailed discussion of the many broad fields of biology that have been impacted by SCS methods over the last 5 years ([Figure 3](#)).

Figure 3

Broad Applications of SCS in Biological and Biomedical Research

Panels illustrating the diverse fields of biology that have been impacted by SCS technologies over the past 5 years. Image credits: neurobiology, Zeynep Saygin (Cell Picture Show); germline transmission, Wang and Navin; organogenesis, Mikael Häggström (Wikimedia Commons); cancer biology, NIH; clinical diagnostics, Wang and Navin; immunology, Olivier Schwartz and the Electron Microscopy Core Facility, Institut Pasteur (Cell Picture Show); microbiology, NIAID; tissue mosaicism, Wang and Navin; embryology, Seth Ruffins, Russell Jacobs, and the Caltech MRI Atlas of Mouse Development (Cell Picture Show); prenatal genetic diagnosis, Shutterstock. All images used with permission.

Microbiology

A formidable challenge in microbiology has been that over 99% of microbial species on planet earth cannot be cultured and expanded in the lab ([Ishoey et al., 2008](#); [Lasken and McLean, 2014](#)). Single-cell DNA and RNA sequencing methods provide a powerful new approach to resolve microbial genomes and delineate cell-to-cell diversity within diverse populations. However, bacteria and other microorganisms often have only femtograms of DNA and RNA, making it even more challenging to amplify than mammalian cells ([Lasken, 2007](#)). In an early study, MDA was used to amplify DNA from the marine *cyanobacterium Prochlorococcus* for pyrosequencing and *de novo* assembly ([Rodrigue et al., 2009](#)). In another study Woyke et al. used FACS and MDA to perform NGS and assemble two marine flavobacteria genomes to 90% coverage ([Woyke et al., 2009](#)). Blainey et al. also used MDA to sequence and assemble the genomes of 5 single cells from an Ammonia-oxidizing archaea ([Blainey et al., 2011](#)). Another study performed SCS of five segmented filamentous bacterial cells to gain insight into their life cycles ([Pamp et al., 2012](#)). While initial studies were often limited to sequencing just a few microbial cells, a subsequent large-scale study was conducted on 201 uncultivated archaeal and bacterial cells from nine diverse habitats. In this study SCS revealed 29 uncharted branches of the tree of life, revealing ‘microbial dark matter’ and challenging the canonical three domains of life ([Rinke et al., 2013](#)). These studies show that SCS is complimentary to metagenomic deep-sequencing methods, and can open up new avenues of investigation into microbial genomes that cannot be cultured in the lab.

Neurobiology

Neurons represent one of the most morphologically diverse populations of cells. Traditional classification has relied mainly on morphological features ([De Carlos and Borrell, 2007](#)), however single-cell RNA sequencing provides a powerful unbiased approach to classify neurons based on their transcriptional profiles. In a study by Qiu et al. single neuron RNA-sequencing was combined with electrophysiology to obtain transcriptional profiles from embryonic mouse hippocampus and neocortical neurons ([Qiu et al., 2012](#)). In another study, single cell RNA-seq was performed *in situ* in spatially defined neuronal regions, which identified cell-to-cell transcriptional variation in hippocampal neurons ([Lovatt et al., 2014](#)). Pollen et al. used low-coverage single cell RNA sequencing and microfluidics to analyze single cells from 11 brain populations, and identified *Notch* signaling as an

important regulator of brain development ([Pollen et al., 2014](#)). In another study, Usoskin et al. used single-cell RNA sequencing to profile 622 sensory neurons in mice, which revealed 11 novel expression classes of sensory neuron cell types ([Usoskin et al., 2014](#)).

Several studies have also begun to investigate DNA heterogeneity in neurons. SCS was recently used to study LINE-1 retrotransposition in the cerebral cortex ([Ervony et al., 2012](#)) and found that each cortex neuron had an average of 0.6 somatic insertions events. In another study, SNS ([Baslan et al., 2012](#); [Navin et al., 2011](#)) was used to identify copy number variants (CNVs) in neurons from three normal and two pathological brain samples ([Cai et al., 2014](#)). The authors reported that large (>1mb) clonal CNVs arose in about 5% of neurons during normal development. In contrast, another study used SNS to profile neuronal copy number diversity in the prefrontal cortex of postmortem brains, which identified many *de novo* CNVs in neurons that were not clonal between different single cells ([McConnell et al., 2013](#)). In another study, SCS using microwells identified copy number changes in a normal postmortem brain and a patient with Down syndrome ([Gole et al., 2013](#)). These initial studies show that SCS provides a novel approach to classify neuronal cell types and identify an unexpected amount of DNA diversity in neuronal populations.

Tissue Mosaicism

The traditional view of somatic tissues is that single normal cells have identical genomes, however this dogma is beginning to be challenged by increasing evidence of genetic mosaicism in normal tissues that arises during normal development ([Biesecker and Spinner, 2013](#)). To date most studies have analyzed bulk tissue samples, and therefore much controversy exists over the prevalence of mosaic mutations and whether they can simply be explained by technical error. SCS methods provide a novel approach to resolve cell-to-cell variations in normal tissues at an unprecedented genomic resolution. SNS was recently used to identify *de novo* CNVs in 13–41% of the neurons in the frontal cortex of postmortem brains, suggesting that CNV mosaic events are common in cortical neurons ([McConnell et al., 2013](#)). This unexpected amount of copy number diversity has previously not been appreciated in the brain. However, a recent study using the same SCS methods (SNS) challenged these data, by suggesting that somatic CNVs are extremely rare in neurons and other normal tissues ([Knouse et al., 2014](#)). In this study 96 single neurons were sequenced from mice and only a single somatic CNV was identified in one neuron. The authors also examined 89 single cells from 4 human patients frontal lobes, and found only 2 cells with aneuploid rearrangements (2.2%). In skin, the authors detected aneuploidy in only 2.7% of mouse keratinocytes and none in human cells. In liver cells they profiled 100 hepatocytes and found only 4% aneuploid cells. Thus, while both studies showed that copy number mosaicism is likely to exist in normal tissues, there is much debate regarding the prevalence of these rearrangements, and whether they might play an important role in human diseases.

Germline Transmission

Sperm cells and oocytes are single cells that fuse to form a zygote and transmit genomic material and evolution has engineered this process to generate genetic variation. Single-cell DNA sequencing provides a novel approach to study the mechanisms that generate germline variation. In one of the first studies on this topic, single sperm cells were sequenced, which revealed an average of 22.8 recombination events, 5–15 gene conversion events and 25–36 *de novo* mutations in each sperm cell ([Wang et al., 2012](#)). The authors also calculated copy number profiles, which showed that 7% of the single sperm cells had aneuploid genomes. Consistent with this study, another group used low-coverage whole-genome sequencing to delineate haplotypes in single sperm cells from one individual, which revealed an average of 25.3 recombination events per cell ([Kirkness et al., 2013](#)). In another

study, Lu et al. applied MALBAC to sequence single sperm cells from an Asian individual, in which they reported aneuploidy in 4% of the cells and 26 recombination events per single sperm cell ([Lu et al., 2012](#)). While most germline studies have focused on sperm, a recent study used MALBAC to analyze fertilized oocytes ([Hou et al., 2013](#)). In this study oocytes from 8 individual females were analyzed, which identified 43 cross-over events per oocyte, a recombination rate that is 1.63X higher than sperm. Interestingly, this study also reported a much higher rate of aneuploidy in oocytes (17.6%) compared to sperm (4–7%). Taken together, these studies have confirmed previous recombination rates and revealed a striking amount of genomic diversity that arises in germ cells during the transmission of genetic material to offspring.

Embryogenesis

Extensive transcriptional regulation and epigenetic reprogramming occurs during the earliest stages of embryonic development, as the zygote forms the three major cell lineages (endoderm, ectoderm and mesoderm). The genomic regulation of these early events and maintenance of pluripotency has been challenging to study due to the limited amount of input material. To address this problem, RNA SCS was used to analyze transcriptional reprogramming *in vitro* during the transition from the inner cell mass of blastocysts to pluripotent embryonic stem cells ([Tang et al., 2010](#)). In another study, RNA SCS was used to profile single cells from human pre-implantation embryos and embryonic stem cells which detected over 1000 heterogeneous transcripts within the same blastomere ([Yan et al., 2013](#)). In another study, RNA SCS was used to study transcriptome dynamics from oocyte to morula development in human and mouse embryos, which delineated a step-wise progression of pathways that regulate cell cycle, gene regulation, translation and metabolism ([Xue et al., 2013](#)). Another study used single cell bisulfite sequencing to measure cytosine DNA modifications in mouse embryonic stem cells, which showed massive global demethylation during embryonic development ([Guo et al., 2013](#)). Collectively, these studies have begun to dissect the complex transcriptional regulation and epigenomic reprogramming that occurs during the earliest stages of embryogenesis.

Organogenesis

In most tissues, traditional classification of cell types has previously been limited to a few dozen markers that have been used for decades. RNA SCS methods provide a powerful new unbiased approach to perform transcriptional profiling and identify groups of cells that share common expression programs, representing distinct cell types. In the first study to apply this approach RNA SCS was used to analyze lung epithelium development ([Treutlein et al., 2014](#)). From these data, the development of lung progenitor cells were traced as they formed alveolar air sacs that regulate gas exchange. In this study the authors identified hundreds of novel markers for distinguishing the four major cell types and used them to reconstruct cell lineages during alveolar sac differentiation. In another study, RNA SCS was used to analyze gene expression patterns of single cells during kidney development in mice at E11.5, E12.5 and P4 ([Brunskill et al., 2014](#)). These data revealed a multi-lineage priming model in which many genes and pathways were repressed during nephrogenesis, rather than being activated from a ‘blank slate’. These initial studies demonstrate the utility of applying unbiased RNA SCS methods to classify cell types and identify novel markers of cell lineages during organ development.

Immunology

The immune system is broadly classified into the adaptive and innate components, which comprise a large variety of cell types that work together in a concerted fashion to recognize and clear antigens. Although the major immune cell types have been known for decades, there is little known about the transcriptional heterogeneity within cell types in responses to antigens. In one study, RNA SCS was used to analyze mouse bone-marrow derived dendritic cells that were stimulated under different conditions *in vitro* and found that individual cells show variable responses that are mediated by interferon paracrine signaling ([Shalek et al., 2014](#)). In another study, RNA SCS was used to identify bimodal gene expression patterns in bone-marrow-derived dendritic cells stimulated by lipopolysaccharide that was modulated through an interferon feedback circuit ([Shalek et al., 2013](#)). Another study performed unbiased RNA SCS to profile 4000 single cells from mouse spleen in response to antigen activation with LPS which revealed seven classes of immune cells and identified 1575 variable gene responses after antigen activation ([Jaitin et al., 2014](#)). These studies show that unbiased RNA SCS methods can be used to investigate heterogeneous transcriptional responses in immune cells after antigen activation.

Cancer Research

Tumors evolve from single normal cells. During this process the cancer cells accumulate mutations and diversify to form distinct lineages and subpopulations. This intratumor heterogeneity confounds the clinical diagnosis and therapeutic treatment of patients. Clonal diversity is likely to play a key role in tumor progression during processes such as invasion, clonal evolution and metastasis by providing fuel for evolution to select upon. Genomic diversity also enables tumor cell populations to survive selective pressures in the tumor microenvironment, including hypoxia, chemotherapy, immune surveillance and geographic barriers. However, to date studying clonal diversity has been difficult in bulk populations of tumor cells using standard sequencing methods. DNA and RNA SCS methods provide powerful new tools for delineating clonal diversity and understanding the role of rare cells during cancer progression.

To date, most SCS studies of cancer have focused on intratumor heterogeneity and clonal evolution in primary tumors. The first study used SNS to investigate aneuploidy evolution in single cells from patients with triple-negative (ER-/PR-/HER2-) breast cancers ([Navin et al., 2011](#)). These data revealed that copy number aberrations evolved in punctuated bursts of evolution, followed by stable clonal expansions to form the tumor mass. In another study, single-cell exome sequencing (NUC-SEQ) showed that point mutations evolved gradually over time generating extensive clonal diversity and many rare (<1%) mutations in the tumor mass ([Wang et al., 2014](#)). Single-cell exome sequencing has also been applied to study clonal diversity in renal carcinoma ([Xu et al., 2012](#)) and a *JAK2*-positive myeloproliferative neoplasm ([Hou et al., 2012](#)), which identified a monoclonal population of cells that shared a common genetic lineage. Similarly, single cell exome sequencing was applied to study a muscle-invasive bladder cancer ([Li et al., 2012](#)), and a colon cancer patient ([Yu et al., 2014](#)), which identified two distinct subpopulations of cells in each of which tumor diverged, but also shared a common set of founder mutations. Another study used DNA SCS to delineate clonal diversity in glioblastoma, which revealed convergent evolution of EGFR mutations in different subclones from the same primary tumors ([Francis et al., 2014](#)).

DNA SCS has also been used to study clonal evolution in hematopoietic cancers. In one study, single cancer cells were sequenced from three patients diagnosed with MDS-derived secondary AML to reconstruct mutational chronology ([Hughes et al., 2014](#)). In another study 1,479 single cells were sequenced from six acute lymphoblastic leukemia (ALL) patients using targeted panels, which identified the presence of multiple clonal subpopulations in many AML patients. Clonal dynamics have also been investigated in xenografts, which showed extensive selection in the first transplantation

passages, followed by clonal dominance (Eirew et al., 2014). Collectively, these studies provide strong evidence for clonal evolution in many human tumors (Campbell and Polyak, 2007; Greaves and Maley, 2012; Navin and Hicks, 2010) by showing that single cells can continue to acquire new mutations and evolve to form the primary tumor mass.

Recent work has begun to use SCS to investigate metastatic dissemination and circulating tumor cells (CTCs) in the blood. In one study, RNA SCS was used to profile CTCs in the blood of melanoma patients (Ramskold et al., 2012). In another study, DNA SCS was used to analyze CTCs from six patients with metastatic colon cancer, showing that many of the driver mutations in the primary tumor could be detected in the CTCs (Heitzer et al., 2013). In another study, MALBAC was used to perform exome sequencing and copy number profiling of single CTCs from 7 metastatic lung adenocarcinoma cancer patients (Ni et al., 2013). In another study, single-cell exome sequencing was applied to a patient with metastatic prostate cancer, which identified 51% of the mutations in the primary and metastatic tumors in the CTC populations (Lohr et al., 2014). RNA SCS was also recently used to investigate CTC clusters in metastatic seeding in breast cancer (Aceto et al., 2014). Another study applied SNS and morphometric imaging to investigate copy number evolution in response to Abiraterone therapy in metastatic prostate cancer (Dago et al., 2014). Collectively, these studies have improved our understanding of CTCs and metastasis in human cancers.

RNA SCS has also been used to study cell plasticity and cancer stem cells in human tumors. An unbiased study of hundreds of single cell transcriptomes in five glioblastoma patients showed that cancer cells displayed a large range of intermediate phenotypes, that do not fall into distinct classes of epithelial or mesenchymal cell types. In summary, SCS methods have already had a large impact on improving our fundamental understanding of intratumor heterogeneity, clonal evolution and metastatic dissemination in human cancers. For a more detailed review on SCS applications in cancer research, please refer to the following review articles (Navin, 2014; Van Loo and Voet, 2014).

CLINICAL APPLICATIONS

SCS methods have direct translational applications in cancer treatment and pre-natal genetic diagnosis (PGD). In cancer research, intratumor heterogeneity presents a major challenge for clinical diagnostics, because single samples may not represent the tumor as a whole. While regional sequencing and deep-sequencing can resolve some clonal substructure, they cannot fully delineate the clonal substructure of a tumor and are inherently unable to determine which combination of mutations occur in each clone. SCS provides a powerful tool for resolving intratumor heterogeneity, and guiding targeted therapy towards the most malignant clones. SCS can also be used to calculate a diversity index for each cancer patient, which may have prognostic utility for predicting poor survival and poor response to chemotherapy. SCS technologies will also have direct applications for non-invasive monitoring, by sequencing single CTCs in the blood to track mutations in the primary and metastatic tumors. Several studies have already shown that over 50% of the mutations in the primary and metastatic tumors can be detected in CTCs of lung cancer (Ni et al., 2013), prostate cancer (Lohr et al., 2014) and colon cancer patients (Heitzer et al., 2013). By sequencing CTCs at multiple time-points over the course of therapy, oncologists can track mutational evolution and make rapid changes to their therapeutic strategies before resistance emerges. SCS methods will also have clinical applications in the early detection of tumor cells in bodily fluids (urine, sputum, blood) and fine-needle-aspirates samples.

Another major area of clinical utility is pre-implantation genetic diagnosis (PGD) and *in vitro* fertilization (IVF). During this procedure a biopsy of a single cell is collected from a set of blastomeres for DNA SCS to screen for genetic disorders prior to implantation into the uterus. In the past, these

methods have traditionally been limited to cytogenetic analysis and single-cell PCR. SCS provides the advantage of being able to profile thousands of mutations and copy number changes associated with diseases that can be screened from one cell using a single assay. In a proof-of-concept study, SCS was used to profile genomic copy number and structural variants in single cells from blastomeres derived from a human zygote after IVF (Voet et al., 2013). In another study, MALBAC was used to sequence polar bodies to identify copy number changes and point mutations prior to implantation (Hou et al., 2013). These preliminary PGD studies demonstrate the technical feasibility of screening oocytes and blastomeres to avoid the genetic transmission of diseases, paving the way for future clinical trials. For a comprehensive review on this topic please refer to the following article (Van der Aa et al., 2013).

COMPUTATIONAL METHODS

While SCS methods are generating torrents of large-scale genomic datasets, the computational methods for analyzing these data are severely lacking. SCS data is distinct from standard NGS data and their analysis tools, due to inherent technical errors and noise, including coverage non-uniformity, sparse data, false-positive errors, amplification biases and allelic dropout events. Some of the first SCS analysis methods focused on quantifying single cell copy number profiles from read count data. To calculate SCS copy number profiles, a variable-binning algorithm was developed that normalizes errors in mappability in the human genome, by adjusting genomic intervals based on the expected number of reads (Baslan et al., 2012; Navin et al., 2011). This processing pipeline was developed into a user-friendly web server platform with impressive visualization tools called Ginkgo (<http://gb.cshl.edu/ginkgo>). Another copy number method uses SCS read count data generated from DOP-PCR that corrects for GC bias and performs binary segmentation followed by dynamic thresholding (Zhang et al., 2013).

Several computational methods have also been developed for analyzing RNA SCS datasets to mitigate technical error. In one method RNA spike-in controls were used to quantify technical noise during WTA (Katayama et al., 2013). In another method unique molecular identifiers (UMI) were used to label RNA before WTA and sequencing, to eliminate amplification bias (Islam et al., 2014).

Computational methods have also been developed to model noise in RNA SCS data using a low-magnitude Poisson processes (Brennecke et al., 2013). Another RNA SCS method called Monocle represents each cell as a point in a high-dimension space, and uses dimensionality reduction to extract essential features over time (Trapnell et al., 2014). Another study developed a latent variable model for single cell RNA data to reduce technical noise from over amplification and cell cycle genes (Buettnner et al., 2015). Several algorithms have also been developed for assembly of microbial genomes from single cells. One method called E+V-SC uses lower initial coverage cutoff and then progressively increases the cutoff to incorporate more bases (Chitsaz et al., 2011). Another method called IDBA-UD uses similar filtering with progressive coverage thresholds strategy and error correction (Peng et al., 2012). A third method, SPAdes, tackled the uneven coverage problem by constructing paired assembly graphs utilizing read-pairs (Bankevich et al., 2012). In summary, while some initial tools have been developed, new quantitative methods are still urgently needed for analyzing DNA and RNA SCS datasets.

Alternatives to Single Cell Sequencing

SCS is not the appropriate technology to address every question in biology. In many studies alternative approaches will provide more powerful tools for investigating population diversity and identifying rare mutations. Methods such as deep-sequencing (Shah et al., 2012) or multi-region sequencing(Gerlinger et al., 2012) provide a more economical approach for resolving complex population substructure and

have the advantage of providing genotyping information on thousands of cells. In cases where living tissue or cells are available, single cells can be subcloned to generate isogenic cell lines or organoids that act as proxies for single cells ([Boj et al., 2015](#); [Sachs and Clevers, 2014](#)). These systems have the advantage of providing an unlimited amount of genetic material for analysis and can be used for functional assays. However, a notable limitation is that most cells are not capable of expanding in culture, which can introduce a strong bias in the representation of the final cells that are derived from a population. Furthermore as cells adapt to the cell culture environment they may alter transcriptional or epigenetic programs. In conclusion, alternative methods to SCS may be a better choice when functional studies are required, or when very rare cells must be detected in a population (without prior isolation or enrichment).

CONCLUSIONS & FUTURE DIRECTIONS

SCS methods have provided great insight into our understanding of biological diversity and rare cells that have previously been difficult to resolve in genomic data from bulk tissue samples. These tools have had a broad impact on many diverse fields of biology over the past 5 years, and several common applications have emerged: 1) delineating population diversity, 2) tracing cell lineages, 3) classifying cell types, and 4) genomic profiling of rare cells. While many initial studies have been published, there are still many applications that remain unexplored. In microbiology, SCS methods have yet to be applied to study viruses in single host cells, to understand how they infect and replicate differently in certain cell types. In neurobiology, SCS methods can provide important information on transcriptional programs in response to stimuli, including auditory, sensory and visual stimulation. In development, single-cell RNA sequencing can be used to study cell lineages in many organ systems to identify new markers and cell types. In tissue mosaicism, future studies should be directed at investigating the diversity of point mutations and indels in different tissue types which are likely to show even more diversity than copy number variations. Cancer immunotherapy is another exciting application, where SCS tools have great potential for illuminating phenomenon such as immunoediting and antigenicity in the context of intratumor heterogeneity. In cancer research, SCS can also help to understand the role of clonal diversity in complex biological processes, such as transformation, invasion and the evolution of chemoresistance ([Navin, 2014](#); [Van Loo and Voet, 2014](#)).

Future efforts in technology development should focus on *in situ* SCS methods that can measure genomic data on single cells while preserving their spatial context in tissues ([Crosetto et al., 2014](#)). Future technologies should also be directed at linking phenotypes and genotypes in single cells, by combining methods such as live-cell imaging with SCS methods. Forthcoming technologies should also focus on collecting combinations of genomic information from the same single cell in parallel (ex. DNA and RNA, or RNA and epigenomic modifications). Some progress was recently made in this area, by demonstrating the feasibility of measuring both copy number states and RNA expression profiles in the same single cells ([Dey et al., 2015](#)). Another important area of technology development is highly-multiplexed single cell DNA and RNA sequencing, to enable the profiling of thousands of single cells in parallel, at a substantially lower cost. A recent technique using microwells and DNA beads with barcodes shows promise for enabling the profiling 10,000 – 100,000 single cells in parallel ([Fan et al., 2015](#)). Several companies (Fluidigm, Wafergen, Cellular Research) are also focusing their efforts on developing higher-throughput single cell RNA and DNA sequencing methods, which are expected to come to market soon. While most SCS studies are still cost-prohibitive, we expect that this barrier will largely be dissolved over the next few years, as the costs of NGS technologies (Illumina, Life Technologies) continues to plummet through new technical innovations and fierce industrial competition.

In closing, while the SCS field is still relatively new, it has already made a large impact on many diverse fields of biology and has led to great improvements to our fundamental understanding of human diseases. We expect that the demand and application of SCS tools will continue to grow tremendously in the coming years, as these methods become more refined, high-throughput, inexpensive and easier to use in standard research and clinical laboratories.

Supplementary Material

01

[Click here to view.](#) (6.3K, pdf)

02

[Click here to view.](#) (58K, xlsx)

03

[Click here to view.](#) (1.5M, pdf)

Acknowledgements

N.N. is a Nadia's Gift Foundation Damon Runyon-Rachleff Innovator (DRR-25-13). This work is supported by grants to N.N. from NCI (1RO1CA169244-01). N.N. is a T.C. Hsu Endowed Scholar. Y.W. was supported by an Agilent University relations grant. This work was also supported by the Moonshot Knowledge Gap Award, the Center for Genetics & Genomics and the Center for Epigenetics at the MD Anderson Cancer Center. This work was also supported by a grant to N.N. from the Eric & Liz Lefkofsky Family Foundation.

Footnotes

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

References

1. Aceto N, Bardia A, Miyamoto DT, Donaldson MC, Wittner BS, Spencer JA, Yu M, Pely A, Engstrom A, Zhu H, et al. Circulating tumor cell clusters are oligoclonal precursors of breast cancer metastasis. *Cell.* 2014;158:1110–1122. [\[PMC free article\]](#) [\[PubMed\]](#) [\[Google Scholar\]](#)
2. Adams DL, Martin SS, Alpaugh RK, Charpentier M, Tsai S, Bergan RC, Ogden IM, Catalona W, Chumsri S, Tang CM, et al. Circulating giant macrophages as a potential biomarker of solid tumors. *Proc Natl Acad Sci U S A.* 2014;111:3514–3519. [\[PMC free article\]](#) [\[PubMed\]](#) [\[Google Scholar\]](#)

3. Altomare L, Borgatti M, Medoro G, Manaresi N, Tartagni M, Guerrieri R, Gambari R. Levitation and movement of human tumor cells using a printed circuit board device based on software-controlled dielectrophoresis. *Biotechnology and bioengineering*. 2003;82:474–479. [PubMed] [Google Scholar]
4. Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM, Nikolenko SI, Pham S, Prjibelski AD, et al. SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. *Journal of computational biology : a journal of computational molecular cell biology*. 2012;19:455–477. [PMC free article] [PubMed] [Google Scholar]
5. Baslan T, Kendall J, Rodgers L, Cox H, Riggs M, Stepansky A, Troge J, Ravi K, Esposito D, Lakshmi B, et al. Genome-wide copy number analysis of single cells. *Nat Protoc.* 2012;7:1024–1041. [PMC free article] [PubMed] [Google Scholar]
6. Bianconi E, Piovesan A, Facchini F, Beraudi A, Casadei R, Frabetti F, Vitale L, Pelleri MC, Tassani S, Piva F, et al. An estimation of the number of cells in the human body. *Annals of human biology*. 2013;40:463–471. [PubMed] [Google Scholar]
7. Biesecker LG, Spinner NB. A genomic view of mosaicism and human disease. *Nat Rev Genet.* 2013;14:307–320. [PubMed] [Google Scholar]
8. Blainey PC, Mosier AC, Potanina A, Francis CA, Quake SR. Genome of a low-salinity ammonia-oxidizing archaeon determined by single-cell and metagenomic analysis. *PLoS One.* 2011;6:e16626. [PMC free article] [PubMed] [Google Scholar]
9. Blainey PC, Quake SR. Dissecting genomic diversity, one cell at a time. *Nat Methods.* 2014;11:19–21. [PMC free article] [PubMed] [Google Scholar]
10. Boj SF, Hwang CI, Baker LA, Chio Ii, Engle DD, Corbo V, Jager M, Ponz-Sarvise M, Tiriac H, Spector MS, et al. Organoid models of human and mouse ductal pancreatic cancer. *Cell.* 2015;160:324–338. [PMC free article] [PubMed] [Google Scholar]
11. Brennecke P, Anders S, Kim JK, Kolodziejczyk AA, Zhang X, Proserpio V, Baying B, Benes V, Teichmann SA, Marioni JC, et al. Accounting for technical noise in single-cell RNA-seq experiments. *Nat Methods.* 2013;10:1093–1095. [PubMed] [Google Scholar]
12. Brunskill EW, Park JS, Chung E, Chen F, Magella B, Potter SS. Single cell dissection of early kidney development: multilineage priming. *Development.* 2014;141:3093–3101. [PMC free article] [PubMed] [Google Scholar]
13. Buettner F, Natarajan KN, Casale FP, Proserpio V, Scialdone A, Theis FJ, Teichmann SA, Marioni JC, Stegle O. Computational analysis of cell-to-cell heterogeneity in single-cell RNA-sequencing data reveals hidden subpopulations of cells. *Nat Biotechnol.* 2015;33:155–160. [PubMed] [Google Scholar]
14. Cai X, Evrony GD, Lehmann HS, Elhosary PC, Mehta BK, Poduri A, Walsh CA. Single-cell, genome-wide sequencing identifies clonal somatic copy-number variation in the human brain. *Cell reports.* 2014;8:1280–1289. [PMC free article] [PubMed] [Google Scholar]
15. Campbell LL, Polyak K. Breast tumor heterogeneity: cancer stem cells or clonal evolution? *Cell cycle.* 2007;6:2332–2338. [PubMed] [Google Scholar]
16. Chitsaz H, Yee-Greenbaum JL, Tesler G, Lombardo MJ, Dupont CL, Badger JH, Novotny M, Rusch DB, Fraser LJ, Gormley NA, et al. Efficient de novo assembly of single-cell bacterial genomes from short-read data sets. *Nat Biotechnol.* 2011;29:915–921. [PMC free article] [PubMed] [Google Scholar]
17. Choi JH, Ogunniyi AO, Du M, Du M, Kretschmann M, Eberhardt J, Love JC. Development and optimization of a process for automated recovery of single cells identified by microengraving. *Biotechnology progress.* 2010;26:888–895. [PubMed] [Google Scholar]

18. Cristofanilli M, Budd GT, Ellis MJ, Stopeck A, Matera J, Miller MC, Reuben JM, Doyle GV, Allard WJ, Terstappen LW, et al. Circulating tumor cells, disease progression, and survival in metastatic breast cancer. *N Engl J Med.* 2004;351:781–791. [[PubMed](#)] [[Google Scholar](#)]
19. Crosetto N, Bienko M, van Oudenaarden A. Spatially resolved transcriptomics and beyond. *Nat Rev Genet.* 2014 [[PubMed](#)] [[Google Scholar](#)]
20. Dago AE, Stepansky A, Carlsson A, Luttgen M, Kendall J, Baslan T, Kolatkar A, Wigler M, Bethel K, Gross ME, et al. Rapid phenotypic and genomic change in response to therapeutic pressure in prostate cancer inferred by high content analysis of single circulating tumor cells. *PLoS One.* 2014;9:e101777. [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
21. de Bourcy CF, De Vlaminck I, Kanbar JN, Wang J, Gawad C, Quake SR. A quantitative comparison of single-cell whole genome amplification methods. *PLoS One.* 2014;9:e105585. [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
22. De Carlos JA, Borrell J. A historical reflection of the contributions of Cajal and Golgi to the foundations of neuroscience. *Brain research reviews.* 2007;55:8–16. [[PubMed](#)] [[Google Scholar](#)]
23. Dean FB, Hosono S, Fang L, Wu X, Faruqi AF, Bray-Ward P, Sun Z, Zong Q, Du Y, Du J, et al. Comprehensive human genome amplification using multiple displacement amplification. *Proc Natl Acad Sci U S A.* 2002;99:5261–5266. [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
24. Dey SS, Kester L, Spanjaard B, Bienko M, van Oudenaarden A. Integrated genome and transcriptome sequencing of the same cell. *Nat Biotechnol.* 2015;33:285–289. [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
25. Eirew P, Steif A, Khattri J, Ha G, Yap D, Farahani H, Gelmon K, Chia S, Mar C, Wan A, et al. Dynamics of genomic clones in breast cancer patient xenografts at single-cell resolution. *Nature.* 2014 [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
26. Evrony GD, Cai X, Lee E, Hills LB, Elhosary PC, Lehmann HS, Parker JJ, Atabay KD, Gilmore EC, Poduri A, et al. Single-neuron sequencing analysis of L1 retrotransposition and somatic mutation in the human brain. *Cell.* 2012;151:483–496. [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
27. Fan HC, Fu GK, Fodor SP. Expression profiling. Combinatorial labeling of single cells for gene expression cytometry. *Science.* 2015;347:1258367. [[PubMed](#)] [[Google Scholar](#)]
28. Fiegler H, Geigl JB, Langer S, Rigler D, Porter K, Unger K, Carter NP, Speicher MR. High resolution array-CGH analysis of single cells. *Nucleic Acids Res.* 2007;35:e15. [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
29. Francis JM, Zhang CZ, Maire CL, Jung J, Manzo VE, Adalsteinsson VA, Homer H, Haidar S, Blumenstiel B, Pedamallu CS, et al. EGFR variant heterogeneity in glioblastoma resolved through single-nucleus sequencing. *Cancer discovery.* 2014;4:956–971. [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
30. Gerlinger M, Rowan AJ, Horswell S, Larkin J, Endesfelder D, Gronroos E, Martinez P, Matthews N, Stewart A, Tarpey P, et al. Intratumor heterogeneity and branched evolution revealed by multiregion sequencing. *N Engl J Med.* 2012;366:883–892. [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
31. Gole J, Gore A, Richards A, Chiu YJ, Fung HL, Bushman D, Chiang HI, Chun J, Lo YH, Zhang K. Massively parallel polymerase cloning and genome sequencing of single cells using nanoliter microwells. *Nat Biotechnol.* 2013;31:1126–1132. [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
32. Greaves M, Maley CC. Clonal evolution in cancer. *Nature.* 2012;481:306–313. [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]

33. Guo H, Zhu P, Wu X, Li X, Wen L, Tang F. Single-cell methylome landscapes of mouse embryonic stem cells and early embryos analyzed using reduced representation bisulfite sequencing. *Genome Res.* 2013;23:2126–2135. [PMC free article] [PubMed] [Google Scholar]
34. Heitzer E, Auer M, Gasch C, Pichler M, Ulz P, Hoffmann EM, Lax S, Waldspuehl-Geigl J, Mauermann O, Lackner C, et al. Complex tumor genomes inferred from single circulating tumor cells by array-CGH and next-generation sequencing. *Cancer Res.* 2013;73:2965–2975. [PubMed] [Google Scholar]
35. Hou Y, Fan W, Yan L, Li R, Lian Y, Huang J, Li J, Xu L, Tang F, Xie XS, et al. Genome analyses of single human oocytes. *Cell.* 2013;155:1492–1506. [PubMed] [Google Scholar]
36. Hou Y, Song L, Zhu P, Zhang B, Tao Y, Xu X, Li F, Wu K, Liang J, Shao D, et al. Single-cell exome sequencing and monoclonal evolution of a JAK2-negative myeloproliferative neoplasm. *Cell.* 2012;148:873–885. [PubMed] [Google Scholar]
37. Hughes AE, Magrini V, Demeter R, Miller CA, Fulton R, Fulton LL, Eades WC, Elliott K, Heath S, Westervelt P, et al. Clonal architecture of secondary acute myeloid leukemia defined by single-cell sequencing. *PLoS genetics.* 2014;10:e1004462. [PMC free article] [PubMed] [Google Scholar]
38. Ishoey T, Woyke T, Stepanauskas R, Novotny M, Lasken RS. Genomic sequencing of single microbial cells from environmental samples. *Current opinion in microbiology.* 2008;11:198–204. [PMC free article] [PubMed] [Google Scholar]
39. Islam S, Zeisel A, Joost S, La Manno G, Zajac P, Kasper M, Lonnerberg P, Linnarsson S. Quantitative single-cell RNA-seq with unique molecular identifiers. *Nat Methods.* 2014;11:163–166. [PubMed] [Google Scholar]
40. Jaitin DA, Kenigsberg E, Keren-Shaul H, Elefant N, Paul F, Zaretsky I, Mildner A, Cohen N, Jung S, Tanay A, et al. Massively parallel single-cell RNA-seq for markerfree decomposition of tissues into cell types. *Science.* 2014;343:776–779. [PMC free article] [PubMed] [Google Scholar]
41. Katayama S, Tohonen V, Linnarsson S, Kere J. SAMstrt: statistical test for differential expression in single-cell transcriptome with spike-in normalization. *Bioinformatics.* 2013;29:2943–2945. [PMC free article] [PubMed] [Google Scholar]
42. Kirkness EF, Grindberg RV, Yee-Greenbaum J, Marshall CR, Scherer SW, Lasken RS, Venter JC. Sequencing of isolated sperm cells for direct haplotyping of a human genome. *Genome Res.* 2013;23:826–832. [PMC free article] [PubMed] [Google Scholar]
43. Knouse KA, Wu J, Whittaker CA, Amon A. Single cell sequencing reveals low levels of aneuploidy across mammalian tissues. *Proc Natl Acad Sci U S A.* 2014;111:13409–13414. [PMC free article] [PubMed] [Google Scholar]
44. Lasken RS. Single-cell genomic sequencing using Multiple Displacement Amplification. *Current opinion in microbiology.* 2007;10:510–516. [PubMed] [Google Scholar]
45. Lasken RS. Single-cell sequencing in its prime. *Nat Biotechnol.* 2013;31:211–212. [PubMed] [Google Scholar]
46. Lasken RS, McLean JS. Recent advances in genomic DNA sequencing of microbial species from single cells. *Nat Rev Genet.* 2014;15:577–584. [PMC free article] [PubMed] [Google Scholar]
47. Li Y, Xu X, Song L, Hou Y, Li Z, Tsang S, Li F, Im KM, Wu K, Wu H, et al. Single-cell sequencing analysis characterizes common and cell-lineage-specific mutations in a muscle-invasive bladder cancer. *GigaScience.* 2012;1:12. [PMC free article] [PubMed] [Google Scholar]
48. Lohr JG, Adalsteinsson VA, Cibulskis K, Choudhury AD, Rosenberg M, Cruz-Gordillo P, Francis JM, Zhang CZ, Shalek AK, Satija R, et al. Whole-exome sequencing of circulating tumor cells provides a window into metastatic prostate cancer. *Nat Biotechnol.* 2014;32:479–

484. [PMC free article] [PubMed] [Google Scholar]
49. Lovatt D, Ruble BK, Lee J, Dueck H, Kim TK, Fisher S, Francis C, Spaethling JM, Wolf JA, Grady MS, et al. Transcriptome in vivo analysis (TIVA) of spatially defined single cells in live tissue. *Nat Methods*. 2014;11:190–196. [PMC free article] [PubMed] [Google Scholar]
50. Lu S, Zong C, Fan W, Yang M, Li J, Chapman AR, Zhu P, Hu X, Xu L, Yan L, et al. Probing meiotic recombination and aneuploidy of single sperm cells by whole-genome sequencing. *Science*. 2012;338:1627–1630. [PMC free article] [PubMed] [Google Scholar]
51. Macaulay IC, Voet T. Single cell genomics: advances and future perspectives. *PLoS genetics*. 2014;10:e1004126. [PMC free article] [PubMed] [Google Scholar]
52. Mardis ER. A decade's perspective on DNA sequencing technology. *Nature*. 2011;470:198–203. [PubMed] [Google Scholar]
53. McConnell MJ, Lindberg MR, Brennand KJ, Piper JC, Voet T, Cowing-Zitron C, Shumilina S, Lasken RS, Vermeesch JR, Hall IM, et al. Mosaic copy number variation in human neurons. *Science*. 2013;342:632–637. [PMC free article] [PubMed] [Google Scholar]
54. Nagano T, Lubling Y, Stevens TJ, Schoenfelder S, Yaffe E, Dean W, Laue ED, Tanay A, Fraser P. Single-cell Hi-C reveals cell-to-cell variability in chromosome structure. *Nature*. 2013;502:59–64. [PMC free article] [PubMed] [Google Scholar]
55. Nagrath S, Sequist LV, Maheswaran S, Bell DW, Irimia D, Ulkus L, Smith MR, Kwak EL, Digumarthy S, Muzikansky A, et al. Isolation of rare circulating tumour cells in cancer patients by microchip technology. *Nature*. 2007;450:1235–1239. [PMC free article] [PubMed] [Google Scholar]
56. Navin N, Hicks J. Future medical applications of single-cell sequencing in cancer. *Genome Med*. 2011;3:31. [PMC free article] [PubMed] [Google Scholar]
57. Navin N, Kendall J, Troge J, Andrews P, Rodgers L, McIndoo J, Cook K, Stepansky A, Levy D, Esposito D, et al. Tumour evolution inferred by single-cell sequencing. *Nature*. 2011;472:90–94. [PMC free article] [PubMed] [Google Scholar]
58. Navin NE. Cancer genomics: one cell at a time. *Genome Biol*. 2014;15:452. [PMC free article] [PubMed] [Google Scholar]
59. Navin NE, Hicks J. Tracing the tumor lineage. *Mol Oncol*. 2010;4:267–283. [PMC free article] [PubMed] [Google Scholar]
60. Ni X, Zhuo M, Su Z, Duan J, Gao Y, Wang Z, Zong C, Bai H, Chapman AR, Zhao J, et al. Reproducible copy number variation patterns among single circulating tumor cells of lung cancer patients. *Proc Natl Acad Sci U S A*. 2013;110:21083–21088. [PMC free article] [PubMed] [Google Scholar]
61. Pamp SJ, Harrington ED, Quake SR, Relman DA, Blainey PC. Single-cell sequencing provides clues about the host interactions of segmented filamentous bacteria (SFB). *Genome Res*. 2012;22:1107–1119. [PMC free article] [PubMed] [Google Scholar]
62. Peng Y, Leung HC, Yiu SM, Chin FY. IDBA-UD: a de novo assembler for single-cell and metagenomic sequencing data with highly uneven depth. *Bioinformatics*. 2012;28:1420–1428. [PubMed] [Google Scholar]
63. Pollen AA, Nowakowski TJ, Shuga J, Wang X, Leyrat AA, Lui JH, Li N, Szpankowski L, Fowler B, Chen P, et al. Low-coverage single-cell mRNA sequencing reveals cellular heterogeneity and activated signaling pathways in developing cerebral cortex. *Nat Biotechnol*. 2014;32:1053–1058. [PMC free article] [PubMed] [Google Scholar]
64. Powell AA, Talasaz AH, Zhang H, Coram MA, Reddy A, Deng G, Telli ML, Advani RH, Carlson RW, Mollick JA, et al. Single cell profiling of circulating tumor cells: transcriptional heterogeneity and diversity from breast cancer cell lines. *PLoS One*. 2012;7:e33788.

[PMC free article] [PubMed] [Google Scholar]

65. Qiu S, Luo S, Evgrafov O, Li R, Schroth GP, Levitt P, Knowles JA, Wang K. Single-neuron RNA-Seq: technical feasibility and reproducibility. *Frontiers in genetics*. 2012;3:124. [PMC free article] [PubMed] [Google Scholar]
66. Ramskold D, Luo S, Wang YC, Li R, Deng Q, Faridani OR, Daniels GA, Khrebtukova I, Loring JF, Laurent LC, et al. Full-length mRNA-Seq from single-cell levels of RNA and individual circulating tumor cells. *Nat Biotechnol*. 2012;30:777–782. [PMC free article] [PubMed] [Google Scholar]
67. Rinke C, Schwientek P, Sczyrba A, Ivanova NN, Anderson IJ, Cheng JF, Darling A, Malfatti S, Swan BK, Gies EA, et al. Insights into the phylogeny and coding potential of microbial dark matter. *Nature*. 2013;499:431–437. [PubMed] [Google Scholar]
68. Rodrigue S, Malmstrom RR, Berlin AM, Birren BW, Henn MR, Chisholm SW. Whole genome amplification and de novo assembly of single bacterial cells. *PLoS One*. 2009;4:e6864. [PMC free article] [PubMed] [Google Scholar]
69. Sachs N, Clevers H. Organoid cultures for the analysis of cancer phenotypes. *Curr Opin Genet Dev*. 2014;24:68–73. [PubMed] [Google Scholar]
70. Saliba AE, Westermann AJ, Gorski SA, Vogel J. Single-cell RNA-seq: advances and future challenges. *Nucleic Acids Res*. 2014;42:8845–8860. [PMC free article] [PubMed] [Google Scholar]
71. Sandberg R. Entering the era of single-cell transcriptomics in biology and medicine. *Nat Methods*. 2014;11:22–24. [PubMed] [Google Scholar]
72. Shah SP, Roth A, Goya R, Oloumi A, Ha G, Zhao Y, Turashvili G, Ding J, Tse K, Haffari G, et al. The clonal and mutational evolution spectrum of primary triple-negative breast cancers. *Nature*. 2012;486:395–399. [PMC free article] [PubMed] [Google Scholar]
73. Shalek AK, Satija R, Adiconis X, Gertner RS, Gaublomme JT, Raychowdhury R, Schwartz S, Yosef N, Malboeuf C, Lu D, et al. Single-cell transcriptomics reveals bimodality in expression and splicing in immune cells. *Nature*. 2013;498:236–240. [PMC free article] [PubMed] [Google Scholar]
74. Shalek AK, Satija R, Shuga J, Trombetta JJ, Gennert D, Lu D, Chen P, Gertner RS, Gaublomme JT, Yosef N, et al. Single-cell RNA-seq reveals dynamic paracrine control of cellular variation. *Nature*. 2014;510:363–369. [PMC free article] [PubMed] [Google Scholar]
75. Shapiro E, Biezuner T, Linnarsson S. Single-cell sequencing-based technologies will revolutionize whole-organism science. *Nat Rev Genet*. 2013;14:618–630. [PubMed] [Google Scholar]
76. Talseth-Palmer BA, Bowden NA, Hill A, Meldrum C, Scott RJ. Whole genome amplification and its impact on CGH array profiles. *BMC Res Notes*. 2008;1:56. [PMC free article] [PubMed] [Google Scholar]
77. Tang F, Barbacioru C, Bao S, Lee C, Nordman E, Wang X, Lao K, Surani MA. Tracing the derivation of embryonic stem cells from the inner cell mass by single-cell RNA-Seq analysis. *Cell Stem Cell*. 2010;6:468–478. [PMC free article] [PubMed] [Google Scholar]
78. Tang F, Barbacioru C, Wang Y, Nordman E, Lee C, Xu N, Wang X, Bodeau J, Tuch BB, Siddiqui A, et al. mRNA-Seq whole-transcriptome analysis of a single cell. *Nat Methods*. 2009;6:377–382. [PubMed] [Google Scholar]
79. Telenius H, Carter NP, Bebb CE, Nordenskjold M, Ponder BA, Tunnacliffe A. Degenerate oligonucleotide-primed PCR: general amplification of target DNA by a single degenerate primer. *Genomics*. 1992;13:718–725. [PubMed] [Google Scholar]

80. Trapnell C, Cacchiarelli D, Grimsby J, Pokharel P, Li S, Morse M, Lennon NJ, Livak KJ, Mikkelsen TS, Rinn JL. The dynamics and regulators of cell fate decisions are revealed by pseudotemporal ordering of single cells. *Nat Biotechnol.* 2014;32:381–386. [PMC free article] [PubMed] [Google Scholar]
81. Treutlein B, Brownfield DG, Wu AR, Neff NF, Mantalas GL, Espinoza FH, Desai TJ, Krasnow MA, Quake SR. Reconstructing lineage hierarchies of the distal lung epithelium using single-cell RNA-seq. *Nature.* 2014;509:371–375. [PMC free article] [PubMed] [Google Scholar]
82. Usoskin D, Furlan A, Islam S, Abdo H, Lonnerberg P, Lou D, Hjerling-Leffler J, Haeggstrom J, Kharchenko O, Kharchenko PV, et al. Unbiased classification of sensory neuron types by large-scale single-cell RNA sequencing. *Nature neuroscience.* 2014 [PubMed] [Google Scholar]
83. Van der Aa N, Zamani Esteki M, Vermeesch JR, Voet T. Preimplantation genetic diagnosis guided by single-cell genomics. *Genome Med.* 2013;5:71. [PMC free article] [PubMed] [Google Scholar]
84. Van Gelder RN, von Zastrow ME, Yool A, Dement WC, Barchas JD, Eberwine JH. Amplified RNA synthesized from limited quantities of heterogeneous cDNA. *Proc Natl Acad Sci U S A.* 1990;87:1663–1667. [PMC free article] [PubMed] [Google Scholar]
85. Van Loo P, Voet T. Single cell analysis of cancer genomes. *Curr Opin Genet Dev.* 2014;24:82–91. [PubMed] [Google Scholar]
86. Voet T, Kumar P, Van Loo P, Cooke SL, Marshall J, Lin ML, Zamani Esteki M, Van der Aa N, Mateiu L, McBride DJ, et al. Single-cell paired-end genome sequencing reveals structural variation per cell cycle. *Nucleic Acids Res.* 2013;41:6119–6138. [PMC free article] [PubMed] [Google Scholar]
87. Wang J, Fan HC, Behr B, Quake SR. Genome-wide single-cell analysis of recombination activity and de novo mutation rates in human sperm. *Cell.* 2012;150:402–412. [PMC free article] [PubMed] [Google Scholar]
88. Wang Y, Waters J, Leung ML, Unruh A, Roh W, Shi X, Chen K, Scheet P, Vattathil S, Liang H, et al. Clonal evolution in breast cancer revealed by single nucleus genome sequencing. *Nature.* 2014;512:155–160. [PMC free article] [PubMed] [Google Scholar]
89. Woyke T, Xie G, Copeland A, Gonzalez JM, Han C, Kiss H, Saw JH, Senin P, Yang C, Chatterji S, et al. Assembling the marine metagenome, one cell at a time. *PLoS One.* 2009;4:e5299. [PMC free article] [PubMed] [Google Scholar]
90. Xu X, Hou Y, Yin X, Bao L, Tang A, Song L, Li F, Tsang S, Wu K, Wu H, et al. Single-cell exome sequencing reveals single-nucleotide mutation characteristics of a kidney tumor. *Cell.* 2012;148:886–895. [PubMed] [Google Scholar]
91. Xue Z, Huang K, Cai C, Cai L, Jiang CY, Feng Y, Liu Z, Zeng Q, Cheng L, Sun YE, et al. Genetic programs in human and mouse early embryos revealed by single-cell RNA sequencing. *Nature.* 2013;500:593–597. [PMC free article] [PubMed] [Google Scholar]
92. Yan L, Yang M, Guo H, Yang L, Wu J, Li R, Liu P, Lian Y, Zheng X, Yan J, et al. Single-cell RNA-Seq profiling of human preimplantation embryos and embryonic stem cells. *Nature structural & molecular biology.* 2013;20:1131–1139. [PubMed] [Google Scholar]
93. Yong Wang JW, Leung Marco L, Unruh Anna, Roh Whijae, Shi Xiuqing, Chen Ken, Scheet Paul, Vattathil Selina, Liang Han, Multani Asha, Zhang Hong, Zhao Rui, Michor Franziska, Meric-Bernstam Funda, Navin Nicholas E. Clonal Evolution in Breast Cancer Revealed by Single Nucleus Genome Sequencing. *Nature.* 2014 [PMC free article] [PubMed] [Google Scholar]

94. Yu C, Yu J, Yao X, Wu WK, Lu Y, Tang S, Li X, Bao L, Li X, Hou Y, et al. Discovery of biclonal origin and a novel oncogene SLC12A5 in colon cancer by single-cell sequencing. *Cell research*. 2014;24:701–712. [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
95. Yu M, Stott S, Toner M, Maheswaran S, Haber DA. Circulating tumor cells: approaches to isolation and characterization. *The Journal of cell biology*. 2011;192:373–382. [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
96. Zhang C, Zhang C, Chen S, Yin X, Pan X, Lin G, Tan Y, Tan K, Xu Z, Hu P, et al. A single cell level based method for copy number variation analysis by low coverage massively parallel sequencing. *PLoS One*. 2013;8:e54236. [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
97. Zong C, Lu S, Chapman AR, Xie XS. Genome-wide detection of singlenucleotide and copy-number variations of a single human cell. *Science*. 2012;338:1622–1626. [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]