# Session IV **Sources of Heterogeneity**

Thomas J. Leeper

Government Department London School of Economics and Political Science

- 1 Other Survey Experimental Designs
- 2 Attention and Satisficing
- 3 Moderators and Effect Heterogeneity
  - Manipulate the Moderator
  - Blocking/Block Randomization
  - Post-hoc Approaches

- 1 Other Survey Experimental Designs
- 2 Attention and Satisficing
- 3 Moderators and Effect Heterogeneity
  - Manipulate the Moderator
  - Blocking/Block Randomization
  - Post-hoc Approaches

### **Beyond One-shot Designs**

- Surveys can be used as a measurement instrument for a field treatment or a manipulation applied in a different survey panel wave
  - 1 Measure effect duration in two-wave panel
  - 2 Solicit pre-treatment outcome measures in a two-wave panel
  - 3 Measure effects of field treatment in post-test only design
  - 4 Randomly encourage field treatment in pre-test and measure effects in post-test

### **Beyond One-shot Designs**

- Surveys can be used as a measurement instrument for a field treatment or a manipulation applied in a different survey panel wave
  - 1 Measure effect duration in two-wave panel
  - Solicit pre-treatment outcome measures in a two-wave panel
  - 3 Measure effects of field treatment in post-test only design
  - 4 Randomly encourage field treatment in pre-test and measure effects in post-test
- Problems? Compliance & nonresponse

#### I. Effect Duration

- Use a two- (or more-) wave panel to measure duration of effects
  - T1: Treatment and outcome measurement
  - T2+: Outcome measurement
- Two main concerns
  - Attrition
  - Panel conditioning

### II. Within-Subjects Designs

- Estimate treatment effects as a difference-in-differences
- Instead of using the post-treatment mean-difference in *Y* to estimate the causal effect, use the difference in pre-post differences for the two groups:


$$(\hat{Y}_{0,t+1} - \hat{Y}_{0,t}) - (\hat{Y}_{j,t+1} - \hat{Y}_{j,t})$$


### II. Within-Subjects Designs


- Estimate treatment effects as a difference-in-differences
- Instead of using the post-treatment mean-difference in *Y* to estimate the causal effect, use the difference in pre-post differences for the two groups:


$$(\hat{Y}_{0,t+1} - \hat{Y}_{0,t}) - (\hat{Y}_{j,t+1} - \hat{Y}_{j,t})$$


Advantageous because variance for paired samples decreases as correlation between  $t_0$  and  $t_1$  observations increases

















<sup>&</sup>lt;sup>1</sup>Shadish, Cook, and Campbell (2002)

As soon as time comes into play, we have to worry about threats to validity.<sup>1</sup>

History (simultaneous cause)

<sup>&</sup>lt;sup>1</sup>Shadish, Cook, and Campbell (2002)

- History (simultaneous cause)
- 2 Maturation (time trends)

<sup>&</sup>lt;sup>1</sup>Shadish, Cook, and Campbell (2002)

- History (simultaneous cause)
- 2 Maturation (time trends)
- Testing (observation changes respondents)

<sup>&</sup>lt;sup>1</sup>Shadish, Cook, and Campbell (2002)

- History (simultaneous cause)
- 2 Maturation (time trends)
- 3 Testing (observation changes respondents)
- 4 Instrumentation (changing operationalization)

<sup>&</sup>lt;sup>1</sup>Shadish, Cook, and Campbell (2002)

- History (simultaneous cause)
- 2 Maturation (time trends)
- 3 Testing (observation changes respondents)
- Instrumentation (changing operationalization)
- Instability (measurement error)

<sup>&</sup>lt;sup>1</sup>Shadish, Cook, and Campbell (2002)

- History (simultaneous cause)
- 2 Maturation (time trends)
- Testing (observation changes respondents)
- 4 Instrumentation (changing operationalization)
- 5 Instability (measurement error)
- 6 Attrition

<sup>&</sup>lt;sup>1</sup>Shadish, Cook, and Campbell (2002)

Examples:

- Examples:
  - Citizens randomly sent a letter by post encouraging them to reduce water usage

- Examples:
  - Citizens randomly sent a letter by post encouraging them to reduce water usage
  - 2 Different local media markets randomly assigned to receive different advertising

- Examples:
  - Citizens randomly sent a letter by post encouraging them to reduce water usage
  - 2 Different local media markets randomly assigned to receive different advertising
- Survey is used to measure outcomes, when treatment assignment is already known

- Examples:
  - Citizens randomly sent a letter by post encouraging them to reduce water usage
  - 2 Different local media markets randomly assigned to receive different advertising
- Survey is used to measure outcomes, when treatment assignment is already known
- Issues

- Examples:
  - Citizens randomly sent a letter by post encouraging them to reduce water usage
  - 2 Different local media markets randomly assigned to receive different advertising
- Survey is used to measure outcomes, when treatment assignment is already known
- Issues
  - Nonresponse
  - Noncompliance

### IV. Treatment Encouragement

- Design:
  - T1: Encourage treatment
  - T2: Measure effects
- Examples:
  - 1 Albertson and Lawrence

### IV. Treatment Encouragement

- Design:
  - T1: Encourage treatment
  - T2: Measure effects
- Examples:
  - 1 Albertson and Lawrence
- Issues

### IV. Treatment Encouragement

- Design:
  - T1: Encourage treatment
  - T2: Measure effects
- Examples:
  - 1 Albertson and Lawrence
- Issues
  - Nonresponse
  - Noncompliance

### **Treatment Noncompliance**

#### Definition:

"when subjects who were assigned to receive the treatment go untreated or when subjects assigned to the control group are treated" <sup>2</sup>

<sup>&</sup>lt;sup>2</sup>Gerber & Green. 2012. Field Experiments, p.132.

### **Treatment Noncompliance**

#### Definition:

"when subjects who were assigned to receive the treatment go untreated or when subjects assigned to the control group are treated" <sup>2</sup>

### Several strategies

- "As treated" analysis
- "Intention to treat" analysis
- Estimate a LATE

<sup>&</sup>lt;sup>2</sup>Gerber & Green. 2012. Field Experiments, p.132.

### **Analyzing Noncompliance**

- If noncompliance only occurs in one group, it is *asymmetric* or *one-sided*
- We can ignore non-compliance and analyze the "intention to treat" effect, which will underestimate our effects because some people were not treated as assigned:  $ITT = \overline{Y}_1 \overline{Y}_0$

### **Analyzing Noncompliance**

- If noncompliance only occurs in one group, it is *asymmetric* or *one-sided*
- We can ignore non-compliance and analyze the "intention to treat" effect, which will underestimate our effects because some people were not treated as assigned:  $ITT = \overline{Y}_1 \overline{Y}_0$

## **Local Average Treatment Effect**

- IV estimate is *local* to the variation in *X* that is due to variation in *D*
- This matters if effects are *heterogeneous*
- LATE is effect for those who *comply*
- Four subpopulations:
  - Compliers: X = 1 only if D = 1
  - Always-takers: X = 1 regardless of D
  - Never-takers: X = 0 regardless of D
  - Defiers: X = 1 only if D = 0
- Exclusion restriction! Monotonicity!

Moderators

# **Questions?**

- 1 Other Survey Experimental Designs
- 2 Attention and Satisficing
- 3 Moderators and Effect Heterogeneity
  - Manipulate the Moderator
  - Blocking/Block Randomization
  - Post-hoc Approaches

How should we deal with respondents that appear to not be paying attention, not "taking" the treatment, or not responding to outcome measures?

- Keep them
- 2 Throw them away

# Best Practice: Protocol

- Excluding respondents based on survey behavior is one of the easiest ways to "p-hack" an experimental dataset
  - Inattention, satisficing, etc. will tend to reduce the size of the SATE
- So regardless of how you handle these respondents, these should be decisions that are made *pre-analysis*

Pre-Treatment

#### Pre-Treatment

Satisficing behaviors

#### Pre-Treatment

- Satisficing behaviors
- Inattention

#### Pre-Treatment

- Satisficing behaviors
- Inattention
- Covariate-based selection

#### Pre-Treatment

- Satisficing behaviors
- Inattention
- Covariate-based selection
- Pretreated

### Pre-Treatment

- Satisficing behaviors
- Inattention
- Covariate-based selection
- Pretreated

#### Post-Treatment

Speeding on treatment

### Pre-Treatment

- Satisficing behaviors
- Inattention
- Covariate-based selection
- Pretreated

- Speeding on treatment
- "Failing" a manipulation check

### Pre-Treatment

- Satisficing behaviors
- Inattention
- Covariate-based selection
- Pretreated

- Speeding on treatment
- "Failing" a manipulation check
- Drop-off

# **Pre-Treatment Exclusion**

This is totally fine from a causal inference perspective


# **Pre-Treatment Exclusion**


- This is totally fine from a causal inference perspective
- Advantages:
  - Focused on engaged respondents
  - Likely increase impact of treatment

# Pre-Treatment Exclusion


- This is totally fine from a causal inference perspective
- Advantages:
  - Focused on engaged respondents
  - Likely increase impact of treatment
- Disadvantages:
  - Changing definition of sample (and thus population)

This is much more problematic because it involves controlling for a post-treatment variable





Risk that estimate of  $\beta_1$  is diminished because effect is being carried through the manipulation check.




Introduction of "collider bias" wherein values of the manipulation check are affected by other factors.


Any post-treatment exclusion is problematic and should be avoided

- Any post-treatment exclusion is problematic and should be avoided
- Can estimate a LATF
  - Interpretation: Effect of manipulation check among those whose value of the check can be changed by the treatment manipulation

- Any post-treatment exclusion is problematic and should be avoided
- Can estimate a LATF
  - Interpretation: Effect of manipulation check among those whose value of the check can be changed by the treatment manipulation
- Non-response or attrition is the same as researcher-imposed exclusion
  - Not problematic if MCAR
  - Nothing really to be done if caused by treatment



Introduction of "collider bias" wherein values of the manipulation check are affected by other factors.



- Any post-treatment exclusion is problematic and should be avoided
- Can estimate a LATF
  - Interpretation: Effect of manipulation check among those whose value of the check can be changed by the treatment manipulation

- Any post-treatment exclusion is problematic and should be avoided
- Can estimate a LATF
  - Interpretation: Effect of manipulation check among those whose value of the check can be changed by the treatment manipulation
- Non-response or attrition is the same as researcher-imposed exclusion
  - Not problematic if MCAR
  - Nothing really to be done if caused by treatment

# **Questions?**


- 1 Other Survey Experimental Designs
- 2 Attention and Satisficing
- 3 Moderators and Effect Heterogeneity
  - Manipulate the Moderator
  - Blocking/Block Randomization
  - Post-hoc Approaches


If we think there might be effect heterogeneity, what can we do?

- Best solution: manipulate the moderator
- Next best: block on the moderator
- Least best: post-hoc exploratory approaches

Simply: Manipulating the moderator variable is the best way to estimate a heterogeneous effect

Why is this true?





- How close do you feel to your ethnic or racial group?
- Some people have said that taxes need to be raised to take care of pressing national needs. How willing would you be to have your taxes raised to improve education in public schools?

 $<sup>^3</sup>$ Transue. 2007. "Identity Salience, Identity Acceptance, and Racial Policy Attitudes: American National Identity as a Uniting Force." *American Journal of Political Science* 51(1): 78–91.

- How close do you feel to other Americans?
- Some people have said that taxes need to be raised to take care of pressing national needs. How willing would you be to have your taxes raised to improve education in public schools?

 $<sup>^3</sup>$ Transue. 2007. "Identity Salience, Identity Acceptance, and Racial Policy Attitudes: American National Identity as a Uniting Force." *American Journal of Political Science* 51(1): 78–91.

- How close do you feel to your ethnic or racial group?
- Some people have said that taxes need to be raised to take care of pressing national needs. How willing would you be to have your taxes raised to improve educational opportunities for minorities?

 $<sup>^3</sup>$ Transue. 2007. "Identity Salience, Identity Acceptance, and Racial Policy Attitudes: American National Identity as a Uniting Force." *American Journal of Political Science* 51(1): 78–91.

- How close do you feel to other Americans?
- Some people have said that taxes need to be raised to take care of pressing national needs. How willing would you be to have your taxes raised to improve educational opportunities for minorities?

 $<sup>^3</sup>$ Transue. 2007. "Identity Salience, Identity Acceptance, and Racial Policy Attitudes: American National Identity as a Uniting Force." *American Journal of Political Science* 51(1): 78–91.

### 2x2 Factorial Design

#### Condition

Educ. for Minorities  $Y_1$ Schools

## 2x2 Factorial Design

Condition	Americans	Own Race	
Educ. for Minorities	$Y_{1,0}$	$Y_{1,1}$	
Schools	$Y_{0,0}$	$Y_{0,1}$	

# Two ways to estimate this

#### Dummy variable regression:

$$Y = \beta_0 + \beta_1 X_{0,1} + \beta_2 X_{1,0} + \beta_3 X_{1,1} + \epsilon$$

Interaction effect:

$$Y = \beta_0 + \beta_1 X 1_1 + \beta_2 X 2_1 + \beta_3 X 1_1 * X 2_1 + \epsilon$$

### **Considerations**

- Need to have hypotheses about heterogeneity a priori
- Factorial designs can quickly become unwieldy and expensive

### Probably obvious, but...

Factors	rs Conditions per factor Total Conditions		n
1	2	2	400
1	3	3	600
1	4	4	800
2	2	4	800
2	3	6	1200
2	4	8	1600
3	3	9	1800
3	4	12	2400
4	4	16	3200

Assumes power to detect a relatively small effect, but no consideration of multiple comparisons.

### **Considerations**

- Need to have hypotheses about heterogeneity a priori
- Factorial designs can quickly become unwieldy and expensive

### Considerations

- Need to have hypotheses about heterogeneity a priori
- Factorial designs can quickly become unwieldy and expensive
- Need to consider what CATEs are of theoretical interest
  - Treatment-control
  - Treatment-treatment

### **Questions?**

## Conjoint Designs I

- "Classic vignettes" taken to an extreme
- Address heterogeneity w/r/t SUTO
- Example: Judge whether to admit an immigrant to your country

"Classic vignettes" taken to an extreme

Moderators

- Address heterogeneity w/r/t SUTO
- Example: Judge whether to admit an immigrant to your country
- Respondents see a series of vignettes that are fully randomized along any number of dimensions
  - Sex, Education, Language proficiency, etc.

# Conjoint Designs I

- "Classic vignettes" taken to an extreme
  - Address heterogeneity w/r/t SUTO
- Example: Judge whether to admit an immigrant to your country
- Respondents see a series of vignettes that are fully randomized along any number of dimensions
  - Sex, Education, Language proficiency, etc.
- Outcome is judgment (binary or rating scale)

# Conjoint Designs II

#### Why is this useful?

- Understand complex decision-making
- Within-subjects comparisons
- Heterogeneous effects across versions of treatment
- Pilot testing: Sensitivity of design to specification of compound vignette

Please read the descriptions of the potential immigrants carefully. Then, please indicate which of the two immigrants you would personally prefer to see admitted to the United States.


	Immigrant 1	Immigrant 2
Prior Trips to the U.S.	Entered the U.S. once before on a tourist visa	Entered the U.S. once before on a tourist visa
Reason for Application	Reunite with family members already in U.S.	Reunite with family members already in U.S.
Country of Origin	Mexico	Iraq
Language Skills	During admission interview, this applicant spoke fluent English	During admission interview, this applicant spoke fluent English
Profession	Child care provider	Teacher
Job Experience	One to two years of job training and experience	Three to five years of job training and experience
Employment Plans	Does not have a contract with a U.S. employer but has done job interviews	Will look for work after arriving in the U.S.
Education Level	Equivalent to completing two years of college in the U.S.	Equivalent to completing a college degree in the U.S.
Gender	Female	Male

Immigrant 1 Immigrant 2

If you had to choose between them, which of these two immigrants should be given priority to come to the United States to live?







# **Conjoint Designs III**

# **Conjoint Designs III**

- As long as profiles are randomized, this is just a complex factorial design where we can estimate marginal effect of each attribute
  - Treatment-control SATE, conditional on all other randomized factors

# **Conjoint Designs III**

- As long as profiles are randomized, this is just a complex factorial design where we can estimate marginal effect of each attribute
  - Treatment-control SATE, conditional on all other randomized factors
- Assumptions:
  - Fully randomized profiles
  - No "carry-over" effects
  - No profile order effects

### **Questions?**

# **Activity!**

- Work in groups of 2–3
- Consider a research question of interest to you
  - Opinions on policies
  - Product purchasing decisions
  - Information selection
  - Attitudes toward out-groups
  - etc.
- Try to describe a basic conjoint design
  - What is the outcome?
  - What are the factors/features?

Stratification:Sampling::Blocking:Experiments

Stratification:Sampling::Blocking:Experiments

 Basic idea: randomization occurs within strata defined before treatment assignment

#### Stratification:Sampling::Blocking:Experiments

- Basic idea: randomization occurs within strata defined before treatment assignment
- CATE is estimate for each stratum; aggregated to SATE

#### Stratification:Sampling::Blocking:Experiments

- Basic idea: randomization occurs within strata defined before treatment assignment
- CATE is estimate for each stratum: aggregated to SATE
- Why?
  - Eliminate chance imbalances
  - Optimized for estimating CATEs
  - More precise SATE estimate

Exp.		Con	itrol		T	reat	mer	nt
1	М	М	М	М	F	F	F	F
2	M	M	M	F	M	F	F	F
3	M	M	F	F	M	M	F	F
4	M	F	F	F	M	M	M	F
5	F	F	F	F	M	M	M	M

```
# population of men and women
pop <- rep(c("Male", "Female"), each = 4)</pre>
```

# randomly assign into treatment and control
split(sample(pop, 8, FALSE), c(rep(0,4), rep(1,4)))

Obs.	$X_{1i}$	$X_{2i}$	$D_i$
1	Male	Old	0
2	Male	Old	1
3	Male	Young	1
4	Male	Young	0
5	Female	Old	1
6	Female	Old	0
7	Female	Young	0
8	Female	Young	1

- Blocking ensures ignorability of all covariates used to construct the blocks
- Incorporates covariates explicitly into the design

- Blocking ensures ignorability of all covariates used to construct the blocks
- Incorporates covariates explicitly into the design
- When is blocking *statistically* useful?

- Blocking ensures ignorability of all covariates used to construct the blocks
- Incorporates covariates explicitly into the *design*
- When is blocking *statistically* useful?
  - If those covariates affect values of potential outcomes, blocking reduces the variance of the SATE

- Blocking ensures ignorability of all covariates used to construct the blocks
- Incorporates covariates explicitly into the design
- When is blocking *statistically* useful?
  - If those covariates affect values of potential outcomes, blocking reduces the variance of the SATE
  - Most valuable in small samples

- Blocking ensures ignorability of all covariates used to construct the blocks
- Incorporates covariates explicitly into the *design*
- When is blocking *statistically* useful?
  - If those covariates affect values of potential outcomes, blocking reduces the variance of the SATE
  - Most valuable in small samples
  - Not valuable if all blocks have similar potential outcomes

## Statistical Properties I

Complete randomization:

$$SATE = \frac{1}{n_1} \sum Y_{1i} - \frac{1}{n_0} \sum Y_{0i}$$

Block randomization:

$$SATE_{blocked} = \sum_{1}^{J} \left( \frac{n_{j}}{n} \right) \left( \widehat{CATE}_{j} \right)$$

Moderators

Obs.	$X_{1i}$	$X_{2i}$	$D_i$	$Y_i$	CATE
1	Male	Old	0	5	
2	Male	Old	1	10	
3	Male	Young	1	4	
4	Male	Young	0	1	
5	Female	Old	1	6	
6	Female	Old	0	2	
7	Female	Young	0	6	
8	Female	Young	1	9	

Obs.	$X_{1i}$	$X_{2i}$	$D_i$	$Y_i$	CATE
1	Male	Old	0	5	_
2	Male	Old	1	10	5
3	Male	Young	1	4	
4	Male	Young	0	1	
5	Female	Old	1	6	
6	Female	Old	0	2	
7	Female	Young	0	6	
8	Female	Young	1	9	

Obs.	$X_{1i}$	$X_{2i}$	$D_i$	$Y_i$	CATE
1	Male	Old	0	5	5
2	Male	Old	1	10	5
3	Male	Young	1	4	2
4	Male	Young	0	1	3
5	Female	Old	1	6	
6	Female	Old	0	2	
7	Female	Young	0	6	
8	Female	Young	1	9	

cing Moderators

Obs.	$X_{1i}$	$X_{2i}$	$D_i$	$Y_i$	CATE
1	Male	Old	0	5	Е
2	Male	Old	1	10	5
3	Male	Young	1	4	2
4	Male	Young	0	1	3
5	Female	Old	1	6	1
6	Female	Old	0	2	4
7	Female	Young	0	6	
8	Female	Young	1	9	

Obs.	$X_{1i}$	$X_{2i}$	$D_i$	$Y_i$	CATE
1	Male	Old	0	5	E
2	Male	Old	1	10	5
3	Male	Young	1	4	2
4	Male	Young	0	1	3
5	Female	Old	1	6	1
6	Female	Old	0	2	4
7	Female	Young	0	6	2
8	Female	Young	1	9	3

$$SATE = \left(\frac{2}{8} * 5\right) + \left(\frac{2}{8} * 3\right) + \left(\frac{2}{8} * 4\right) + \left(\frac{2}{8} * 3\right)$$
= 3.75

## **SATE Estimation**

$$SATE = \left(\frac{2}{8} * 5\right) + \left(\frac{2}{8} * 3\right) + \left(\frac{2}{8} * 4\right) + \left(\frac{2}{8} * 3\right)$$
  
= 3.75

The blocked and unblocked estimates are the same here because Pr(Treatment) is constant across blocks and blocks are all the same size.

### SATE Estimation

- We can use weighted regression to estimate this in an OLS framework
- Weights are the inverse prob. of being treated w/in block
  - Pr(Treated) by block:
  - $p_{ij} = Pr(D_i = 1|J = j)$ Weight (Treated):  $w_{ij} = \frac{1}{p_{ij}}$
  - Weight (Control):  $w_{ij} = \frac{1}{1 p_{ii}}$

# Statistical Properties II

Complete randomization:

$$\widehat{SE}_{SATE} = \sqrt{\frac{\widehat{Var}(Y_0)}{n_0} + \frac{\widehat{Var}(Y_1)}{n_1}}$$

Block randomization:

$$\widehat{SE}_{SATE_{blocked}} = \sqrt{\sum_{1}^{J} \left(\frac{n_{j}}{n}\right)^{2} \widehat{Var}(SATE_{j})}$$

# Statistical Properties II

Complete randomization:

$$\widehat{SE}_{SATE} = \sqrt{\frac{\widehat{Var}(Y_0)}{n_0} + \frac{\widehat{Var}(Y_1)}{n_1}}$$

Block randomization:

$$\widehat{SE}_{SATE_{blocked}} = \sqrt{\sum_{1}^{J} \left(\frac{n_{j}}{n}\right)^{2} \widehat{Var}(SATE_{j})}$$

When is the blocked design more efficient?

### **Practicalities**

- Blocked randomization only works in exactly the same situations where stratified sampling works
  - Need to observe covariates pre-treatment in order to block on them
  - Work best in a panel context
- In a single cross-sectional design that might be challenging
  - Some software can block "on the fly"

## **Questions?**

#### Three Post-hoc Approaches

- Suggestive evidence
- Regression using treatment-by-covariate interactions

Moderators

Automated approaches

### Three Post-hoc Approaches

- Suggestive evidence
- Regression using treatment-by-covariate interactions
- Automated approaches
- (Replication and meta-analysis)

We can never know  $Var(TE_i)!$ 

We can never know  $Var(TE_i)!$  But...

Quantile-quantile plots

Equality of variance tests

We can never know  $Var(TE_i)$ ! But...


- Quantile-quantile plots
  - $\blacksquare$  Compare the distribution of  $Y_0$ 's to distribution of  $Y_1$ 's
  - If homogeneity, a vertical shift in  $Y_1$ 's
  - If heterogeneity, a slope  $\neq 1$
- Equality of variance tests


We can never know  $Var(TE_i)!$  But...

- Quantile-quantile plots
  - Compare the distribution of  $Y_0$ 's to distribution of  $Y_1$ 's
  - If homogeneity, a vertical shift in  $Y_1$ 's
  - $lue{}$  If heterogeneity, a slope eq 1
- Equality of variance tests
  - If homogeneity, variance should be equal
  - If heterogeneity, variances should differ

# **QQ** Plots

```
# y 0 data
set.seed(1)
n < -200
y0 \leftarrow rnorm(n) + rnorm(n, 0.2)
# y 1 data (homogeneous effects)
y1a \leftarrow y0 + 2 + rnorm(n, 0.2)
# y 1 data (heterogeneous effects)
y1b \leftarrow y0 + rep(0:1, each = n/2) + rnorm(n, 0.2)
qqplot(y0, y1a, pch=19, xlim=c(-3,5), ylim=c(-3,5), asp=1)
curve((x), add = TRUE)
qqplot(y0, y1b, pch=19, xlim=c(-3,5), ylim=c(-3,5), asp=1)
curve((x), add = TRUE)
```





# **Equality of Variance tests**

```
> var.test(y0, y1a)
```

F test to compare two variances

```
data: y0 and y1a
F = 0.60121, num df = 199, denom df = 199,
  p-value = 0.0003635
alternative hypothesis:
  true ratio of variances is not equal to 1
95 percent confidence interval:
 0.4549900 0.7944289
sample estimates:
ratio of variances
         0.6012131
```

# **Equality of Variance tests**

```
> var.test(y0, y1b)
```

F test to compare two variances

```
data: y0 and y1b
F = 0.53483, num df = 199, denom df = 199,
  p-value = 1.224e-05
alternative hypothesis:
  true ratio of variances is not equal to 1
95 percent confidence interval:
 0.4047531 0.7067133
sample estimates:
ratio of variances
         0.5348312
```

## **Questions?**

### Aside: Regression Adjustment in **Experiments, Generally**

- Recall the general advice that we do not need covariates in the regression to "control" for omitted variables (because there are none)
- Including covariates can reduce variance of our SATE by explaining more of the variation in Y

#### Scenario

Imagine two regression models. Which is correct?

- Mean-difference estimate of SATE is "not significant"
- Regression estimate of SATE. controlling for sex, age, and education, is "significant"

## **Scenario**

Imagine two regression models. Which is correct?

- Mean-difference estimate of SATE is "not significant"
- Regression estimate of SATE, controlling for sex, age, and education, is "significant"

This is a small-sample dynamic, so make these decisions pre-analysis!

- The regression paradigm allows us to estimate CATEs using interaction terms
  - $\blacksquare$  X is an indicator for treatment
  - *M* is an indicator for possible moderator

- The regression paradigm allows us to estimate CATEs using interaction terms
  - X is an indicator for treatment.
  - *M* is an indicator for possible moderator
- SATE:  $Y = \beta_0 + \beta_1 X + e$

- The regression paradigm allows us to estimate CATEs using interaction terms
  - X is an indicator for treatment
  - lacksquare M is an indicator for possible moderator
- SATE:  $Y = \beta_0 + \beta_1 X + e$
- CATEs:

$$Y = \beta_0 + \beta_1 X + \beta_2 M + \beta_3 X * M + e$$

- The regression paradigm allows us to estimate CATEs using interaction terms
  - X is an indicator for treatment
  - lacktriangleq M is an indicator for possible moderator
- SATE:  $Y = \beta_0 + \beta_1 X + e$
- CATEs:

$$Y = \beta_0 + \beta_1 X + \beta_2 M + \beta_3 X * M + e$$

- Homogeneity:  $\beta_3 = 0$
- Heterogeneity:  $\beta_3 \neq 0$

 Coefficients on moderators have no causal interpretation without further conditioning on observables

- Coefficients on moderators have no causal interpretation without further conditioning on observables
- Nearly unlimited potential moderators
  - First-order interactions with every covariate in dataset
  - Second-, third-order, etc. interactions
- Thus, multiple comparisons problem!

- Coefficients on moderators have no causal interpretation without further conditioning on observables
- Nearly unlimited potential moderators
  - First-order interactions with every covariate in dataset
  - Second-, third-order, etc. interactions
- Thus, multiple comparisons problem!
- Power (esp. if M is continuous)

## **Questions?**

### **BART**

■ Estimate CATEs in a fully automated fashion


#### **BART**

- Estimate CATEs in a fully automated fashion
- "Bayesian Additive Regression Trees"
  - Essentially an ensemble machine learning method


#### BART

- Estimate CATEs in a fully automated fashion
- "Bayesian Additive Regression Trees"
  - Essentially an ensemble machine learning method
- Iteratively split a sample into more and more homogeneous groups until some threshold is reached using binary (cutpoint) decisions
- Repeat this a bunch of times, aggregating across results

Moderators



Green & Kern. 2012. "Modeling Heterogeneous Treatment Effects in Survey Experiments with Bayesian Additive Regression Trees." *Public Opinion Quarterly* 76(3): 491–511.



### **Considerations**

- BART is totally automated, conditional on the set of covariates used
- Only really works with dichotomous covariates
- Not widely used or tested
- Totally post-hoc and atheoretical

## Replication!

- If we think effects are homogeneous (across SUTO), then replications in other SUTO conditions should provide us the same SATE (within sampling error)
- If we think effects are heterogeneous, then replications should give systematically different SATE (or CATE) estimates

### Replication!

■ If we think effects are homogeneous (across SUTO), then replications in other SUTO conditions should provide us the same SATE (within sampling error)

Moderators

- If we think effects are heterogeneous, then replications should give systematically different SATE (or CATE) estimates
  - Identify those patterns of heterogeneity using meta-analysis

## Replication!

- If we think effects are homogeneous (across SUTO), then replications in other SUTO conditions should provide us the same SATE (within sampling error)
- If we think effects are heterogeneous, then replications should give systematically different SATE (or CATE) estimates
  - Identify those patterns of heterogeneity using meta-analysis
  - Regress effect estimates from multiple studies on SUTO features of each study

### **Conclusion**

■ Do we want to know SATE, CATE(s), or both?

### Conclusion

- Do we want to know SATE, CATE(s), or both?
- Decide in advance
  - Include in protocol
  - Design study to estimate CATE(s)

### **Conclusion**

- Do we want to know SATE, CATE(s), or both?
- Decide in advance
  - Include in protocol
  - $\blacksquare$  Design study to estimate CATE(s)
- Estimation of CATEs
  - Block randomization
  - Post-hoc procedures

### **Questions?**



# Apparent Satisficing

- Some common measures:
  - "Straightlining"
  - Non-differentiation
  - Acquiescence
  - Nonresponse
  - DK responding
  - Speeding
- Difficult to detect and distinguish from "real" responses

# Metadata/Paradata

- Timing
  - Some survey tools will allow you to time page
  - Make a prior rules about dropping participants for speeding

# Metadata/Paradata

- Timing
  - Some survey tools will allow you to time page
  - Make a prior rules about dropping participants for speeding
- Mousetracking or eyetracking
  - Mousetracking is unobtrusive
  - Eyetracking requires participants opt-in

# Metadata/Paradata

- Timing
  - Some survey tools will allow you to time page
  - Make a prior rules about dropping participants for speeding
- Mousetracking or eyetracking
  - Mousetracking is unobtrusive
  - Eyetracking requires participants opt-in
- Record focus/blur browser events

### **Direct Measures**

How closely have you been paying attention to what the questions on this survey actually mean?

### **Direct Measures**

- How closely have you been paying attention to what the questions on this survey actually mean?
- While taking this survey, did you engage in any of the following behaviors? Please check all that apply.
  - Use your mobile phone
  - Browse the internet
  - . . . .

#### Instructional Manipulation Check

We would like to know if you are reading the questions on this survey. If you are reading carefully, please ignore this question, do not select any answer below, and click "next" to proceed with the survey.

Strongly disagree Somewhat disagree Neither agree nor disagree Somewhat agree Strongly agree

#### Instructional Manipulation Check

Do you agree or disagree with the decision to send British forces to fight ISIL in Syria? We would like to know if you are reading the questions on this survey. If you are reading carefully, please ignore this question, do not select any answer below, and click "next" to proceed with the survey.

Strongly disagree Somewhat disagree Neither agree nor disagree Somewhat agree Strongly agree

