

Summarized chain of thought (Isomorphism of free group factors)

OpenAI

Problem

Determine whether $L(F_2)$ and $L(F_3)$ are isomorphic as von Neumann algebras, and prove the answer. Here F_n is the discrete free group on n generators. Its left regular representation on the complex Hilbert space $\ell^2(F_n)$ is defined by $(\lambda_n(g)\xi)(h) = \xi(g^{-1}h)$. Define $L(F_n) = \{\lambda_n(g): g \in F_n\}''$, the double commutant, equivalently the weak operator closure of the unital complex algebra generated by these unitaries.

The precise question is whether there exists a unital normal $*$ -isomorphism $\Phi: L(F_2) \rightarrow L(F_3)$. Normal means preservation of suprema of bounded increasing nets of self-adjoint operators, equivalently ultraweak continuity. The isomorphism may send the canonical group generators to arbitrary elements: no requirement to respect a chosen group presentation is imposed. The canonical normalized traces are $\tau_n(T) = \langle T \delta_e, \delta_e \rangle$; a normal $*$ -isomorphism necessarily preserves them.

1 From factor rigidity to the missing width bound

The assistant first seeks an intrinsic obstruction to an isomorphism of the two free group factors. From Dykema–Rădulescu’s rescaling results it takes the dichotomy that interpolated free group factors are either all isomorphic or pairwise nonisomorphic, together with the equivalent requirement that $Q_1 * Q_2 \cong Q_1 * Q_2 * L(F_\infty)$ for every pair of type II_1 factors [16]. It considers transferring Boutonnet–Drimbe–Ioana–Popa’s distinction between nonseparable abelian free powers to the separable problem [5]. It examines Dykema–Zhao’s nonseparable free-product constructions as a possible comparison, but finds no way to transport an arbitrary separable-factor isomorphism to those objects [17]. Peterson’s separation of $L(SL_3(\mathbb{Z}))$ from $L(F_2)$ by elementary equivalence likewise does not give it a distinction between ranks two and three [40].

The assistant next tries rigid free-product components and Gaussian crossed products. Popa’s discussion of generation and rigidity motivates these reductions [41]. It uses the free-group-factor dichotomy, also recorded in Dykema’s interpolation work, to consider arbitrarily many semicircular generators [18]. The assistant finds that rigidity leaves an uncontrolled conjugating unitary that might encode the extra free directions. It also finds no way to induce an isomorphism of the proposed crossed products, so it cannot apply the unique-Cartan theorem of Popa–Vaes to its Gaussian construction [42]. It also considers differentiating a trace-scaling flow, but Chakraborty–Novac’s discussion ties the existence of such a flow on the stabilization of $L(F_2)$ back to the same isomorphism problem [11]. Jung’s strong 1-boundedness distinguishes a different class of factors, while the 1-bounded entropy developed by Hayes has the same infinite value throughout the free-group family [33, 26]. Generator-dependent entropy quantities encounter another obstacle:

VERBATIM EXCERPT

Unknown W^* invariance due to bad approximations. Could try to circumvent using local topological dimension/mean dimension rather than Minkowski.

The replacement proposal studies matrix microstates through width dimension, the dimension needed for continuous encoding with small fibers. If s free semicircular generators were recoverable from $n < s$

self-adjoint generators, polynomial approximation could give a width upper bound nN^2 for $N \times N$ microstates. A contradictory lower bound must remain proportional to sN^2 as moment constraints become arbitrarily stringent.

The assistant finds that high probability does not supply that bound: a commutative rounding example retains approximate recovery while sharply reducing coarse dimension, and its finite-quotient models lose too many parameters. The assistant identifies a possible distinction from unrestricted continuous encoding:

VERBATIM EXCERPT

Crucially P, Q are unitarily equivariant trace-polynomial approximants with degree independent of N . Entrywise rounding isn't implementable by polynomial operations (coordinate-dependent). Might exploit equivariance, dimension-theoretic invariance reinforced by matrix amplification.

The assistant pursues equivariance, then finds that its tensor-twisted embeddings preserve free distributions while retaining infinitely many Fourier coefficients, preventing the parameter count it sought. The assistant leaves the width lower bound unresolved and considers Connes–Shlyakhtenko's algebraic L^2 -homology as a possible generator-independent obstruction, while noting that Popa–Vaes's vanishing of continuous first L^2 -cohomology prevents the continuous variant from distinguishing the factors [15, 43].

2 Small projections and the recurring boundary obstruction

The assistant adjoins nonseparable auxiliary factors, then finds that its extensions reduce to a separable support algebra. It cannot transport the invariant distinguishing nonseparable abelian free powers [5] through an arbitrary separable isomorphism. Popa–Vaes's vanishing theorem [43] prompts it to consider derivations on restricted domains.

Factoring polynomial approximations through fewer variables bounds free-Jacobian rank, but the assistant cannot control derivatives in the limit. Its commutative checkerboard example suggests that strong convergence of bounded kernels need not preserve module rank; gradient relations supply no uniform estimate.

Proper proximality [6] suggests a boundary approach, with Mei–Ricard's free Hilbert transforms [38] motivating regularity. Using Julg–Valette's tree operator [32], the assistant considers a free group factor $M = L(\mathbb{F}_r)$ and compares maps from $H = L^2(M)$ to H^r with weakened compactness of commutators. It constructs a shift on normalized vectors supported on shrinking orthogonal projections, concludes that the shift removes the index defect while preserving regularity, and abandons the invariant.

The Dykema–Rădulescu absorption equivalence [16] next suggests contradicting $Q_1 * Q_2 \cong Q_1 * Q_2 * L(\mathbb{F}_\infty)$ with rigid components. Under an affirmative answer, this absorption would apply to arbitrary type II_1 factors Q_1, Q_2 . It therefore hopes to choose components whose rigidity would expose the additional free directions. The assistant finds that Ioana–Peterson–Popa rigidity [30] tracks those components without detecting the extra free part. Returning to algebraic L^2 -homology [15], it encounters derivations not determined by weakly dense generators.

It compares Ringrose's Banach-bimodule continuity theorem [46] with affiliated targets. Shen's generator invariant [48] and its amplification scaling [19] leave positivity unknown. The assistant finds that Ge–Popa's tensor-product single-generation result [23] blocks its finite-generation shortcut, while Jung's ultrapower-embedding characterization of amenability [34] limits its back-and-forth construction.

Comparing its inclusions with standard embeddings [20], it tries placing copies of $L(\mathbb{F}_\infty)$ near finite sets. Adjoining missing coordinates imposes conjugacy relations, and it encounters spectral-gap obstructions to perturbing them into free bases. Gaboriau's measure-equivalence invariance of ℓ^2 -Betti numbers [21] and treeing-cost argument [22] suggest a coupling approach. The assistant identifies the

missing step: measurable relabeling gives exact agreement on large sets, whereas its operator approximations give only L^2 closeness.

Using Dykema's compression formula [18], the assistant considers self-similar corners, but identifying a corner with the original factor would still require generator collapse. A matrix-amplification attempt studies a distinguished row. Stripping the first letter of a Wick expansion has bounded total L^2 norm, unlike its earlier derivatives. Composing row formulas for two semicircular presentations might force equal generator counts. But the assistant notices that bulk errors on a projection of trace about $1/k$ produce order-one row errors, preventing its composition argument. Reinterpreting the row space as $L^2(M) \ominus \mathbb{C}$, it recognizes the earlier Julg–Valette obstruction.

Finally, it tries matching bulk embeddings by automorphisms and passing to an ultralimit where a module maps properly into itself. It leaves open the required operator-norm control and the contradiction that this proper inclusion would provide.

3 From rank obstructions to conditional absorption

The assistant's Fock-space construction produces extra representation sectors, but it finds no contradiction: the inverse transformation need not preserve the selected subspace. Applying nonseparable free-power rigidity [5] would require an isomorphism to preserve enlarged components. It finds neither that preservation nor a generalized-wreath-product theorem allowing arbitrary free-group bases [31]. Rough generator changes also obstruct its use of Voiculescu's norm-ideal invariants [51].

Replicating an isomorphism preserves its free-entropy dimension ratio; iteration increases reconstruction complexity alongside the gap. The assistant cannot control continuity moduli or preserve Fock-space compactness. Coordinatewise quantization also prevents its inference from large microstate packing numbers to a topological obstruction.

For Fox derivatives, it compares Thom's affiliated-operator cohomology [50], Alekseev–Kyed automatic continuity [1], Alekseev's account [2], and Popa–Vaes vanishing [43], without finding the required closability statement. The prefix–suffix structure of the derivatives suggests isolating coefficients that retain a common group-theoretic relation. The assistant hopes that this filtering will force a vanishing relation strong enough to compare generator counts. Filtering derivative limits onto a diagonal subgroup leaves a cycle space of dimension $n - 1$. It also notices a domain obstacle:

VERBATIM EXCERPT

Even if prove D closable in measure, does it solve problem? Isom T arbitrary M elements not necessarily in domain.

Bermúdez's relative homology [4] supplies no distinction it can establish over a hyperfinite subfactor.

The assistant then modifies corners of $M_0 = Q_1 * Q_2$. For projections e_i of trace $1/k$, it retains

$$Q_i^0 = (1 - e_i)Q_i(1 - e_i) \oplus \mathbb{C}e_i, \quad P_0 = Q_1^0 * Q_2^0.$$

Invoking Avitzour's criterion [3] for factoriality, it tries recovering each Q_i by adjoining a partial isometry and comparing it with one in the intermediate factor. For a suitable corner projection p , its proposed reduced-word calculation leads to

$$pM_0p \cong pP_0p * L(\mathbb{F}_2).$$

Assuming an affirmative isomorphism answer and this calculation, it seeks absorption of $L(\mathbb{F}_\infty)$ through maps fixing P_0 , approximating the identity uniformly on each component's unit ball. It finds no compactness of preimages or surjectivity of a limiting map; property (T) aligns components without

controlling their free product. Dykema–Rădulescu compression and absorption formulas inform this construction [16].

Jung’s strong 1-boundedness [33] and the Jung–Shlyakhtenko property-(T) bound [35] do not give it control of arbitrary free-product generators. It rejects the Christensen–Wang complemented-subspace route [14], finding the required complementedness obstructed by an expected copy of $L(\mathbb{F}_\infty)$.

Using interpolation formulas from Dykema and Dykema–Zhao [18, 17], it writes $N_r = L(\mathbb{F}_r)$ and obtains

$$N_r^t \cong N_{1+(r-1)/t^2}, \quad N_r * N_s \cong N_{r+s},$$

$$N_2 \cong N_3 \iff 1/\sqrt{2} \in \mathcal{F}(N_2), \quad \mathcal{F}(N_2) = \{t > 0 : N_2^t \cong N_2\}.$$

Here N_r^t denotes amplification by t , and $\mathcal{F}(N_2)$ is the fundamental group of N_2 . The reduction expresses the desired isomorphism through membership of a particular scalar in this group, but the assistant finds no way to establish or exclude that membership. It next investigates whether nonseparability obstructs asymptotic freeness in its conditional construction, continuing reduced-word calculations without obtaining an isomorphism or contradiction. It cites Schafhauser–Tikuisis–White’s problem list for the unresolved question [47].

4 Embedding obstructions meet uncontrolled changes of generators

The assistant used long alternating words with nonperiodic letters to conjugate $W^*(a, b^2)$ freely away from component ultrapowers, defeating its intended obstruction. Its cancellation bound depended on syllable length rather than the individual letters, with the outer letter orthogonal to the smaller algebra. It found no suitable free-product structure for nonseparable intertwining [5], and questioned whether deformation arguments forced component embeddings [30].

It tried conjugating a free shift to its square, finding no entropy distinction [8]. Analytic Gibbs transport preserved the number of variables [24]; supported semicirculars suggested hiding directions on small supports [18].

Commuting Haar witnesses c_1, c_2 , free from b , gave a free triple $b, c_1 b c_1^*, c_2 b c_2^*$. Using free dimension and standard embeddings [20], it compared partition dimensions $3d$ and $1 + d$, where $d = 1 - \sum_j t_j^2$. Although $1/2 < d < 2/3$ suggested room in rank two, it could not identify the embeddings. The assistant recognized that an abstract factor isomorphism need not preserve the selected partition algebra or the witnesses used to distinguish its presentations:

VERBATIM EXCERPT

| Joint law free P , embeddings not approximately conjugate. Already nonstandard vs standard yields.

Jung’s amenability characterization reinforced that obstacle [34]. The assistant revisited the problem survey [47], interpolation [18, 44], absorption equivalence [16], and nonseparable rescaling [17].

Recovering extra generators from a pair required uncontrolled functions. Shrinking-corner encodings consumed recovery information; rescaling magnified errors. Jacobian ranks failed to pass through approximations without derivative control. Continuous-cohomology vanishing discouraged extending derivatives [43]; strong 1-boundedness did not distinguish these ranks [33]. The assistant sought non-commutative versions of cost and ℓ^2 -invariance [22, 21], then tested trace scaling and generator invariants without an obstruction [45, 48].

Fock-space creation rows gave a rectangular Hilbert-space unitary, but no finite-factor dimension comparison. Free Hilbert transforms suggested prefix regularity [38]; the tree operator returned it to the failed index approach [32].

Iterating a hypothetical transformation from n semicircular generators to $m > n$ would produce m^k outputs from n^k inputs. But errors obeyed $e_j \leq \delta_j + L_j e_{j-1}$, with uncontrolled Lipschitz constants:

VERBATIM EXCERPT

| Since f fixed measurable, $L(\delta)$ can grow without bound arbitrarily; automorphism may pathological.

Fixing final accuracy forced increasingly precise earlier layers. Free subgroup words restored polynomial control but lost microstate extendability. Orbital entropy did not resolve invariance under rough generator changes [28].

The assistant found that tail permutations were outer and supplied no implementing unitaries for central sequences. Its Baire argument lost a uniform interior radius. Even assuming Sobolev regularity, closability of δ did not imply closability of $P\delta$ for arbitrary module projections. It abandoned graph-product rigidity [10, 29]; maximal-amenable uniqueness suggested incidence geometry but no rank calculation [27]. Free Stein dimension supplied algebraic invariance, while the computation and monotonicity needed for a limiting extension remained unavailable to it [12, 13].

Finally, it restricted attention to finite Fourier-support images of canonical unitaries. It hoped subgroup surjectivity and abelianization would detect rank there, while continuing to seek control of arbitrary generator changes without quantitative regularity.

5 Recovering a missing root while losing freeness

The assistant moved from possible rank invariants for free group factors $L(\mathbb{F}_n)$ toward constructive changes of generators. It first examined matrix entries and free compression: Wick words follow paths through matrix indices, and free compression interpolates between a change of generators and its covariance. It found no rank obstruction: rare contractions can preserve invertibility despite unfavorable dimension counts, while iterative encodings force increasingly rough decoders.

It reconsidered Ringrose's automatic continuity theorem [46], Alekseev–Kyed's measure-continuity results [1], and Popa–Vaes's vanishing theorem [43], but could not extend their continuity hypotheses to arbitrary affiliated-valued derivations. It also revisited the Peterson–Thom property and its structural consequences [27] as possible controls on rough changes of generators, without obtaining a rank invariant.

The assistant next used the Boutonnet–Drimbe–Ioana–Popa sans-rank, which distinguishes free powers of a nonseparable diffuse abelian algebra [5]. Dykema–Zhao's rescaling formulas supplied a further test, but changing the separable free-factor parameter left the nonseparable component count unchanged [17]. The assistant found that Nielsen substitutions mixed components and destroyed the commutation relations required to transfer a separable isomorphism to those enlargements. It suspected that enlarging the construction to tolerate such substitutions would erase its finite component count, but did not calculate the rank of the enlargement. The Dykema–Rădulescu equivalence links collapse of free group factors to absorption of $L(\mathbb{F}_\infty)$ by free products of two factors [16]. It tried to exploit Ioana–Peterson–Popa's rigidity results for these components [30], but separate conjugacies did not control the free complement.

A constructive turn drew on the Ornstein–Weiss factor-map example [39] and Bowen's Bernoulli-shift classification [7] as classical comparisons, then used free Haar vertex variables x_g on the Cayley tree of $\mathbb{F}_d$. Edge increments $x_g^* x_{gs_i}$ are free, but reconstruction requires the missing root x_e . A classical analogy suggests breaking the symmetry:

VERBATIM EXCERPT

Aha! For classical vertex labels in a group with non-uniform distribution, edge differences determine absolute labels almost surely because the global shift ambiguity can be resolved by empirical frequencies (only one shift yields correct distribution).

For free vertex unitaries with nonzero mean β , telescoping products and weak convergence give $x_g^* x_h \rightarrow \beta x_g^*$, recovering every vertex from the increments. The cost is correlations between neighboring edges: generation returns while exact freeness disappears. It considered free monotone transport [24] to remove the correlations, but did not establish the required regularity. It then argued that averages of conjugate site variables converge in operator norm, strengthening the recovery argument.

Returning to the missing-root construction, the assistant argued that a spectral gap bars nearby free generating tuples when $d \geq 2$: the root would become an impossible approximate intertwiner. It concluded:

VERBATIM EXCERPT

Thus generating set (nonfree) accumulates at a free tuple separated from all free generating tuples; no Baire smoothing there.

The assistant thus distinguished nearly free generators from genuinely free generating tuples, while leaving its original isomorphism question unresolved. Its return to the partial reduction also used the problem-status discussion of Schafhauser, Tikuisis, and White [47].

6 From width obstructions to a proposed Gaussian absorption flow

The assistant again sought a generator-independent microstate invariant:

VERBATIM EXCERPT

Difference $P(V)$ - U small but derivative of difference can be huge; no closability. Same old.

Jung's hyperfinite inequality did not carry its finite-Borel-operation estimates through strong limits [36]. It recalled zero-entropy free shifts [49] and found that separable maximal abelian subalgebras lost the nonseparable argument's cardinal distinction [5]. Projection-sparsity and amplification estimates still lacked a positive lower bound [48, 19]; the commutator-based candidate invariant again confronted passage from algebraic to von Neumann generation [15].

Urysohn width suggested obstructing an encoding of free tuples through fewer matrix generators. Spectral clipping and Borsuk–Ulam controlled an entire operator norm ball, but the assistant could not restrict the obstruction uniformly to a prescribed free law. Projection onto one coordinate defeated its global-negation argument. It compared packing rank and strong 1-boundedness [37, 33], noted transport's regularity restrictions [24], and found insufficient matrix-dimension growth in mixtures of ultra-power embeddings [9].

It then tried absorbing a free generator through small perturbations. Straight cocycle paths produced nearly translation-invariant elements, obstructing approximation of a free complement. For Haar unitaries U_i and a complementary semicircular S , it proposed

$$\dot{U}_i = i \left(\sum_g h_i(g) U_g S U_g^* \right) U_i, \quad \dot{S} = 0,$$

with finitely supported real coefficients. Comparing cyclomorphy [52] and coalgebra constructions [53], it questioned its divergence calculation and preferred direct cumulant pairing. It recast derivatives of

$S_g = U_g S U_g^*$ as single-variable conjugations: the same-variable coefficient commuted with S_g , while the remaining conjugator lay in the other variables' algebra.

Fox derivatives of long words were intended to amplify small coefficients into displacement involving S . For free prefixes p_r , target amplitude R , and cutoff d , it set

$$P = 1 + \sum_{r=1}^k \lambda(p_r), \quad Q = PP^*, \quad E = 1_{(dk, \infty)}(Q), \quad h = RP^*Q^{-1}E\delta_e.$$

It obtained $Ph = RE\delta_e$ and $\|h\|_2^2 \leq R^2/(dk)$. Haagerup's inequality bounded the prefix sum by $2\sqrt{k}$ [25]. Reduced-word counting suggested Catalan moments and a Marchenko–Pastur limit without an atom at zero. Choosing a cutoff, then a word length, and finally finite real approximants was intended to make both h and $Ph - R\delta_e$ small. Preserving the semicircular law would keep residuals small without differentiating them.

The assistant proposed a measurable Haar reparameterization of S and a free-basis change to approximate the specified extra generator. Diagonal iteration budgeted future displacements against earlier polynomial lengths to seek a generating limiting tuple. The assistant wanted each later perturbation to preserve the approximations already obtained for earlier targets. Interpolation and amplification would transfer its rank-three/rank-four comparison to ranks two and three [18, 44]. Checking operator-norm approximation against reduced group C^* -algebra invariants led it to distinguish measurable coordinate changes from continuous calculus. It then asked whether a continuous Haar function could generate the semicircular algebra while approximating the oscillatory exponential.

7 Logarithmic perturbations and limiting generation

The assistant sought to absorb one free Haar generator C into small perturbations of the others, denoted $A_1, \dots, A_n$, with A_g the corresponding reduced word. It replaced a semicircular auxiliary variable by the bounded Borel logarithm $S = \text{Arg}(C)$, so that $\tau(S) = 0$ and $e^{iS} = C$. For the free conjugates $S_g = A_g S A_g^*$, it split multiplication into creation, annihilation, and diagonal terms to obtain the proposed estimate

$$\left\| \sum_g h(g) S_g \right\| \leq 3\pi \|h\|_2.$$

This would control generator displacement using small coefficient vectors without requiring semicircular stability.

The assistant also considered Shen's single-generation results [48], without establishing an application that would settle this generator-count comparison.

The argument combined a polynomial flow with a prefix-sum operator. The assistant argued for trace preservation by conjugating individual free variables by elements generated by the others, then used polynomial degree estimates to justify analytic continuation of trace moments. Catalan moment counting and a spectral cutoff inverse were intended to produce coefficients that moved generators little while moving a long word substantially.

The assistant corrected the target of the approximation: the moved word had to approach CA_w , not C , since the flow fixed C and preserved its freeness from the moved generators. An inverse Nielsen automorphism would convert that estimate into approximation of C . Summable perturbations were then meant to yield a norm-convergent tuple. For generation of the ambient von Neumann algebra, the assistant used a separate diagonal argument establishing L^2 -density of the word algebra; recovering earlier coordinates could require discontinuous logarithms, so operator-norm convergence alone did not give norm generation.

In comparing its vector fields with existing constructions, the assistant recalled Voiculescu’s *Cyclo-morphy* as a framework for trace-preserving deformations [52]. It also revisited the nonseparable free-product theorem of Boutonnet, Drimbe, Ioana, and Popa [5]: its proposed countable diagonal construction could not absorb an entire nonseparable abelian component.

The remaining compatibility questions concerned relative freeness, fixed-subalgebra constraints, and Julg–Valette index obstructions. The assistant used Voiculescu’s noncommutative Weyl–von Neumann theorem to argue that absorption of an ample representation defeated the simplest proposed index argument [54]. It then recalled the L^p -bounded free Hilbert transforms of Mei and Ricard while testing whether additional regularity could restore an obstruction [38]. For the fixed algebra $N = W^*(A_2, \dots, A_n, C)$, the assistant argued that its chosen positive long word x was Haar free from N , and that multiplying it by $C \in N$ preserved this relative distribution. It therefore found no contradiction from these moment identities. Large finite coefficient sums could make the local analytic radius very small; the assistant answered that a positive radius and global real-time existence would still permit continuation to time one.

To transfer its proposed adjacent-rank comparison to the requested ranks, the assistant invoked the interpolation, amplification, and free-product formulas of Dykema and Rădulescu, from Dykema’s *Interpolated free group factors* [18], using the accounts in Dykema–Zhao’s free-product and rescaling paper and Dykema–Rădulescu’s *Rescalings of free products of II_1 -factors* [17, 16].

The assistant ended with a proposed affirmative resolution based on the perturbation and limiting-generation argument: a bijective unital normal $*$ -isomorphism preserving the normalized traces. This conclusion depends on the trace-preserving flow and limiting-generation steps described above.

References

- [1] Vadim Alekseev and David Kyed. *Measure continuous derivations on von Neumann algebras and applications to L^2 -cohomology*. Journal of Operator Theory **73**(1) (2015), 91–111. <https://doi.org/10.7900/jot.2013sep23.2018>.
- [2] Vadim Alekseev. *Groups, operator algebras and approximation*. Habilitation thesis, Technische Universität Dresden (2019). Submitted December 2019; published in the university repository in 2021. <https://nbn-resolving.org/urn:nbn:de:bsz:14-qucosa2-761277>.
- [3] Daniel Avitzour. *Free products of C^* -algebras*. Transactions of the American Mathematical Society **271**(2) (1982), 423–435. <https://doi.org/10.2307/1998890>.
- [4] Miguel Bermudez. *L^2 -homology for inclusions of von Neumann algebras*. arXiv preprint arXiv:1403.6044 (2014). <https://arxiv.org/abs/1403.6044>.
- [5] Rémi Boutonnet, Daniel Drimbe, Adrian Ioana, and Sorin Popa. *Non-isomorphism of A^{*n} , $2 \leq n \leq \infty$, for a non-separable abelian von Neumann algebra A* . Geometric and Functional Analysis **34**(2) (2024), 393–408. <https://doi.org/10.1007/s00039-024-00669-8>.
- [6] Rémi Boutonnet, Adrian Ioana, and Jesse Peterson. *Properly proximal groups and their von Neumann algebras*. Annales Scientifiques de l’École Normale Supérieure **54**(2) (2021), 445–482. <https://doi.org/10.24033/asens.2462>.
- [7] Lewis Bowen. *A measure-conjugacy invariant for free group actions*. Annals of Mathematics **171**(2) (2010), 1387–1400. <https://doi.org/10.4007/annals.2010.171.1387>.
- [8] Nathaniel P. Brown, Ken Dykema, and Dimitri Shlyakhtenko. *Topological Entropy of Free Product Automorphisms*. Acta Mathematica **189**(1) (2002), 1–35. <https://doi.org/10.1007/bf02392643>.

- [9] Nathaniel P. Brown. *Topological dynamical systems associated to II_1 factors*. Advances in Mathematics **227**(4) (2011), 1665–1699. <https://doi.org/10.1016/j.aim.2011.04.003>.
- [10] Martijn Caspers and Enli Chen. *Internal graphs of graph products of hyperfinite II_1 -factors*. arXiv preprint arXiv:2505.05179 (2025). Revised 2026; to appear in Journal of Noncommutative Geometry. <https://arxiv.org/abs/2505.05179>.
- [11] Soham Chakraborty and Sergiu Novac. *Non-uniqueness of continuous trace-scaling flows on certain full factors*. arXiv preprint arXiv:2608.15385 (2026). <https://arxiv.org/abs/2608.15385>.
- [12] Ian Charlesworth and Brent Nelson. *Free Stein Irregularity and Dimension*. Journal of Operator Theory **85**(1) (2021), 101–133. <https://doi.org/10.7900/jot.2019aug29.2271>.
- [13] Ian Charlesworth and Brent Nelson. *On Free Stein Dimension*. Indiana University Mathematics Journal **74**(4) (2025), 883–918. <https://doi.org/10.1512/iumj.2025.74.60370>.
- [14] Erik Christensen and Liguang Wang. *On a von Neumann algebra which is a complemented subspace*. arXiv preprint arXiv:1312.0450 (2013). <https://arxiv.org/abs/1312.0450>.
- [15] Alain Connes and Dimitri Shlyakhtenko. *L^2 -Homology for von Neumann Algebras*. Journal für die reine und angewandte Mathematik **586** (2005), 125–168. <https://doi.org/10.1515/crll.2005.2005.586.125>.
- [16] Ken Dykema and Florin Rădulescu. *Rescalings of free products of II_1 factors*. Proceedings of the American Mathematical Society **131**(6) (2003), 1813–1816. <https://doi.org/10.1090/S0002-9939-02-06749-7>.
- [17] Ken Dykema and Junchen Zhao. *Free products and rescalings involving non-separable abelian von Neumann algebras*. Journal of Functional Analysis **290**(4) (2026), 111264. <https://doi.org/10.1016/j.jfa.2025.111264>.
- [18] Ken Dykema. *Interpolated free group factors*. Pacific Journal of Mathematics **163**(1) (1994), 123–135. <https://doi.org/10.2140/pjm.1994.163.123>.
- [19] Ken Dykema, Allan Sinclair, Roger Smith, and Stuart White. *Generators of II_1 Factors*. Operators and Matrices **2**(4) (2008), 555–582. <https://doi.org/10.7153/oam-02-35>.
- [20] Ken Dykema. *Free products of hyperfinite von Neumann algebras and free dimension*. Duke Mathematical Journal **69**(1) (1993), 97–119. <https://doi.org/10.1215/S0012-7094-93-06905-0>.
- [21] Damien Gaboriau. *Invariants ℓ^2 de relations d’équivalence et de groupes*. Publications Mathématiques de l’IHÉS **95** (2002), 93–150. <https://doi.org/10.1007/s102400200002>.
- [22] Damien Gaboriau. *Coût des relations d’équivalence et des groupes*. Inventiones Mathematicae **139**(1) (2000), 41–98. <https://doi.org/10.1007/s002229900019>.
- [23] Liming Ge and Sorin Popa. *On some decomposition properties for factors of type II_1* . Duke Mathematical Journal **94**(1) (1998), 79–101. <https://doi.org/10.1215/S0012-7094-98-09405-4>.
- [24] Alice Guionnet and Dimitri Shlyakhtenko. *Free Monotone Transport*. Inventiones Mathematicae **197**(3) (2014), 613–661. <https://doi.org/10.1007/s00222-013-0493-9>.
- [25] Uffe Haagerup. *An example of a non nuclear C^* -algebra, which has the metric approximation property*. Inventiones Mathematicae **50**(3) (1978), 279–293. <https://doi.org/10.1007/BF01410082>.

- [26] Ben Hayes. *1-bounded entropy and regularity problems in von Neumann algebras*. International Mathematics Research Notices **2018**(1) (2018), 57–137. <https://doi.org/10.1093/imrn/rnw237>.
- [27] Ben Hayes, David Jekel, and Srivatsav Kunnawalkam Elayavalli. *Consequences of the random matrix solution to the Peterson-Thom conjecture*. Analysis & PDE **18**(7) (2025), 1805–1834. <https://doi.org/10.2140/apde.2025.18.1805>.
- [28] Fumio Hiai, Takuho Miyamoto, and Yoshimichi Ueda. *Orbital approach to microstate free entropy*. International Journal of Mathematics **20**(2) (2009), 227–273. <https://doi.org/10.1142/s0129167x09005261>.
- [29] Camille Horbez and Adrian Ioana. *Rigidity for graph product von Neumann algebras*. arXiv preprint arXiv:2508.03662 (2025). <https://arxiv.org/abs/2508.03662>.
- [30] Adrian Ioana, Jesse Peterson, and Sorin Popa. *Amalgamated free products of weakly rigid factors and calculation of their symmetry groups*. Acta Mathematica **200**(1) (2008), 85–153. <https://doi.org/10.1007/s11511-008-0024-5>.
- [31] Adrian Ioana, Sorin Popa, and Stefaan Vaes. *A class of superrigid group von Neumann algebras*. Annals of Mathematics **178**(1) (2013), 231–286. <https://doi.org/10.4007/annals.2013.178.1.4>.
- [32] Pierre Julg and Alain Valette. *K-theoretic amenability for $SL_2(\mathbb{Q}_p)$, and the action on the associated tree*. Journal of Functional Analysis **58**(2) (1984), 194–215. [https://doi.org/10.1016/0022-1236\(84\)90039-9](https://doi.org/10.1016/0022-1236(84)90039-9).
- [33] Kenley Jung. *Strongly 1-bounded von Neumann algebras*. Geometric and Functional Analysis **17**(4) (2007), 1180–1200. <https://doi.org/10.1007/s00039-007-0624-9>.
- [34] Kenley Jung. *Amenability, tubularity, and embeddings into R^ω* . Mathematische Annalen **338**(1) (2007), 241–248. <https://doi.org/10.1007/s00208-006-0074-y>.
- [35] Kenley Jung and Dimitri Shlyakhtenko. *Any generating set of an arbitrary property T von Neumann algebra has free entropy dimension ≤ 1* . Journal of Noncommutative Geometry **1**(2) (2007), 271–279. <https://doi.org/10.4171/JNCG/7>.
- [36] Kenley Jung. *A hyperfinite inequality for free entropy dimension*. Proceedings of the American Mathematical Society **134**(7) (2006), 2099–2108. <https://doi.org/10.1090/s0002-9939-06-08237-2>.
- [37] Kenley Jung. *The Rank Theorem and L^2 -invariants in Free Entropy: Global Upper Bounds*. arXiv preprint arXiv:1602.04726 (2016). <https://arxiv.org/abs/1602.04726>.
- [38] Tao Mei and Éric Ricard. *Free Hilbert Transforms*. Duke Mathematical Journal **166**(11) (2017), 2153–2182. <https://doi.org/10.1215/00127094-2017-0007>.
- [39] Donald S. Ornstein and Benjamin Weiss. *Entropy and isomorphism theorems for actions of amenable groups*. Journal d’Analyse Mathématique **48**(1) (1987), 1–141. <https://doi.org/10.1007/BF02790325>.
- [40] Jesse Peterson. *On ultraproduct approximations and property (T) factors*. arXiv preprint arXiv:2605.16669 (2026). <https://arxiv.org/abs/2605.16669>.
- [41] Sorin Popa. *Some open problems on III factors of group actions*. Talk at the Fields Institute (2007). 29 October 2007. https://www.fields.utoronto.ca/talks/Some-open-problems-III_1-factors-group-actions.

- [42] Sorin Popa and Stefaan Vaes. *Unique Cartan decomposition for II_1 factors arising from arbitrary actions of free groups*. Acta Mathematica **212**(1) (2014), 141–198. <https://doi.org/10.1007/s11511-014-0110-9>.
- [43] Sorin Popa and Stefaan Vaes. *Vanishing of the first continuous L^2 -cohomology for II_1 factors*. International Mathematics Research Notices **2015**(12) (2015), 3899–3907. <https://doi.org/10.1093/imrn/rnu054>.
- [44] Florin Rădulescu. *Random matrices, amalgamated free products and subfactors of the von Neumann algebra of a free group, of noninteger index*. Inventiones Mathematicae **115** (1994), 347–389. <https://doi.org/10.1007/BF01231764>.
- [45] Florin Rădulescu. *A one-parameter group of automorphisms of $L(F_\infty) \otimes B(H)$ scaling the trace*. Comptes Rendus de l'Académie des Sciences de Paris, Série I, Mathématique **314**(13) (1992), 1027–1032. <https://art.torvergata.it/handle/2108/121502>.
- [46] John Robert Ringrose. *Automatic Continuity of Derivations of Operator Algebras*. Journal of the London Mathematical Society, Second Series **5**(3) (1972), 432–438. <https://doi.org/10.1112/jlms/s2-5.3.432>.
- [47] Christopher Schafhauser, Aaron Tikuisis, and Stuart White. *Nuclear C^* -algebras: 99 problems*. arXiv preprint arXiv:2506.10902 (2025). <https://arxiv.org/abs/2506.10902>.
- [48] Junhao Shen. *Type II_1 factors with a single generator*. Journal of Operator Theory **62**(2) (2009), 421–438. <https://jot.theta.ro/jot/archive/2009-062-002/2009-062-002-010.pdf>.
- [49] Erling Størmer. *Entropy of some automorphisms of the II_1 -factor of the free group in infinite number of generators*. Inventiones Mathematicae **110** (1992), 63–73. <https://doi.org/10.1007/BF01231324>.
- [50] Andreas Thom. *L^2 -cohomology for von Neumann algebras*. Geometric and Functional Analysis **18**(1) (2008), 251–270. <https://doi.org/10.1007/s00039-007-0634-7>.
- [51] Dan-Virgil Voiculescu. *Some results on norm-ideal perturbations of Hilbert space operators*. Journal of Operator Theory **2**(1) (1979), 3–37. <https://jot.theta.ro/jot/archive/1979-002-001/1979-002-001-001.html>.
- [52] Dan-Virgil Voiculescu. *Cyclomorphy*. International Mathematics Research Notices **2002**(6) (2002), 299–332. Preprint version: arXiv:math/0105096v1, <https://arxiv.org/abs/math/0105096v1>. <https://doi.org/10.1155/S1073792802105046>.
- [53] Dan-Virgil Voiculescu. *The coalgebra of the free difference quotient and free probability*. International Mathematics Research Notices **2000**(2) (2000), 79–106. <https://doi.org/10.1155/S1073792800000064>.
- [54] Dan-Virgil Voiculescu. *A non-commutative Weyl-von Neumann theorem*. Revue Roumaine de Mathématiques Pures et Appliquées **21** (1976), 97–113. <https://math.berkeley.edu/~dvv/Publications.pdf>.