

RESEARCH ARTICLE

View Article Online
View Journal

Cite this: DOI: 10.1039/c8qo00191j

Domino cyclization/trifluoromethylation of 2-alkynylanilines using fluoroform-derived CuCF_3 : synthesis of 3-(trifluoromethyl)indoles†

Yibin Ye,  Kelvin Pak Shing Cheung, Lisi He and Gavin Chit Tsui  *

Received 22nd February 2018,

Accepted 14th March 2018

DOI: 10.1039/c8qo00191j

rsc.li/frontiers-organic


By employing easily accessible 2-alkynylanilines and the well-established fluoroform-derived CuCF_3 reagent, a novel class of 3-(trifluoromethyl)indoles can be synthesized in good yields with no ambiguity of the CF_3 position. The method utilizes a domino cyclization/trifluoromethylation strategy to construct the indole cores with the concomitant installation of the CF_3 group. The ultimate source of CF_3 is the low-cost industrial by-product fluoroform (CF_3H).

Trifluoromethylated heterocycles have found substantial applications in pharmaceuticals and agrochemicals.^{1,2} A number of important marketed drugs, including anti-HIV efavirenz (Sustiva), anti-inflammatory celecoxib (Celebrex) and anti-diabetic sitagliptin (Januvia), contain the trifluoromethylated heterocyclic core. Indoles constitute some of the most abundant and biologically active natural products and pharmaceutical compounds.³ The increasing interest in fluorinated indole derivatives towards drug discovery led to the recent development of methods accessing *trifluoromethylated indoles*, a class of trifluoromethylated heterocycles showing promising therapeutic potential.⁴

The construction of the substituted indole cores can be facilitated by the intramolecular cyclization of 2-alkynylanilines with the incorporation of functional groups at the C-3 position.⁵ This conceivably convenient approach, however, has not been exploited for the preparation of 3-(trifluoromethyl)indoles. To the best of our knowledge, there has been only one such example from Hou's group using stoichiometric amounts of CuBr as a promoter and the expensive Umemoto's reagent as an electrophilic CF_3 source.⁶ The majority of the current methods for the synthesis of 3-(trifluoromethyl)indoles rely on the trifluoromethylation of pre-formed indole cores or the use of CF_3 -containing building blocks. The direct C–H trifluoromethylation of indole cores provided powerful late-stage functionalization; however, mixtures of inseparable C-2 and C-3 trifluoromethylated products were often obtained.⁷ The

regioselectivity issues could be partially solved by using blocking groups^{7h-m} and pre-installed functionalities (halides or boronic acids),⁸ although the substrate scopes were limited and the reactions were less atom-economical. On the other hand, CF_3 -containing building blocks (e.g. alkenes and alkynes) have been used in cycloadditions and cyclizations to construct the indole cores.⁹ In most cases, tedious synthetic routes were needed to prepare the CF_3 -containing substrates, thus lowering the overall efficiency.

In the context of our interest in synthesizing novel trifluoromethylated heterocycles,¹⁰ we discovered that the 2-alkynylaniline **1a** can be converted into 3-(trifluoromethyl)indole **2a** in one step (82% isolated yield) using the *fluoroform-derived* $[\text{CuCF}_3]$ reagent (Scheme 1). Fluoroform (CF_3H) is a large-volume by-product from Teflon manufacturing and is commercially available at a very low cost. It is also non-toxic and ozone-friendly, and therefore, would be a highly attractive CF_3 source for preparing valuable trifluoromethylated compounds.¹¹ Grushin's group pioneered the preparation and applications of fluoroform-derived $[\text{CuCF}_3]$.¹² We have also successfully employed this reagent in the trifluoromethylation



Scheme 1 (a) Preparation of fluoroform-derived $[\text{CuCF}_3]$ by Grushin's method. (b) Synthesis of 3-(trifluoromethyl)indole **2a** from 2-alkynylaniline **1a**.

Department of Chemistry, The Chinese University of Hong Kong, Shatin, New Territories, Hong Kong SAR. E-mail: gctsui@cuhk.edu.hk

†Electronic supplementary information (ESI) available: Experimental procedures, characterization data and NMR spectra. CCDC 1551421. For ESI and crystallographic data in CIF or other electronic format see DOI: 10.1039/c8qo00191j

of alkynes, alkenes and arynes.¹³ $[\text{CuCF}_3]$ was prepared from copper(I) chloride/potassium *tert*-butoxide/fluoroform in DMF and stabilized with triethylamine trihydrofluoride ($\text{Et}_3\text{N}\cdot 3\text{HF}$) according to Grushin's procedure with modifications (Scheme 1a).^{12a,13a} Using this reagent under standard ambient conditions, the 3-(trifluoromethyl)indole core was furnished smoothly *via* a domino¹⁴ cyclization/trifluoromethylation sequence (Scheme 1b). Other known conditions for copper-mediated trifluoromethylations were also investigated (Table S1†),¹⁵ including the use of the $(\text{PPh}_3)_3\text{CuCF}_3$ complex,^{16a} electrophilic Togni's reagent,^{16b} and nucleophilic TMSCF_3 ,^{16c} which all showed inferior reactivities compared to fluoroform-derived $[\text{CuCF}_3]$ mainly due to the competing background cyclization to form the 3-*H* indole side product.¹⁷

The *N*-protecting group and reaction temperature had profound impacts on the reactivity (Table 1). The secondary sulfonamides were the most reactive substrates. Carbamates, amides, mono-/dialkylamines and unprotected aniline were completely unreactive (Table S2†).¹⁵ While substrate **1a** cleanly afforded the desired product **2a** in a good yield at room temperature (*cf.* Scheme 1b), it gave a small amount of desulfonylated 3-(trifluoromethyl)indole **5** at 80 °C (Table 1, entry 1).¹⁸ Substrates bearing mesyl (Ms) **3a**, nosyl (Ns) **3b** and triflyl (Tf) **3c** groups gave a mixture of products **4a–c** and **5**, even at room temperature (Table 1, entries 2–4). At 80 °C, the desulfonylated product predominated and by using substrate **3a** we could isolate compound **5** in 62% yield (Table 1, entry 2).¹⁹

The scope of the reaction was subsequently investigated using *N*-Ts 2-alkynylanilines **1b–x** (Table 2). The starting materials were conveniently synthesized in two steps from 2-iodoanilines *via* Sonogashira coupling of terminal alkynes and then *N*-tosylation with good yields. Upon treatment with fluoroform-derived $[\text{CuCF}_3]$, the domino cyclization/trifluoromethylation took place furnishing a novel class of diversely functionalized 3-(trifluoromethyl)indoles **2b–x**. The substituent groups were well tolerated at the acetylenic and aniline ring positions of the substrate. For R^1 , electron-rich arenes

Table 2 Scope of 3-(trifluoromethyl)indoles **2**^a

Table 2 displays the scope of the reaction, showing various 3-(trifluoromethyl)indoles (**2**) synthesized from 2-alkynylanilines (**1**) under the conditions: $[\text{CuCF}_3]$ (from fluoroform), DMF, 23 °C, 15 h, open to air.

Substrates **1** are shown with substituents R^1 and R^2 . The products **2** are shown with their respective yields.

Substrates **1** and corresponding products **2** (Yields):

- 1a** (X = H) → **2a** (85%^b)
- 1b** (X = 4-OMe) → **2b** (94%)
- 1c** (X = 3-OMe) → **2c** (90%)
- 1d** (X = 2-OMe) → **2d** (93%)
- 1e** (X = 4-Me) → **2e** (91%)
- 1f** (X = 3-Me) → **2f** (82%)
- 1g** (X = 2-Me) → **2g** (90%)
- 1h** (X = 4-CF₃) → **2h** (80%)
- 1i** (X = 4-CO₂Et) → **2i** (76%)
- 1j** (X = 4-NO₂) → **2j** (77%)
- 1k** (X = 4-Ac) → **2k** (78%)
- 1l** (X = 2-phenyl) → **2l** (75%)
- 1m** (X = 2-naphthyl) → **2m** (77%)
- 1n** (X = 2-pyridyl) → **2n** (63%)
- 1o** (X = 2-thienyl) → **2o** (68%)
- 1p** (X = 2-pyridyl) → **2p** (71%)
- 1q** (X = 2-thienyl) → **2q** (70%)
- 1r** (X = 2-thienyl) → **2r** (85%)
- 1s** (X = 2-phenyl) → **2s** (80%)
- 1t** (X = 2-phenyl) → **2t** (89%)
- 1u** (X = 2-phenyl) → **2u** (70%)
- 1v** (X = 2-phenyl) → **2v** (81%)
- 1w** (X = 2-phenyl) → **2w** (65%)
- 1x** (X = 2-phenyl) → **2x** (73%)
- 1y** (X = 2-phenyl) → **2y** (79%)

Structure of **2z** (Bazedoxifene) is shown, labeled as (prevention of postmenopausal osteoporosis). Its X-ray structure is also displayed.

^a Conditions: $[\text{CuCF}_3]$ in DMF solution (2.0 equiv., prepared from $\text{CuCl}/t\text{-BuOK}/\text{fluoroform}$ and stabilized by $\text{Et}_3\text{N}\cdot 3\text{HF}$) and **1** (0.2 mmol, 0.2 M in DMF). Isolated yields. ^b 1 mmol scale.

Table 1 Effects of the *N*-protecting group and reaction temperature^a

Table 1 displays the effects of the *N*-protecting group and reaction temperature on the reaction of 2-alkynylanilines (**1a**, **3a–c**) with $[\text{CuCF}_3]$ (from fluoroform) in DMF.

Substrates **1a**, **3a–c** are shown with substituents R (Ts, Ms, Ns, Tf). The products are 3-(trifluoromethyl)indoles (**2a**, **4a–c**) and 3-(trifluoromethyl)indole (**5**).

Yields are shown in parentheses.


Entry	R	Temp = 23 °C Yield ^b (%)	Temp = 80 °C Yield ^b (%)
1	Ts 1a	83(82)/0 2a/5	72/13 2a/5
2	Ms 3a	55(54)/17 4a/5	0/64(62) 4a/5
3	Ns 3b	60(60)/20 4b/5	11/63 4b/5
4	Tf 3c	50(53)/5 4c/5	0/62 4c/5

^a Conditions: $[\text{CuCF}_3]$ in DMF solution (2.0 equiv., prepared from $\text{CuCl}/t\text{-BuOK}/\text{fluoroform}$ and stabilized by $\text{Et}_3\text{N}\cdot 3\text{HF}$), **1a** or **3a–c** (0.2 mmol, 0.2 M in DMF). $\text{R} = \text{Ts}$ (*p*-toluenesulfonyl), Ms (methanesulfonyl), Ns (4-nitrobenzenesulfonyl), Tf (trifluoromethanesulfonyl). ^b Yield was determined by ^{19}F NMR analysis using benzotrifluoride as the internal standard. Isolated yields are shown in parentheses.


(**2b–g**) were generally higher-yielding than the electron-deficient ones (**2h–k**). The substituents at the *para*, *meta* and *ortho* positions of the benzene ring all gave good yields (**2b–g**). Other groups such as naphthyl (**2l**), phenanthryl (**2m**), pyridyl (**2p–q**) and thienyl (**2r**) were also demonstrated. For the alkyl substituent groups (**2n–o**), the yields were lower due to the increased amounts of background cyclization products. When R^1 was CO_2Me , only a trace amount of the desired product was detected and mainly the starting material was recovered (81%). The substrate containing terminal alkyne gave a complex mixture of products. On the aniline ring, methyl, fluoro and even chloro substituents (R^2) were tolerated at the C-5/6 positions (**2s–w**), despite the fact that aryl chlorides were known to react with fluoroform-derived $[\text{CuCF}_3]$ in aromatic trifluoromethylation reactions.^{12e} Furthermore, the 3- CF_3 -7-azaindole **2x** was synthesized in 73% yield, which could be an interesting candidate for medicinal chemistry owing to the pharmaceutical potential of 7-azaindoles.²⁰

Using this method, we could construct the 3- CF_3 indole core **2y** in 79% yield. X-ray crystallography unambiguously confirmed its structure and the C-3 position of the CF_3 group.²¹ The FDA approved drug bazedoxifene for the management of postmenopausal osteoporosis contains the analogous indole core.²² Therefore, compound **2y** could serve as a key intermediate towards the 3- CF_3 analog of bazedoxifene in a few steps according to literature procedures (detosylation, installing the azepane fragment and debenzoylation)^{22a} for the structure-activity relationship (SAR) studies.

The control experiment showed that when fluoroform-derived $[\text{CuCF}_3]$ was prepared under argon and used in the reaction under an argon atmosphere, no desired product **2a** was obtained (eqn (1), cf. Scheme 1). Instead, the background cyclization product **2a'** was observed exclusively. To probe the deprotection process, the *N*-Ts **2a** and *N*-Ms **4a** indole products were resubjected to the standard conditions at 80 °C (eqn (2)). It was found that fluoroform-derived $[\text{CuCF}_3]$ facilitated desulfonylation to give *N*-H product **5**, which was significantly more efficient for **4a** than **2a**, consistent with previous observations (cf. Table 1, entries 1–2). NMR studies had shown the intermediacy of **4a** in the formation of **5** from **3a** (Fig. S1†).¹⁵



Based on known literature evidence and the above studies, we propose a plausible mechanism for the domino cyclization/trifluoromethylation of 2-alkynylanilines **1** to form 3-(trifluoromethyl)indoles **2** (Scheme 2). The initial fluoroform-derived $[\text{Cu(I)CF}_3]$ reagent is easily oxidized to the $[\text{Cu(II)CF}_3]$ complex in air.^{12a,b,13a} This *pre-oxidation* step during the preparation of a CuCF_3 reagent is important as $[\text{Cu(I)CF}_3]$ was ineffective in the formation of the desired product **2a** (cf. eqn (1)). The activation of the alkyne moiety by coordination with $[\text{Cu(II)CF}_3]$, followed by 5-*endo-dig* cyclization furnish the indole core **B**.^{17a–c,10a} The unproductive protodemetalation of **B** would lead to 3-*H* by-product **2'**; this pathway was particularly dominant with other copper/ CF_3 sources and $[\text{Cu(I)CF}_3]$.¹⁵ Alternatively, removing a proton by using a base, *i.e.* KF and Et_3N are present in the reaction mixture from the CuCF_3 reagent,^{12a} gives the 3-copper indole intermediate **C**. The final reductive elimination delivers the 3- CF_3 indole product **2**.^{12b} Although Cu(II) complexes are invoked here for intermediates **B** and **C**, the involvement of Cu(III) and polynuclear species should not be completely ruled out, especially under aerobic conditions.²³ It is noteworthy that the nature of the fluoroform-derived $[\text{CuCF}_3]$ reagent, *i.e.* Cu(I) versus Cu(II) , is much more important than the requirement of air for the reaction.



Scheme 2 Proposed mechanism.

The control experiment showed that using $[\text{Cu(II)CF}_3]$ and carrying out the reaction with **1a** under an argon atmosphere also gave 74% yield (NMR) of **2a**. In general, running the reactions open to air was more convenient and gave higher yields.

In conclusion, a useful synthetic method has been developed for the preparation of trifluoromethylated indoles with the following features: (1) 3-(trifluoromethyl)indoles, mostly new compounds, can be accessed from the common starting material 2-alkynylanilines, with no ambiguity of the CF_3 position; (2) reaction conditions are mild, not moisture or air-sensitive, and tolerate a wide range of functional groups; (3) energy-efficient domino strategies are employed to form multiple bonds in one pot without the isolation and purification of intermediates; and (4) the ultimate CF_3 source is the industrial by-product fluoroform (CF_3H).

Conflicts of interest

There are no conflicts to declare.

Acknowledgements


This work was supported by the Research Grants Council of Hong Kong (CUHK 24301217) and the Chinese University of Hong Kong (the Faculty Strategic Fund for Research from the Faculty of Science and the Direct Grant for Research 4053276).

Notes and references

- For reviews on trifluoromethylated heterocycles, see: (a) V. Muzalevskiy, A. Shastin, E. Balenkova, G. Haufe and V. Nenajdenko, *Synthesis*, 2009, 3905; (b) A. A. Gakh and Y. Shermolovich, *Curr. Top. Med. Chem.*, 2014, **14**, 952; (c) *Fluorine in Heterocyclic Chemistry*, ed. V. Nenajdenko,

- Springer International Publishing, 2014; (d) V. A. Petrov, *Fluorinated Heterocyclic Compounds: Synthesis, Chemistry, and Applications*, John Wiley & Sons, New Jersey, 2009.
- 2 For selected recent reviews on fluorination and trifluoromethylation methods, see: (a) S. Krishnamoorthy and G. Prakash, *Synthesis*, 2017, 3394; (b) C. Alonso, E. Martínez de Marigorta, G. Rubiales and F. Palacios, *Chem. Rev.*, 2015, **115**, 1847; (c) X. Yang, T. Wu, R. J. Phipps and F. D. Toste, *Chem. Rev.*, 2015, **115**, 826; (d) C. N. Neumann and T. Ritter, *Angew. Chem., Int. Ed.*, 2015, **54**, 3216.
 - 3 (a) *Indoles*, ed. R. J. Sundberg, Academic Press, London, 1996; (b) J. A. Joule and K. Mills, *Heterocyclic Chemistry*, Blackwell Publishing Ltd, 2010; (c) S. M. Bronner, A. E. Goetz and N. K. Garg, *Synlett*, 2011, 2599; (d) R. Vallakati and J. A. May, *Synlett*, 2012, 2577; (e) I. Chakraborty and S. Jana, *Synthesis*, 2013, 3325.
 - 4 (a) M. Bandini, A. Melloni and A. Umani-Ronchi, *Angew. Chem., Int. Ed.*, 2004, **43**, 550; (b) T. B. Poulsen and K. A. Jorgensen, *Chem. Rev.*, 2008, **108**, 2903; (c) M. Bandini and A. Eichholzer, *Angew. Chem., Int. Ed.*, 2009, **48**, 9608; (d) M. Zeng and S.-L. You, *Synlett*, 2010, 1289.
 - 5 For reviews, see: (a) G. Battistuzzi, S. Cacchi and G. Fabrizi, *Eur. J. Org. Chem.*, 2002, 2671; (b) K. Krüger, A. Tillack and M. Beller, *Adv. Synth. Catal.*, 2008, **350**, 2153. For selected examples, see: (c) J. Barluenga, M. Trincado, E. Rubio and J. M. González, *Angew. Chem., Int. Ed.*, 2003, **42**, 2406; (d) Y. Yin, W. Ma, Z. Chai and G. Zhao, *J. Org. Chem.*, 2007, **72**, 5731; (e) X. Han and X. Lu, *Org. Lett.*, 2010, **12**, 3336; (f) K. Inamoto, N. Asano, Y. Nakamura, M. Yonemoto and Y. Kondo, *Org. Lett.*, 2012, **14**, 2622; (g) S. Li and S. Ma, *Adv. Synth. Catal.*, 2012, **354**, 2387; (h) A. Arcadi, E. Pietropaolo, A. Alvino and V. Michelet, *Org. Lett.*, 2013, **15**, 2766; (i) J. Liu, X. Xie and Y. Liu, *Chem. Commun.*, 2013, **49**, 11794; (j) L. Yang, Y. Ma, F. Song and J. You, *Chem. Commun.*, 2014, **50**, 3024; (k) J. Yu, D. Zhang-Negrerie and Y. Du, *Org. Lett.*, 2016, **18**, 3322; (l) Q. Shi, P. Li, Y. Zhang and L. Wang, *Org. Chem. Front.*, 2017, **4**, 1322.
 - 6 G. Ge, X. Huang, C. Ding, H. Li, S. Wan and X. Hou, *Chin. J. Chem.*, 2014, **32**, 727.
 - 7 (a) P. G. Janson, N. O. Ilchenko, A. Diez-Varga and K. J. Szabó, *Tetrahedron*, 2015, **71**, 922; (b) R. Rey-Rodriguez, P. Retailleau, P. Bonnet and I. Gillaizeau, *Chem. – Eur. J.*, 2015, **21**, 3572; (c) E. Mejía and A. Togni, *ACS Catal.*, 2012, **2**, 521; (d) L. Chu and F.-L. Qing, *J. Am. Chem. Soc.*, 2012, **134**, 1298; (e) N. Iqbal, S. Choi, E. Ko and E. J. Cho, *Tetrahedron Lett.*, 2012, **53**, 2005; (f) D. A. Nagib and D. W. C. MacMillan, *Nature*, 2011, **480**, 224; (g) X. Mu, S. Chen, X. Zhen and G. Liu, *Chem. – Eur. J.*, 2011, **17**, 6039; (h) K. Natte, R. V. Jagadeesh, L. He, J. Rabeah, J. Chen, C. Taeschler, S. Ellinger, F. Zaragoza, H. Neumann, A. Brückner and M. Beller, *Angew. Chem., Int. Ed.*, 2016, **55**, 2782; (i) A. G. O'Brien, A. Maruyama, Y. Inokuma, M. Fujita, P. S. Baran and D. G. Blackmond, *Angew. Chem., Int. Ed.*, 2014, **53**, 11868; (j) S. P. Pitre, C. D. McTiernan, H. Ismaili and J. C. Scaiano, *ACS Catal.*, 2014, **4**, 2530; (k) C. Xu, J. Liu, W. Ming, Y. Liu, J. Liu, M. Wang and Q. Liu, *Chem. – Eur. J.*, 2013, **19**, 9104; (l) S. Seo, J. B. Taylor and M. F. Greaney, *Chem. Commun.*, 2013, **49**, 6385; (m) Y. Ji, T. Brueckl, R. D. Baxter, Y. Fujiwara, I. B. Seiple, S. Su, D. G. Blackmond and P. S. Baran, *Proc. Natl. Acad. Sci. U. S. A.*, 2011, **108**, 14411.
 - 8 (a) X. Zhang, J. Wang and Z. Wan, *Org. Lett.*, 2015, **17**, 2086; (b) Z. Gonda, S. Kovács, C. Wéber, T. Gáti, A. Mészáros, A. Kotschy and Z. Novák, *Org. Lett.*, 2014, **16**, 4268; (c) T. D. Senecal, A. T. Parsons and S. L. Buchwald, *J. Org. Chem.*, 2011, **76**, 1174; (d) T. Liu, X. Shao, Y. Wu and Q. Shen, *Angew. Chem., Int. Ed.*, 2011, **51**, 540; (e) C.-P. Zhang, Z.-L. Wang, Q.-Y. Chen, C.-T. Zhang, Y.-C. Gu and J.-C. Xiao, *Angew. Chem., Int. Ed.*, 2011, **50**, 1896.
 - 9 (a) Z. Xia, K. Wang, J. Zheng, Z. Ma, Z. Jiang, X. Wang and X. Lv, *Org. Biomol. Chem.*, 2012, **10**, 1602; (b) Z.-J. Wang, J.-G. Yang, F. Yang and W. Bao, *Org. Lett.*, 2010, **12**, 3034; (c) T. Konno, J. Chae, T. Ishihara and H. Yamanaka, *J. Org. Chem.*, 2004, **69**, 8258; (d) J. Ichikawa, Y. Iwai, R. Nadano, T. Mori and M. Ikeda, *Chem. – Asian J.*, 2008, **3**, 393; (e) Y.-Q. Fang and M. Lautens, *Org. Lett.*, 2005, **7**, 3549; (f) J. Chae, T. Konno, T. Ishihara and H. Yamanaka, *Chem. Lett.*, 2004, **33**, 314.
 - 10 (a) Q. Wang, L. He, K. K. Li and G. C. Tsui, *Org. Lett.*, 2017, **19**, 658; (b) K. P. S. Cheung and G. C. Tsui, *Org. Lett.*, 2017, **19**, 2881; (c) Q. Wang and G. C. Tsui, *J. Org. Chem.*, 2018, **83**, 2971; (d) Y. Ye, K. P. S. Cheung, L. He and G. C. Tsui, *Org. Lett.*, 2018, **20**, 1676.
 - 11 For reviews, see: (a) V. V. Grushin, *Chim. Oggi Chem. Today*, 2014, **32**, 81; (b) S. Kyasa, *Synlett*, 2015, 1911. For recent examples of using fluoroform in synthesis, see: (c) J. B. Geri and N. K. Szymczak, *J. Am. Chem. Soc.*, 2017, **139**, 9811; (d) E. Carbonnel, T. Besset, T. Poisson, D. Labar, X. Pannecoucke and P. Jubault, *Chem. Commun.*, 2017, **53**, 5706; (e) M. B. Haskali and V. W. Pike, *Chem. – Eur. J.*, 2017, **23**, 8156; (f) K. Aikawa, K. Maruyama, J. Nitta, R. Hashimoto and K. Mikami, *Org. Lett.*, 2016, **18**, 3354; (g) S. Okusu, E. Tokunaga and N. Shibata, *Org. Lett.*, 2015, **17**, 3802; (h) S. Potash and S. Rozen, *J. Org. Chem.*, 2014, **79**, 11205; (i) C. S. Thomason and W. R. Dolbier Jr., *J. Org. Chem.*, 2013, **78**, 8904; (j) G. K. S. Prakash and G. A. Olah, *Science*, 2012, **338**, 1324; (k) S. Potash and S. Rozen, *J. Fluorine Chem.*, 2014, **168**, 173.
 - 12 For seminal reports, see: (a) A. Zanardi, M. A. Novikov, E. Martin, J. Benet-Buchholz and V. V. Grushin, *J. Am. Chem. Soc.*, 2011, **133**, 20901; (b) P. Novák, A. Lishchynskiy and V. V. Grushin, *Angew. Chem., Int. Ed.*, 2012, **51**, 7767; (c) P. Novák, A. Lishchynskiy and V. V. Grushin, *J. Am. Chem. Soc.*, 2012, **134**, 16167; (d) A. I. Konovalov, J. Benet-Buchholz, E. Martin and V. V. Grushin, *Angew. Chem., Int. Ed.*, 2013, **52**, 11637; (e) A. Lishchynskiy, M. A. Novikov, E. Martin, E. C. Escudero-Adán, P. Novák and V. V. Grushin, *J. Org. Chem.*, 2013, **78**, 11126.
 - 13 (a) L. He and G. C. Tsui, *Org. Lett.*, 2016, **18**, 2800; (b) L. He, X. Yang and G. C. Tsui, *J. Org. Chem.*, 2017, **82**,

- 6192; (c) X. Yang, L. He and G. C. Tsui, *Org. Lett.*, 2017, **19**, 2446; (d) X. Yang and G. C. Tsui, *Org. Lett.*, 2018, **20**, 1179.
- 14 *Domino Reactions*, ed. L. F. Tietze, Wiley-VCH, Weinheim, 2014.
- 15 See the ESI† for full details.
- 16 (a) O. A. Tomashenko, E. C. Escudero-Adán, M. M. Belmonte and V. V. Grushin, *Angew. Chem., Int. Ed.*, 2011, **50**, 7655; (b) J.-S. Lin, Y.-P. Xiong, C.-L. Ma, L.-J. Zhao, B. Tan and X.-Y. Liu, *Chem. – Eur. J.*, 2014, **20**, 1332; (c) L. Chu and F.-L. Qing, *J. Am. Chem. Soc.*, 2010, **132**, 7262.
- 17 Substrate **1a** was prone to background cyclization in the presence of copper salts. The previously reported conditions for the synthesis of trifluoromethylated pyrazoles using $\text{Cu}(\text{OTf})_2/\text{TMSCF}_3/\text{KF}$ (ref. 10a) only gave 34% of **2a** with significant background cyclization. Lowering the equivalents of fluoroform-derived $[\text{CuCF}_3]$ led to yield decrease (1.0 equiv., 64% of **2a**). See the ESI† for optimization studies. For examples of $\text{Cu}(\text{i})/(\text{ii})$ -mediated cyclization of 2-alkynylanilines, see: (a) K. Hiroya, S. Itoh, M. Ozawa, Y. Kanamori and T. Sakamoto, *Tetrahedron Lett.*, 2002, **43**, 1277; (b) Z. Shen and X. Lu, *Adv. Synth. Catal.*, 2009, **351**, 3107; (c) J. Yu, D. Zhang-Negrerie and Y. Du, *Org. Lett.*, 2016, **18**, 3322.
- 18 We have also reported a $\text{Cu}(\text{OTf})_2$ -promoted domino cyclization/trifluoromethylation/desulfonylation of tosylhydrazones using TMSCF_3 , see ref. 10a.
- 19 The *N*-Ms 2-alkynylanilines were useful for providing the *N*-H 3-(trifluoromethyl)indoles. The reactions tolerated electron-donating/-withdrawing substituents:



- 20 F. Popowycz, S. Routier, B. Joseph and J.-Y. M  rour, *Tetrahedron*, 2007, **63**, 1031.
- 21 CCDC 1551421† (**2y**) contains the supplementary crystallographic data for this paper. Note the distortion of the CF_3 group.
- 22 (a) C. P. Miller, M. D. Collini, B. D. Tran, H. A. Harris, Y. P. Kharode, J. T. Marzolf, R. A. Moran, R. A. Henderson, R. H. W. Bender, R. J. Unwalla, L. M. Greenberger, J. P. Yardley, M. A. Abou-Gharbia, C. R. Lyttle and B. S. Komm, *J. Med. Chem.*, 2001, **44**, 1654; (b) B. S. Komm, Y. P. Kharode, P. V. N. Bodine, H. A. Harris, C. P. Miller and C. R. Lyttle, *Endocrinology*, 2005, **146**, 3999.
- 23 (a) J. X. Qiao and P. Y. S. Lam, *Synthesis*, 2011, 829; (b) A. E. King, T. C. Brunold and S. S. Stahl, *J. Am. Chem. Soc.*, 2009, **131**, 5044; (c) M. Tromp, G. P. F. van Strijdonck, S. S. van Berkel, A. van den Hoogenband, M. C. Feiters, B. de Bruin, S. G. Fiddy, A. M. J. van der Eerden, J. A. van Bokhoven, P. W. N. M. van Leeuwen and D. C. Koningsberger, *Organometallics*, 2010, **29**, 3085.