

## Dilution & Storage Protocol

Hair was placed in 20  $\mu$ L of Dilution Buffer containing 0.5  $\mu$ L of DNA Release Additive. The samples were incubated at room temperature for 2 minutes and then at 98  $^{\circ}$ C for 5 minutes. The samples were spun down and 1  $\mu$ L of the supernatant was used as a template in a 20  $\mu$ L reaction with Phire Tissue Direct PCR Master Mix and gene specific primers (Table 1). PCR was performed as described in Table 2 and PCR products were loaded directly onto gels.

Table 1. PCR Reaction Setup.

Component	20 $\mu$ L rxn	50 $\mu$ L rxn	Final conc.
H <sub>2</sub> O	add to 20 $\mu$ L	add to 50 $\mu$ L	
2X Phire Tissue Direct PCR Master Mix	10 $\mu$ L	25 $\mu$ L	1X
Primer A	X $\mu$ L	X $\mu$ L	0.5 $\mu$ M
Primer B	X $\mu$ L	X $\mu$ L	0.5 $\mu$ M
<b>Sample</b>			
Direct Protocol	–	0.5 mm punch of hair	0.5 mm/L
Dilution & Storage Protocol	1 $\mu$ L	–	

Table 2. PCR Cycling Conditions.

Cycle step	2-step		3-step		Cycles
	Temp.	Time	Temp.	Time	
Initial denaturation	98 $^{\circ}$ C	5 min	98 $^{\circ}$ C	5 min	1
Denaturation	98 $^{\circ}$ C	5 s	98 $^{\circ}$ C	5 s	35
Annealing	–	–	X $^{\circ}$ C	5 s	
Extension	72 $^{\circ}$ C	20 s $\leq$ 1 kb 20 s/kb $>$ 1 kb	72 $^{\circ}$ C	20 s $\leq$ 1 kb 20 s/kb $>$ 1 kb	
Final Extension	72 $^{\circ}$ C +4 $^{\circ}$ C	1 min hold	72 $^{\circ}$ C +4 $^{\circ}$ C	1 min hold	1

## Results and Discussion

The ability of Phire Tissue Direct PCR Master Mix to amplify target DNA from the hair samples was tested using mouse hair. Mouse hair samples were collected with tweezers from the animal's back, and only hair with follicles attached were used for PCR. Robust amplification of different length amplicons was achieved even from a single hair sample (Figure 1). Phire Tissue Direct PCR Master Mix can therefore be used for PCR based genotyping applications by using hair as a starting material.




Figure 1. Different length amplicons were amplified using different number of mice hair and Phire Tissue Direct PCR Master Mix (direct protocol). Fragments sizes 1 – 237 bp, 2 – 515 bp, 3 – 1.1 kb. M – Thermo Scientific™ O'GeneRuler™ Express DNA Ladder.

Multiplex PCR amplification of five different length fragments was achieved using mouse hair and Phire Tissue Direct PCR Master Mix (Figure 2.). This approach can be used in genotyping applications that require simultaneous determination of several markers in one reaction.




Figure 2. Five different length DNA amplicons, ranging from 185 to 650 bp, were amplified from mouse hair in 1-plex to 5-plex reaction with Phire Tissue Direct PCR Master Mix. Dilution & storage protocol and 5 hairs were used for the amplification. M – O'GeneRuler Express DNA Ladder.

The Phire DNA polymerase from Direct PCR Master Mix is optimized to achieve robust and sensitive amplification even from tiny amounts of tissue samples. The performance of Phire Direct PCR Master Mix was compared to that of corresponding kits from other vendors. Figure 3 shows that while other vendors' kits work

7. send sanger before 7 pm.