
a
The plasticity association is attenuated but survives
grey bar, median to max of 24 detection−matched control genes
AKSP
A
WT
0.0
0.1
0.2
0.3
partial Spearman rho with the lineage plasticity axis
as published
minus the 14 genes shared with mOnF
b
The placenta signal does not survive removing generic YAP targets
no contrast remains significant once 4 of 34 genes are dropped
YAPOE_FOS_vs_EV
YAPOE_vs_EV
YAPsh1_vs_scr
FOSkd_vs_scr
YAPsh2_vs_scr
FOSDN_vs_EV_in_Apcloss
0
2
4
mean Wald statistic (cameraPR)
placenta set, 34 genes
minus 4 YAP target genes (Egfr, Krt19, Krt8, Phlda2)
FDR < 0.05
FALSE
TRUE
c
Both immune results are carried by a single gene
the gene named in brackets is the one removed
Cytotoxic resistance
Immune suppression
FOSDN_vs_EV_in_Apcloss (Tnfaip3)
FOSkd_vs_scr (Serpinb9)
YAPOE_FOS_vs_EV (Serpinb9)
YAPOE_vs_EV (Serpinb9)
YAPsh1_vs_scr (Cd24a)
YAPsh2_vs_scr (Cflar)
FOSDN_vs_EV_in_Apcloss (Cd274)
FOSkd_vs_scr (H2−T23)
YAPOE_FOS_vs_EV (Nt5e)
YAPOE_vs_EV (Nt5e)
YAPsh1_vs_scr (Pvr)
YAPsh2_vs_scr (Pvr)
0.001
0.010
0.100
1.000
cameraPR P (log scale); dashed line 0.05
whole set
minus the strongest gene
d
The locus is a bound TEAD and AP−1 element in public ChIP−seq
ChIP−Atlas mm10, 5 kb window; but 0 of the 4 intestinal experiments calls a peak
Fosl1
Jund
Fosl2
Tead1
Fos
Junb
Jun
Tead4
Yap1
5/6 expts
26/34 expts
11/20 expts
19/24 expts
22/51 expts
15/58 expts
32/87 expts
12/32 expts
7/37 expts
50
75
100
percentile of the Tnfrsf12a TSS among all genes, by mean MACS2 score
AP−1
YAP/TEAD