
a
Tnfrsf12a responds to YAP and to AP−1, in both directions
3−4 organoid replicates per arm; batch in the model where present
YAP gain
YAPOE_vs_EV
YAP+AP−1 gain
YAPOE_FOS_vs_EV
AP−1 gain on YAP
YAPOE_FOS_vs_YAPOE
YAP loss
YAPsh1_vs_scr
YAPsh2_vs_scr
AP−1 loss
FOSkd_vs_scr
FOSDN_vs_EV_in_Apcloss
progression
AKSP_vs_WT
A_vs_WT
Apcloss_vs_WT
AKSP_vs_A
oncofetal state
OnFreporter_high_vs_neg
RXR gain
RXROE_vs_EV_WT
RXROE_vs_EV_Apcloss
RXR loss
RXRi_vs_DMSO_AKSP
−2
−1
0
1
2
3
Tnfrsf12a log2 fold change (DESeq2, 95% CI)
n.s.
padj < 0.05
b
The placenta and tumour−intrinsic immune programmes move with YAP and AP−1
cameraPR, competitive test; stars are global BH FDR (* .05, ** .01, *** .001)
oncofetal (authors')
revCSC
YAP regeneration
Lgr5 stem
placenta / trophoblast
EVT core
IFA core
TRA core
immune suppression
cytotoxic resistance
YAPOE_vs_EV
YAPOE_FOS_vs_EV
YAPOE_FOS_vs_YAPOE
YAPsh1_vs_scr
YAPsh2_vs_scr
FOSDN_vs_EV_in_Apcloss
FOSkd_vs_scr
mean Wald statistic
10
0
−10
c
Across 15 contrasts the receptor moves with the programme
Spearman rho = 0.975
Tnfrsf12a, Wald statistic
mOnF programme, mean Wald statistic
YAPOE_FOS_vs_EV
YAPOE_vs_EV
AKSP_vs_WT
A_vs_WT
Apcloss_vs_WT
OnFreporter_high_vs_neg
YAPOE_FOS_vs_YAPOE
RXRi_vs_DMSO_AKSP
AKSP_vs_A
RXROE_vs_EV_WT
RXROE_vs_EV_Apcloss
FOSkd_vs_scr
YAPsh1_vs_scr
YAPsh2_vs_scr
FOSDN_vs_EV_in_Apcloss