

a
AlphaFold model AF-Q9NP84-F1 (v6), mature chain 28–129
Illustrative orientation only: inter-domain geometry is not determined
Extracellular
TWEAK-binding residues
D45 · K48 · M50 · D62
Cysteine-rich domain
(CRD, TNFR-Cys) 36–67
3 disulfides: C36–C49,
C52–C67, C55–C64
Stalk 68–80
N-terminus (E28)
N-segment 28–35
Lipid
bilayer
Transmembrane helix
81–101 (21 aa)
C-terminus (Q129)*
Juxtamembrane basic
cluster 102–109
RRCRRREK
Cytoplasmic tail 110–129
PIEET 113–117 matches the
TRAF2 (P/S/A/T)x(Q/E)E motif
10 Å
Cytoplasm
Bilayer is drawn, not modelled: placed on the TM-helix (81–101) axis, hydrophobic core ±15 Å.
CRD placement relative to the bilayer is not determined (CRD↔TM mean PAE 23 Å; within-domain ≤ 4 Å).
*AlphaFold places 124–129 inside the bilayer (pLDDT < 50); not a physical position.
b
Experimental ECD: PDB 2KMZ (solution NMR, model 1 of 20)
N (E28)
D45
K48
M50
D62
C55–C64
C52–C67
C36–C49
P80
Same orientation as a
(CRD superposed).
CRD Cα RMSD vs AF (32 Cα):
1.74 Å to model 1;
1.30–2.14 Å over 20 models
(median 1.59 Å)
c
1
28
36
67
81
101
110
129
disulfides
SP
N-seg
CRD
Stalk
TM
JM
C-tail
TWEAK-binding residues
AF pLDDT
90
70
50
0
cleaved
pLDDT
<50
50–70
70–90
>90