
a
TNFRSF12A is enriched in regenerative and hybrid states
15 patients, 16,009 epithelial nuclei; REC and iREC are indistinguishable
Intermediate
iREC
REC
Hypoxia
TA2
TA1
Stem NOTUM
Goblet
Colonocyte
Stem
UPR
Enteroendocrine
Tuft
36%
30%
30%
25%
22%
21%
20%
16%
14%
13%
10%
10%
2%
0
10
20
30
40
% of nuclei with TNFRSF12A detected
hybrid (Intermediate)
other epithelial
regenerative (REC / iREC)
Stem NOTUM
b
Both links are real; the oncofetal link is the larger one
per patient, each dimension controlled for the other, cell cycle and depth
bar = median; 11 of 12 patients oncofetal > inflammatory, P = 0.0015
partial Spearman rho with TNFRSF12A
0.0
0.1
0.2
oncofetal
(revCSC)
inflammatory
(IFN−gamma)
c
Specific on the oncofetal axis; matched by a control gene on the other
vs 24 detection−matched control genes: 0 of 24 exceed the receptor on the oncofetal
axis, but 1 of 24 (CLTB) does on the inflammatory axis. Dotted line = equal association.
TNFRSF12A
partial rho, inflammatory axis
0.00
0.05
0.00
0.05
0.10
0.15
partial rho, oncofetal axis
d
Both axes add; neither explains the other
within−patient median split on each score. Thin lines, patients; thick lines, median.
9.4% −> 16.7% (IFN alone) −> 23.5% (OnF alone) −> 34.3% (both)
% of nuclei with TNFRSF12A detected
0
20
40
60
OnF−low
OnF−high
IFN−low
IFN−high