
a
The receptor tracks the oncofetal front; the ligand does not
GeoMx CTA, tumour epithelium, patient−paired (n = 10)
normalised expression (log)
MMP7
+1.05 | 10/10 | q = 0.004
LAMC2
+0.95 | 10/10 | q = 0.004
TNFRSF12A
+0.46 | 10/10 | q = 0.004
TGFB3
+0.18 | 9/10 | q = 0.006
TGFB1
+0.13 | 8/10 | q = 0.192
TNFSF12
+0.02 | 5/10 | q = 0.625
core
front
b
A ligand with sources, a ligand without, and one that is absent
cell types ordered by TGFB1 detection; early−CRC microenvironment scRNA−seq, 2,409 cells, 5 patients
TNFSF12 (TWEAK)
TGFB1
IL36G
telocyte−like CAF
trophocyte−like CAF MMP+
endothelial.tip
macrophage
trophocyte−like CAF MMP−
pericytes
telocyte
MKI67+ lymphocytes
mast cells
myofibroblast.1
T.reg
endothelial.stalk
T.CD8
T.CD4
trophocyte
plasma
SMC
enteric glial
stromal.1
B cell
myofibroblast.2
% of cells with >= 1 count
c
Where the ligand−positive cells are
top eight TNFSF12 sources, same order in both panels; macrophages supply 7.1% of TNFSF12+ cells
TNFSF12 (TWEAK)
TGFB1
pericytes
myofibroblast.1
telocyte
T.reg
T.CD8
macrophage
trophocyte
endothelial.stalk
% of all ligand−positive cells