Vector spaces

A set V with field F and operations $+$ and $\cdot$ such that $\forall u, v, w \in V, \forall a, b \in F$:

- $u + v \in V$
- $a \cdot u \in V$
- $u + v = v + u$
- $(u + v) + w = u + (v + w)$
- $\exists 0 \in V : u + 0 = u$
- $\exists (-u) \in V : u + (-u) = 0$
- $1 \cdot u = u$
- $(ab) \cdot u = a \cdot (b \cdot u)$
- $(a + b) \cdot u = a \cdot u + b \cdot u$
- $a \cdot (u + v) = a \cdot u + a \cdot v$

subspace a subset S of V that is itself a vector space under the same operations. It is sufficient that $S \neq \varnothing$ and $\forall u, v \in S \, \forall a, b \in F, au + bv \in S$.

linear combination for a subset S of V, a sum of scalar multiples of vectors in S, $\sum_{v \in S} a_v v$

linear independence the property of a subset S of V where $\sum_{v \in S} a_v v = 0 \implies \forall v \in S, a_v = 0$

span $\operatorname{Span}(S) := \left\{ \sum_{v \in S} a_v v \,\middle|\, \forall s \in S, a_s \in F \right\}$, the set of all linear combinations of vectors in S.

basis a subset B of V that is linearly independent and spans V.

dimension $\dim(V) = |B|$ for basis B.

sum of vector spaces for subspaces U and W of vector space V, $U + W := \{u + w \mid u \in U \wedge w \in W\}$

Theorems

- Zero vectors are unique for each vector space.
- Each vector has a unique additive inverse.
- Intersections of subspaces are subspaces.
- Spans are subspaces.
- Every vector has a unique representation w.r.t. a basis.
- Every basis of a vector space has the same cardinality.
- for a subset S of V, $|S| > \dim(V) \implies S$ is linearly dependent.
- for a subset S of V, the following imply S is a basis:
  - S is linearly independent and spans V
  - S is linearly independent with $|S| = \dim(V)$
  - S spans V with $|S| = \dim(V)$
- for subspaces W_1 and W_2, $\dim(W_1 + W_2) = \dim(W_1) + \dim(W_2) - \dim(W_1 \cap W_2)$

Standard bases

Linear maps

For vector spaces V and W over field F, a map $f : V \to W$ such that $\forall a, b \in F \, \forall u, v \in V = f(au + bv) = af(u) + bf(v)$.

kernel $\operatorname{Ker}(f) := \{v \mid v \in V \wedge f(v) = 0\}$ is the set of vectors in V that f takes to the zero vector in W. Also called the null space or notated $\operatorname{N}(f)$.

nullity $\operatorname{null}(f) := \dim(\operatorname{Ker}(f))$

image $\operatorname{Im}(f) := \{w \mid w \in W \wedge \exists v \in V : w = f(v)\}$ is the set of outputs of f in W. Also called the range or notated $\operatorname{R}(f)$.

rank $\operatorname{rank}(f) := \dim(\operatorname{Im}(f))$

Theorems

- $f(0_V) = 0_W$
- f is injective $\iff \operatorname{Ker}(f) = \{0\} \iff \operatorname{null}(f) = 0$
- f is surjective $\iff \operatorname{Im}(f) = W \iff \operatorname{rank}(f) = \dim(f)$