

# **Scientific Project**

Summer Semester 2026

Supervisor: Prof. Dr. Stefan Teufel  
Mathematics Department  
University of Tübingen

Fernando Muñoz Martínez

# Contents

|          |                                                                       |          |
|----------|-----------------------------------------------------------------------|----------|
| <b>1</b> | <b>Introduction</b>                                                   | <b>2</b> |
| 1.1      | Research Motivation . . . . .                                         | 2        |
| 1.2      | Problem Statement . . . . .                                           | 2        |
| <b>2</b> | <b>Theoretical Framework</b>                                          | <b>3</b> |
| 2.1      | Basic Definitions . . . . .                                           | 3        |
| 2.1.1    | Metric and open balls on $\mathbb{R}^n$ . . . . .                     | 3        |
| 2.1.2    | Uniformly Discrete Set . . . . .                                      | 3        |
| 2.1.3    | Discrete Operator and Matrix Elements . . . . .                       | 3        |
| 2.1.4    | Finite-Dimensional Subspaces and Orthogonal Projections . . . . .     | 3        |
| 2.1.5    | Short- and Finite-Range Operators . . . . .                           | 3        |
| 2.1.6    | Equivalent Action and Finite Local Complexity . . . . .               | 4        |
| 2.1.7    | Hausdorff Distance . . . . .                                          | 4        |
| 2.2      | The Spectral and Pseudospectral Problems . . . . .                    | 4        |
| 2.2.1    | Normal Operators . . . . .                                            | 4        |
| 2.2.2    | The Pseudospectrum . . . . .                                          | 4        |
| 2.3      | Approximating the (Pseudo-)Spectrum with Full Error Control . . . . . | 6        |
| 2.3.1    | The Pseudospectral Inclusion Bound . . . . .                          | 6        |
| 2.3.2    | The Spectral Gap Bound . . . . .                                      | 7        |
| 2.3.3    | A General Algorithm . . . . .                                         | 7        |
| <b>3</b> | <b>Conclusion and Future Work</b>                                     | <b>9</b> |

# 1 Introduction

## 1.1 Research Motivation

Spectral analysis has proven to be a powerful tool for solving a wide range of problems in mathematics, physics and engineering. Among its many applications, it facilitates the decoupling of problems involving vectors or functions into collections of simpler problems involving only scalars, the analysis of the phenomenon of resonance, and the characterization of the asymptotic behavior of a system [7]. Therefore, the ability to compute the spectra of linear operators is of fundamental importance.

In practice, however, analytical solutions are rarely available. Even in the case of finite-dimensional operators represented by  $n \times n$  matrices, the theorem of Abel and Ruffini shows that no general formula, involving only rational operations and root extractions, can exist for solving algebraic equations of degree higher than fourth [8]. As a result, there is no general closed-form expression for the eigenvalues of arbitrary matrices, making numerical methods necessary.

The situation becomes even more challenging when considering infinite-volume operators or those acting on infinite-dimensional spaces. Nevertheless, their ubiquity in applications has driven the development of multiple numerical algorithms designed to approximate the spectra for different classes of operators.

Motivated by these challenges, the authors of [4] address the problem of computing the spectra for *short-range finite-local-complexity infinite-volume operators*. Prior to their work, no numerical method was capable of approximating the spectra for this specific class of operators with full error control. By overcoming this limitation, their work fills an important gap in the existing literature.

## 1.2 Problem Statement

The broad aim of this work is to validate and contribute to the understanding of the algorithm developed in [4] in concrete situations, e.g., in the computation of spectral densities of quasicrystals. To this end, simulation tools which take advantage of extensively-tested scientific libraries such as NumPy and SciPy were developed in Python. This allowed us to fully focus on the implementation and validation of the algorithm while resting assured that everything else would work seamlessly.

## 2 Theoretical Framework

### 2.1 Basic Definitions

#### 2.1.1 Metric and open balls on $\mathbb{R}^n$

In this dissertation, unless otherwise stated, we define the distance  $d$  on  $\mathbb{R}^n$  as the *maximum distance*

$$d(x, y) = d((x_1, \dots, x_n), (y_1, \dots, y_n)) := \max_{k=1, \dots, n} |x_k - y_k| \quad (1)$$

for all  $x, y \in \mathbb{R}^n$ .

We also define the set  $B_r(x)$  for  $r > 0$  and  $x \in \mathbb{R}^n$  as the open ball using this distance, i.e. as the hypercube of side length  $2r$  centered at  $x$ .

#### 2.1.2 Uniformly Discrete Set

**Definition.** A subset  $\Gamma \subseteq \mathbb{R}^n$  is called *uniformly discrete* if there exists a constant  $q > 0$ , the *packing radius*, such that  $d(x, y) \geq q$  for all  $x, y \in \Gamma$  with  $x \neq y$ .

#### 2.1.3 Discrete Operator and Matrix Elements

**Definition.** A *discrete operator*  $H$  in dimension  $n \in \mathbb{N}$  is a bounded operator on a separable Hilbert space  $\mathcal{H}$ , together with an orthonormal basis  $(e_i)_{i \in \Gamma}$  indexed by a uniformly discrete set  $\Gamma \subset \mathbb{R}^n$ .

We also define the *matrix elements* of a discrete operator  $H$  at points  $x, y \in \Gamma$  as  $H_{xy} = \langle e_x, H e_y \rangle$ .

We shall always represent discrete operators with respect to the special basis  $(e_i)_{i \in \Gamma}$  and identify  $\mathcal{H}$  with  $\ell^2(\Gamma)$  through the basis isomorphism.

#### 2.1.4 Finite-Dimensional Subspaces and Orthogonal Projections

For any  $x \in \mathbb{R}^n$  and  $r > 0$ , the finite-dimensional subspace  $\mathcal{H}_{B_r(x)} \subset \mathcal{H}$  is defined by  $\mathcal{H}_{B_r(x)} := \text{span}\{e_y | y \in B_r(x) \cap \Gamma\}$ , and the orthogonal projection onto  $\mathcal{H}_{B_r(x)}$  is denoted by  $\mathbf{1}_{B_r(x)}$ .

#### 2.1.5 Short- and Finite-Range Operators

**Definition.** Let  $H$  be a discrete operator in dimension  $n$ . Then  $H$  is called *short-range* if there exists  $C$  and  $\varepsilon > 0$  such that

$$|H_{xy}| \leq C d(x, y)^{-(n+\varepsilon)} \quad (2)$$

for all  $x, y \in \Gamma$ .

**Definition.** A discrete operator  $H$  is said to have *finite range* if there exists a constant  $m > 0$  (the maximal hopping length) such that  $H_{xy} = 0$  whenever  $d(x, y) > m$ .

### 2.1.6 Equivalent Action and Finite Local Complexity

**Definition.** A discrete operator  $H$  is said to have an *equivalent action* on two subsets  $A, B \subseteq \Gamma$  if there is  $t \in \mathbb{R}^n$  such that  $B = A + t$  and if there exists a  $U : A \rightarrow S^1 \subset \mathbb{C}$ ,  $z \mapsto U(z)$  such that for any  $a_1, a_2 \in A$  we have

$$H_{b_1 b_2} = U(a_1) H_{a_1 a_2} U(a_2)^* \quad (3)$$

where  $b_1 = a_1 + t$ ,  $b_2 = a_2 + t$ .

**Remark.** Note that for any operator  $H$ , the equivalent action of  $H$  defines an equivalent relation on subsets of  $\Gamma$ .

**Definition.** A discrete operator  $H$  is said to have *finite local complexity* (FLC) if for any  $r > 0$ , the set  $\{\Gamma \cap B_r(x) | x \in \mathbb{R}^n\}$  can be partitioned into finitely many equivalence classes with respect to the equivalent action of  $H$ .

### 2.1.7 Hausdorff Distance

**Definition.** Let  $(X, d)$  be a metric space, and  $\mathcal{F}(X)$  the set of all nonempty bounded closed sets in  $X$ . For  $(A, B) \in \mathcal{F}(X) \times \mathcal{F}(X)$ , the *Hausdorff distance* between  $A$  and  $B$  is defined as [2]

$$d_{\text{Hausdorff}}(A, B) := \max \left\{ \sup_{a \in A} d(a, B), \sup_{b \in B} d(A, b) \right\} \quad (4)$$

where  $d(x_0, E) := \inf_{e \in E} d(x_0, e)$  is the distance of a point  $x_0 \in X$  to a nonempty subset  $E \subseteq X$ .

## 2.2 The Spectral and Pseudospectral Problems

Next, we give some additional definitions relevant to the discussion of error bounds on the approximant spectra.

### 2.2.1 Normal Operators

**Definition.** Let  $A$  be a linear operator on an inner product space  $V$ . Then  $A$  is *normal* if and only if  $AA^* = A^*A$ .

**Remark.** All self-adjoint and all isometric linear operators are normal. One important property of normal operators is that eigenvectors belonging to distinct eigenvalues are orthogonal, that is, for a normal operator  $A$ , let  $Av_1 = \lambda_1 v_1$  and  $Av_2 = \lambda_2 v_2$ , then  $\langle v_1, v_2 \rangle = 0$  whenever  $\lambda_1 \neq \lambda_2$  [3].

### 2.2.2 The Pseudospectrum

From mechanical oscillations to oscillations in quantum mechanics, most of the familiar applications of spectrum analysis involve matrices or operators that are normal or close to normal. Yet this familiarity can be deceiving: for nonnormal matrices or

operators, the spectrum alone may tell us surprisingly little. By way of example, if a system is modeled by a linear normal operator, then the resonant amplification experienced by it in response to a sinusoidally varying driving force with frequency  $\omega$  will be inversely proportional to the distance in the complex plane between  $\omega$  and the nearest eigenvalue. However, for a nonnormal system, the resonant amplification may be orders of magnitude greater. The resonances are simply not determined by the eigenvalues alone [7]. It is precisely this inadequacy that motivates the pseudospectrum.

**Definition.** Let  $H$  be a linear operator with spectrum  $\sigma(H)$ , then the map  $\rho_H : \mathbb{C} \rightarrow \mathbb{R}_+$  defined as

$$\rho_H(z) := \begin{cases} \|(H - z)^{-1}\|^{-1} & \text{for } z \notin \sigma(H) \\ 0 & \text{otherwise.} \end{cases} \quad (5)$$

is called the *lower norm function*.

**Definition (Variational).** Let  $H$  be a linear operator on a Hilbert space  $\mathcal{H}$ , then

$$\rho_H(z) := \inf_{\substack{\psi \in \mathcal{H} \\ \|\psi\|=1}} \|(H - z)\psi\| \quad (6)$$

is the *lower norm function*.

**Definition.** Let  $H$  be a linear operator, then the  $\varepsilon$ -pseudospectrum is defined as

$$\sigma_\varepsilon(H) := \{z \in \mathbb{C} \mid \rho_H(z) \leq \varepsilon\} \quad (7)$$

**Proposition.** Let  $H$  be a linear operator with spectrum  $\sigma(H)$ , then

$$\rho_H(z) \leq d(z, \sigma(H)) \quad (8)$$

with normality of  $H$  being sufficient for equality.

Thus, the  $\varepsilon$ -pseudospectrum of a normal operator is simply the  $\varepsilon$ -fattening of its spectrum.

**Remark.** Pseudospectral analysis offers a more informative object for studying the behavior of operators for which the spectrum alone is insufficient. In particular, it shifts the focus from exact eigenvalues to  $\varepsilon$ -approximate ones, which, as H. J. Landau observed [5], carry over smoothly from the normal case:

The distinction between them arises, and plays a crucial role, in passing to non-Hermitian equations where, as we have seen, true eigenvalues do not in general describe the operator adequately; on the other hand ... systems of mutually orthogonal  $\varepsilon$ -approximate eigenfunctions are easily constructed, and the behavior of  $\varepsilon$ -approximate eigenvalues, as well as their role in explaining the action of the operator, carry over smoothly from the Hermitian case.

**Theorem.** Let  $H$  be a closed operator on a complex Banach space, then the pseudospectra  $\{\sigma_\varepsilon(H)\}_{\varepsilon>0}$  have the following properties [7].

- Each  $\sigma_\varepsilon(H)$  is a nonempty open subset of  $\mathbb{C}$ , and any bounded connected component of  $\sigma_\varepsilon(H)$  intersects the spectrum  $\sigma(H)$ .
- The pseudospectra are strictly nested supersets of the spectrum:  $\bigcap_{\varepsilon>0} \sigma_\varepsilon(H) = \sigma(H)$ , and conversely, for any  $\delta > 0$ ,  $\sigma_{\varepsilon+\delta}(H) \supseteq \sigma_\varepsilon(H) + \Delta_\delta$ , where  $\Delta_\delta$  is the open disk of radius  $\delta$  and  $+$  is the Minkowski sum.

## 2.3 Approximating the (Pseudo-)Spectrum with Full Error Control

As shown in [4], as long as we are dealing with operators of finite local complexity, it is always possible to compute the spectrum in the normal case, and the pseudospectrum in the nonnormal one. In this section we provide the theorems that led to these results.

### 2.3.1 The Pseudospectral Inclusion Bound

An upper bound on  $\rho_H$  can be obtained by performing a singular value decomposition (SVD) on the rectangular matrix  $(H_{yz})_{y \in B_{r+m}(x), z \in B_r(x)}$ , where  $x \in \mathbb{R}^n$  and  $r, m > 0$  are fixed numbers.

Recall that a singular value decomposition of a matrix  $A \in \mathbb{C}^{m \times n}$  is a factorization into a product of matrices  $A = USV^*$ , where  $U \in \mathbb{C}^{m \times m}$  and  $V \in \mathbb{C}^{n \times n}$  are unitary, and  $S \in \mathbb{C}^{m \times n}$  is diagonal. The matrix  $S$  contains the square roots of the nonnegative eigenvalues of both  $AA^*$  and  $A^*A$ , and these are the *singular values* of  $A$  [6].

**Definition.** Let  $H$  be a discrete operator with finite range  $m > 0$ , let  $x \in \mathbb{R}^n$  be any point,  $r > 0$  and  $\lambda \in \mathbb{C}$ . The *uneven section* for these data is the linear operator

$$\begin{aligned} Q_{r,\lambda,x} : \mathcal{H}_{B_r(x)} &\rightarrow \mathcal{H}_{B_{r+m}(x)} \\ Q_{r,\lambda,x}(\psi) &= \mathbf{1}_{B_{r+m}(x)} (H - \lambda) \mathbf{1}_{B_r(x)} \psi \end{aligned}$$

Since  $\mathcal{H}_{B_r(x)}$  and  $\mathcal{H}_{B_{r+m}(x)}$  are finite dimensional, we can represent  $Q_{r,\lambda,x}$  as a rectangular matrix.

**Theorem.** Let  $H$  be a discrete operator with finite range  $m > 0$ , let  $x \in \mathbb{R}^n$  be any point,  $r > 0$  and  $\lambda \in \mathbb{C}$ . Let  $\varepsilon_{r,\lambda,x}$  be the smallest singular value of  $Q_{r,\lambda,x}$ . Then  $\lambda$  is contained in the  $\varepsilon_{r,\lambda,x}$ -pseudospectrum of  $H$ , that is,

$$\rho_H(\lambda) \leq \varepsilon_{r,\lambda,x} \tag{9}$$

*Proof.* We can write the SVD of  $Q_{r,\lambda,x}$  as

$$Q_{r,\lambda,x} = \sum_{i=1}^n |u_i\rangle s_i \langle v_i|$$

where  $(u_i)_{i=1,\dots,n}$  are the first  $n$  columns of  $U$ ,  $(v_i)_{i=1,\dots,n}$  are the columns of  $V$ , and  $(s_i)_{i=1,\dots,n}$  are the diagonal elements of  $\Sigma$ . We assume that the  $(s_i)_{i=1,\dots,n}$  are sorted in increasing order so that  $\varepsilon_{r,\lambda,x} = s_1$ .

Then  $v_1$ , considered as a vector in  $\mathcal{H}$  via the canonical inclusion  $\iota : \mathcal{H}_{B_r(x)} \rightarrow \mathcal{H}$ , is an  $\varepsilon_{r,\lambda,x}$ -quasimode for  $H$ . This follows from

$$\begin{aligned} (H - \lambda) v_1 &= (H - \lambda) \mathbf{1}_{B_r(x)} v_1 && \text{(since } v_1 \text{ is supported on } B_r(x)) \\ &= \mathbf{1}_{B_{r+m}(x)} (H - \lambda) \mathbf{1}_{B_r(x)} v_1 && \text{(by the finite range of } H) \\ &= Q_{r,\lambda,x} v_1 \\ &= s_1 u_1 \end{aligned}$$

and therefore  $\|(H - \lambda) v_1\| = s_1 \|u_1\| = s_1$ . We conclude that  $\|(H - \lambda) v_1\| \leq s_1 = \varepsilon_{r,\lambda,x}$  ( $a = b \Rightarrow a \leq b$ ), and thus  $v_1$  is an  $\varepsilon_{r,\lambda,x}$ -quasimode (by definition of a quasimode), which implies that  $\lambda \in \sigma_{\varepsilon_{r,\lambda,x}}(H)$  by definition of the spectrum (eq. 6).

### 2.3.2 The Spectral Gap Bound

In previous work [1], it was shown that the pseudospectral inclusion bound  $\varepsilon_{r,\lambda,x}$  always converges to  $\rho_H(\lambda)$  as  $r \rightarrow \infty$ , at least for normal, finite-range operators. Nonetheless, since the convergence is not uniform in the operator  $H$ , it cannot be used to compute a lower bound on  $\rho_H(\lambda)$ .

One way to circumvent this is by making use of the infimum of  $\varepsilon_{r,\lambda,x}$  over all  $x \in \mathbb{R}^n$ . More precisely, for

$$\varepsilon_{r,\lambda} := \inf_{x \in \mathbb{R}^n} \varepsilon_{r,\lambda,x}$$

we get the following theorem.

**Theorem.** Let  $n \in \mathbb{N}$ , let  $\Gamma \subseteq \mathbb{R}^n$  be a uniformly discrete set with packing radius  $q > 0$ , let  $H$  be a discrete operator on  $\ell^2(\Gamma)$  with finite range  $m > 0$  and let  $M := \sup_{x,y \in \Gamma} |H_{xy}|$ . Then for every  $\lambda \in \mathbb{C}$  and every  $r > m$  it holds that

$$\rho_H(\lambda) \geq \varepsilon_{r,\lambda} - \frac{C}{r} \tag{10}$$

where

$$C := mM \left( \frac{36m}{q} \right)^{n/2} \tag{11}$$

For the proof of this theorem see [4].

In practice, the importance of this theorem lies in the fact that for FLC operators, the infimum  $\varepsilon_{r,\lambda}$  can be computed by evaluating  $\varepsilon_{r,\lambda,x}$  for a finite number of suitably chosen points  $x \in \mathbb{R}^n$ .

### 2.3.3 A General Algorithm

Finally, combining bounds 9 and 10 leads to a general algorithm for computing the (pseudo-)spectrum for FLC operators with full error control. That is, given any short-range, discrete, normal, FLC operator  $H$  and  $k \in \mathbb{N}$ , one can derive an algorithm which computes an approximation  $\Gamma_k(H) \subseteq \mathbb{C}$  such that

$$d_{\text{Hausdorff}}(\sigma(H), \Gamma_k(H)) \leq 2^{-k} \tag{12}$$



Similarly, for nonnormal operators one can derive an algorithm which computes for any  $\varepsilon > 0$  and approximation  $\Gamma_k(H, \varepsilon)$  such that

$$d_{\text{Hausdorff}}(\sigma_\varepsilon(H), \Gamma_k(H, \varepsilon)) \leq 2^{-k} \quad (13)$$

For a derivation of either of these algorithms, see [\[4\]](#).

### 3 Conclusion and Future Work

## References

- [1] Matthew J. Colbrook, Bogdan Roman, and Anders C. Hansen. “How to Compute Spectra with Error Control”. In: *Phys. Rev. Lett.* 122 (25 2019). DOI: [10.1103/PhysRevLett.122.250201](https://doi.org/10.1103/PhysRevLett.122.250201).
- [2] James Dugundji. *Topology*. Allyn and Bacon, Inc., 1978.
- [3] Frederick W. Byron, Jr. and Robert W. Fuller. *Mathematics of Classical and Quantum Physics*. Dover Publications, Inc., 1992.
- [4] Paul Hege, Massimo Moscolari, and Stefan Teufel. “Computing the Spectrum and Pseudospectrum of Infinite-Volume Operators from Local Patches”. In: *American Mathematical Society* (2025).
- [5] H. J. Landau. “The Notion of Approximate Eigenvalues Applied to an Integral Equation of Laser Theory”. In: *Quarterly of Applied Mathematics* (1977).
- [6] Gilbert Strang. *Linear Algebra and Its Applications*. 4th. Cengage Learning, 2006.
- [7] Lloyd N. Trefethen and Mark Embree. *Spectra and Pseudospectra: The Behavior of Nonnormal Matrices and Operators*. Princeton University Press, 2005.
- [8] J. V. Uspensky. *Theory of Equations*. McGraw-Hill Book Company, Inc., 1948.