

SB - RC - Master equation

After applying the RC-map to the spin-boson Hamiltonian we find.

$$H = H_S + H_E + H_I$$

$$H_S = \frac{\omega}{2} \sigma_z + \Delta_0 b^\dagger b + g \sigma_z (b^\dagger + b) + \frac{g^2}{\Delta_0} \sigma_z^2 + \Delta_0 \Delta (b^\dagger b)^2$$

$$H_E = \sum_n \Delta_n b_n^\dagger b_n$$

$$H_I = (b + b^\dagger) \sum_n h_n (b_n + b_n^\dagger)$$

State time evolution - von Neumann

$$\dot{\rho}(t) = -i [H_S + H_E + H_I, \rho(t)] \quad (1)$$

Move to the interaction picture

$$\rho_I(t) = \exp(i(H_S + H_E)t) \rho(t) \exp(-i(H_S + H_E)t)$$

$$\stackrel{(1)}{\rightarrow} \dot{\rho}_I(t) = -i [H_I(t), \rho_I(t)]. \quad (2)$$

with

$$H_I(t) = \exp(i(H_S + H_E)t) H_I \exp(-i(H_S + H_E)t)$$

Formal integration of (2) yields

$$\rho_I(t) = \rho_I(0) - i \int_0^t dt' [H_I(t'), \rho_I(t')], \quad (3)$$

which after substituting in (2) leads to

$$\dot{\rho}_I(t) = -i [H_I(t), \rho_I(0)] - \int_0^t dt' [H_I(t), [H_I(t'), \rho_I(t')]] \quad (4)$$

Define now the reduced state of the system as

$$\bar{\rho}_I(t) = \text{Tr}_E(\rho_I(t)),$$

and assume that $\rho(0) = \bar{\rho}_S \otimes \rho_E$, with

$$\text{Tr}_E(H_I(t) \rho_I(0)) = 0. \quad (\text{which can be shown to hold for the conditions in our problem})$$

Trace over the environment on both sides of (4)


$$\stackrel{(4)}{\Rightarrow} \dot{\bar{\rho}}_I(t) = - \int_0^t dt' \text{Tr}_E ([H_I(t), [H_I(t'), \bar{\rho}_I(t')]]) \quad (5)$$

* Approximation: At this point we take

$$\rho_I(t) = \bar{\rho}_I(t) \otimes \rho_E.$$

- i) $H_I \ll H_S, H_B$
- ii) The state of the bath doesn't change (No memory).
- iii) The state of the system does change.
- iv) A weaker condition could have been imposed:

For interactions of the form $X A$, then



$$\text{Tr}_E ([A_I(t), [A(t'), \bar{\rho}_I(t')]]) \approx$$

$$\bar{\rho}_I(t) \otimes \underbrace{\text{Tr}_E ([A_I(t), [A(t'), \rho_E]])}_{\text{Bath correlations.}}$$

(Assume they don't change).

$$\stackrel{(5)}{\Rightarrow} \dot{\bar{\rho}}_I(t) = - \int_0^t dt' \text{Tr}_E ([H_I(t), [H_I(t'), \bar{\rho}_I(t') \otimes \rho_E]]) \quad (6)$$

Markov Approximation.

We assume the bath correlation time to be small.

- i) $\bar{\rho}_I(t') \rightarrow \bar{\rho}_I(t)$ (Markov approx: Yields a first order differential equation for $\bar{\rho}_I$).
- ii) We let the upper limit of the integral to go to infinity.

$$\begin{aligned} \stackrel{(6)}{\Rightarrow} \dot{\bar{\rho}}_I(t) &= - \int_0^{+\infty} dt' \text{Tr}_E ([H_I(t), [H_I(t'), \bar{\rho}_I(t) \otimes \rho_E]]) ; \quad \tau = t - t' \\ &= \int_0^{+\infty} d\tau \text{Tr}_E ([H_I(t), [H_I(t-\tau), \bar{\rho}_I(t) \otimes \rho_E]]). \end{aligned} \quad (7)$$

* We can choose appropriate forms for H_S and H_I and find different master equations.

Redfield equation

To recover the redfield equation, let us consider $H_I(t)$ in detail.

$$H_I(t) = \exp(i(H_S + H_E)t)(b + b^\dagger) \sum_k h_k (b_k + b_k^\dagger) \exp(-i(H_S + H_E)t).$$

Using $e^{iH_E t} b_k e^{-iH_E t} = b_k e^{-i\omega_k t}$

$$H_I(t) = \exp(iH_S t)(b + b^\dagger) \exp(-iH_S t) \sum_k h_k (b_k e^{-i\omega_k t} + b_k^\dagger e^{i\omega_k t}) \quad (8)$$

$\Gamma(t)$

Equation (7) on the Schrödinger equation can now be rewritten as

$$\begin{aligned} \dot{\rho}(t) &= -i[H_S, \rho(t)] - \int_0^{+\infty} d\tau \text{Tr}_E ([H_I, [H_I(t-\tau), \bar{\rho}(t) \otimes \rho_E]]) \\ &= -i[H_S, \bar{\rho}(t)] - \int_0^{+\infty} d\tau [\text{Tr}_E (\Pi(0) \Pi(\tau) \rho_E) [(b + b^\dagger), \bar{e}^{iH_S \tau} (b + b^\dagger) \bar{e}^{iH_S \tau}] \bar{\rho}(t)] \\ &\quad + \text{Tr}_E (\Pi(\tau) \Pi(0) \rho_E) [\bar{e}^{iH_S \tau} (b + b^\dagger) \bar{e}^{iH_S \tau} \bar{\rho}(t), (b + b^\dagger)] \\ &= -i[H_S, \bar{\rho}(t)] - \int_0^{+\infty} d\tau C(\tau) [(b + b^\dagger), \bar{e}^{iH_S \tau} (b + b^\dagger) \bar{e}^{iH_S \tau} \bar{\rho}(t)] \\ &\quad - \int_0^{+\infty} d\tau C(-\tau) [\bar{e}^{iH_S \tau} (b + b^\dagger) \bar{e}^{iH_S \tau} \bar{\rho}(t), (b + b^\dagger)] \quad (9) \end{aligned}$$

This is the Redfield equation reported by G. Shuller.

With the correlation function of the residual bath.

$$C(\tau) = \text{Tr}_E (\Pi(0) \Pi(\tau) \rho_E) = \text{Tr}_E (\Pi(-\tau) \Pi(0) \rho_E)$$

which we write also as.

$$C(\tau) = \frac{1}{2\pi} \int_{-\infty}^{+\infty} \delta(\omega) e^{-i\omega\tau} d\tau = C^*(-\tau)$$

To proceed further we use the eigen basis of H_{sys} ($H_{\text{sys}}|a\rangle = E_a|a\rangle$), to write

$$\begin{aligned}\dot{\bar{\rho}}(t) &= -i[H_s, \bar{\rho}(t)] - \sum_{ab} \langle a | (b + b^\dagger) | b \rangle \\ &\quad \times \left\{ - \int_0^{+\infty} d\tau C(\tau) e^{i(E_b - E_a)\tau} [(b + b^\dagger), |a\rangle \langle b| \bar{\rho}(t)] \right. \\ &\quad \left. - \int_0^{+\infty} d\tau C(-\tau) e^{i(E_b - E_a)\tau} [|a\rangle \langle b| \bar{\rho}(t), (b + b^\dagger)] \right\}.\end{aligned}$$

Let us now consider

$$\begin{aligned}\int_0^{+\infty} d\tau C(\tau) e^{i(E_b - E_a)\tau} &= \int_{-\infty}^{+\infty} d\tau \Theta(\tau) C(\tau) e^{i(E_b - E_a)\tau} \\ &= \frac{1}{2\pi} \int_{-\infty}^{+\infty} dw \gamma(w) \int_{-\infty}^{+\infty} d\tau e^{i(E_b - E_a - w)\tau} \frac{1}{2} (1 + \text{sng}(\tau)) \\ &= \frac{\gamma(E_b - E_a)}{2} + \frac{i}{2\pi} P \int_{-\infty}^{+\infty} dw \frac{\gamma(w)}{E_b - E_a - w} \approx \frac{\gamma(E_b - E_a)}{2}.\end{aligned}$$

The equation of motion becomes

$$\dot{\bar{\rho}}(t) = -i[H_s, \bar{\rho}] + \sum_{ab} \langle a | (b + b^\dagger) | b \rangle \left\{ -\gamma_{ab} [(b + b^\dagger), |a\rangle \langle b| \bar{\rho}(t)] \right. \\ \left. - \gamma_{ba} [|a\rangle \langle b| \bar{\rho}(t), (b + b^\dagger)] \right\} \quad (10)$$

The Markovian ME.

We continue with the calculation in the interaction picture starting with equation (7).

$$\dot{\bar{\rho}}_I(t) = - \int_0^{+\infty} d\tau \text{Tr}_E ([H_I(t), [H_I(t-\tau), \bar{\rho}_I(t) \otimes \rho_E]]). \quad (7)$$

We now use Eq.(8) to write.

$$H_I(t) = \exp(iH_{\text{st}}t)(b + b^+) \exp(-iH_{\text{st}}t) \Gamma(t).$$

At this point we evaluate (Luis already did it!!)

$$\begin{aligned} e^{-iH_{\text{st}}\tau}(b + b^+) e^{iH_{\text{st}}\tau} &= \underbrace{(b + \frac{g}{\Omega} J_z)}_{X} e^{-i\Omega\tau} + \underbrace{(b^+ + \frac{g}{\Omega} J_z)}_{X^+} e^{i\Omega\tau} \\ &= X e^{-i\Omega\tau} + X^+ e^{i\Omega\tau} \\ &= X(t) + X^+(t). \end{aligned}$$

Equation (7) can now be rewritten as

$$\dot{\bar{\rho}}_I(t) = - \int_0^{+\infty} d\tau \text{Tr}_E ([((X(t) + X^+(t))\Gamma(t), [(X(t-\tau) + X^+(t-\tau))\Gamma(t-\tau), \bar{\rho}_I(t) \otimes \rho_E]]])$$

After some algebra

$$\begin{aligned} [A, B]^T \cdot (AB - BA)^T &= (B^+ A^+ - A^+ B^+) \\ &= [B^+, A^+] = -[A^+, B^+] \end{aligned}$$

$$\begin{aligned} \dot{\bar{\rho}}_I(t) &= - \int_0^{+\infty} d\tau \{ C(\tau) [(X(t) + X^+(t)), (X(t-\tau) + X^+(t-\tau))] \bar{\rho}_I(t) \\ &\quad + C(-\tau) [\bar{\rho}_I(t)(X(t-\tau) + X^+(t-\tau)), (X(t) + X^+(t))] \} \end{aligned}$$

There are several terms that are obtained from expanding the commutators.

Let us consider just a typical one

$$\begin{aligned} (X(t) + X^+(t))(X(t-\tau) + X^+(t-\tau)) &= (X e^{-i\Omega t} + X^+ e^{i\Omega t}) \\ &\quad \times (X e^{-i\Omega(t-\tau)} + X^+ e^{i\Omega(t-\tau)}) \\ &= X^2 e^{-i\Omega(2t-\tau)} + X^+ X e^{i\Omega t} + X X^+ e^{-i\Omega t} + X^{+2} e^{i\Omega(2t-\tau)} \end{aligned}$$

Rotating wave approximation.

In the above expression we neglect the rapidly rotating terms with t , as they will give negligible contributions.

$$(X(t) + X^+(t))(X(t-\tau) + X^+(t-\tau)) \approx X^+ X e^{i\Omega t} + X X^+ e^{-i\Omega t}.$$

Equation (7) now reduces to

$$\begin{aligned}\dot{\bar{\rho}}_I(t) = & - \int_0^{+\infty} d\tau C(\tau) \left\{ (X^+ X e^{-i\omega\tau} + X X^+ e^{-i\omega\tau}) \bar{\rho}_I(t) \right. \\ & \left. - (X \bar{\rho}_I(t) X^+ e^{i\omega\tau} + X^+ \bar{\rho}_I(t) X e^{i\omega\tau}) \right\} \\ & - \int_0^{+\infty} d\tau C(-\tau) \left\{ \bar{\rho}_I(t) (X^+ X e^{-i\omega\tau} + X X^+ e^{-i\omega\tau}) \right. \\ & \left. - (X \bar{\rho}_I(t) X^+ e^{-i\omega\tau} + X^+ \bar{\rho}_I(t) X e^{-i\omega\tau}) \right\}\end{aligned}$$

We now consider

$$\int_0^{+\infty} d\tau C(\tau) e^{i\omega\tau} = \Gamma(\omega) \quad C(-\tau) = C^*(\tau).$$

$$\int_0^{+\infty} d\tau C(-\tau) e^{i\omega\tau} = \int_0^{+\infty} d\tau C^*(\tau) e^{i\omega\tau} = \left(\int_0^{+\infty} d\tau C(\tau) e^{-i\omega\tau} \right)^* = \Gamma^*(-\omega)$$

$$\Rightarrow \dot{\bar{\rho}}_I(t) = -\Gamma(\omega) (X^+ X \bar{\rho}_I(t) - X \bar{\rho}_I(t) X^+) - \Gamma^*(\omega) (\bar{\rho}_I(t) X^+ X - X \bar{\rho}_I(t) X^+) \\ - \Gamma(-\omega) (X X^+ \bar{\rho}_I(t) - X^+ \bar{\rho}_I(t) X) - \Gamma^*(-\omega) (\bar{\rho}_I(t) X X^+ - X^+ \bar{\rho}_I(t) X)$$

We now use

$$\Gamma(\omega) = \frac{\gamma(\omega)}{2} + \frac{i}{2} \sigma(\omega) \quad \Gamma(-\omega) = \frac{\bar{\gamma}(\omega)}{2} + \frac{i}{2} \bar{\sigma}(\omega)$$

$$\gamma(\omega) = J(\omega) (\bar{n}(\omega) + 1) \quad \bar{\gamma}(\omega) = J(\omega) \bar{n}(\omega)$$

$$\bar{n}(\omega) = (e^{\beta\omega} - 1)^{-1}$$

and arrive to the Lindblad master equation. (We neglect the Lamb shift)

$$\dot{\bar{\rho}}_I(t) = +\gamma (X \bar{\rho}_I(t) X^+ - \frac{1}{2} \{ X^+ X, \bar{\rho}_I(t) \}) + \bar{\gamma} (X^+ \bar{\rho}_I(t) X - \frac{1}{2} \{ X X^+, \bar{\rho}_I(t) \}),$$

Our lovely master equation! (in the interaction picture)

Remember: $X = b + \frac{g}{\omega} \sigma_x$. This non-trivial change is what is going to make the difference!

Can we find a left unitary that separates this?

Going to the Schrödinger picture -

$$\dot{\bar{\rho}}(t) = -i[H_S, \bar{\rho}(t)] + \gamma (x \bar{\rho}(t) x^+ - \frac{1}{2} \{x^+ x, \bar{\rho}(t)\}) + \bar{\gamma} (x^+ \bar{\rho}(t) x - \frac{1}{2} \{x x^+, \bar{\rho}(t)\})$$

master equation! (in the Schrödinger picture)