


Phytoplankton departure from each site's baseline: genera
Bray-Curtis to baseline
0.00
0.05
0.10
0.15
0.20
0.25
0.30
1750
1800
1850
1900
1950
2000
Phytoplankton departure from each site's baseline: genera, without Chaetoceros
Bray-Curtis to baseline
0.0
0.2
0.4
0.6
0.8
1750
1800
1850
1900
1950
2000
Phytoplankton departure from each site's baseline: groups
Bray-Curtis to baseline
0.00
0.05
0.10
0.15
0.20
0.25
0.30
1750
1800
1850
1900
1950
2000
dotted lines = largest within-baseline dissimilarity per site (the natural range)


MC - who moved (late period)
Chaetoceros
Pedinellales_X
Micromonas
unassigned Diatom
Dolichomastigaceae-B
Porosira
unassigned Cryptophyte
Triparma
0.00
0.01
0.02
0.03
0.04
BC contribution
p68 - who moved (late period)
Chaetoceros
Pedinellales_X
Stellarima
unassigned Diatom
Micromonas
Porosira
Dolichomastigaceae-B
Thalassiosira
0.000
0.001
0.002
0.003
0.004
BC contribution
p56 - who moved (late period)
Chaetoceros
Pedinellales_X
Micromonas
unassigned Diatom
Porosira
Dolichomastigaceae-B
unassigned Cryptophyte
Thalassiosira
0.000
0.001
0.002
0.003
BC contribution
BB - who moved (late period)
Chaetoceros
Micromonas
Thalassiosira
Pyramimonas
Phaeocystis
Pedinellales_X
Trieres
Porosira
0.000
0.005
0.010
0.015
0.020
0.025
0.030
0.035
BC contribution
SC - who moved (late period)
Chaetoceros
Micromonas
Pedinellales_X
Phaeocystis
Dolichomastigaceae-B
Thalassiosira
Pyramimonas
unassigned Marine stramenopile
0.000
0.001
0.002
0.003
0.004
0.005
0.006
0.007
BC contribution
increased
decreased