

CHAPTER 13

RANDOM GRAPHS WITH GENERAL DEGREE DISTRIBUTIONS

This chapter describes more sophisticated random graph models that mimic networks with arbitrary degree distributions

IN THE previous chapter we looked at the classic random graph model, in which pairs of vertices are connected at random with uniform probabilities. Although this model has proved tremendously useful as a source of insight into the structure of networks, it also has, as described in Section 12.8, a number of serious shortcomings. Chief among these is its degree distribution, which follows the Poisson distribution and is quite different from the degree distributions seen in most real-world networks. In this chapter we show how we can create more sophisticated random graph models, which incorporate arbitrary degree distributions and yet are still exactly solvable for many of their properties in the limit of large network size.

The fundamental mathematical tool that we will use to derive the results of this chapter is the probability generating function. We have already seen in Section 12.6 one example of a generating function, which was useful in the calculation of the distribution of component sizes in the Poisson random graph. We begin this chapter with a more formal introduction to generating functions and to some of their properties which will be useful in later calculations. Readers interested in pursuing the mathematics of generating functions further may like to look at the book by Wilf [329].[185](#)

13.1 GENERATING FUNCTIONS

Suppose we have a probability distribution for a non-negative integer variable, such that separate instances, occurrences, or draws of this variable are independent and have value k with probability p_k . A good example of such a distribution is the distribution of the degrees of randomly chosen vertices in a network. If the fraction of vertices in a network with degree k is p_k then p_k is also the probability that a randomly chosen vertex from the network will have degree k .

The *generating function* for the probability distribution p_k is the polynomial

$$g(z) = p_0 + p_1z + p_2z^2 + p_3z^3 + \dots = \sum_{k=0}^{\infty} p_k z^k.$$

(13.1)

Sometimes a function of this kind is called a *probability generating function* to distinguish it from another common type of function, the *exponential generating function*. We will not use exponential generating functions in this book, so for us all generating functions will be probability generating functions.

If we know the generating function for a probability distribution p_k then we can recover the values of p_k by differentiating:

$$p_k = \frac{1}{k!} \left. \frac{d^k g}{dz^k} \right|_{z=0}.$$

(13.2)

Thus the generating function gives us complete information about the probability distribution and vice versa. The distribution and the generating function are really just two different representations of the same thing. As we will see, it is easier in many cases to work with the generating function than with the probability distribution and doing so leads to many useful new results about networks.

13.1.1 EXAMPLES

Right away let us look at some examples of generating functions. Suppose our variable k takes only the values 0, 1, 2, and 3, with probabilities p_0, p_1, p_2 , and p_3 , respectively, and no other values. In that case the corresponding generating function would take the form of a cubic polynomial:

$$g(z) = p_0 + p_1 z + p_2 z^2 + p_3 z^3.$$

(13.3)

For instance, if we had a network in which vertices of degree 0, 1, 2, and 3 occupied 40%, 30%, 20%, and 10% of the network respectively then

$$g(z) = 0.4 + 0.3 z + 0.2 z^2 + 0.1 z^3.$$

(13.4)

As another example, suppose that k follows a Poisson distribution with mean c :

$$p_k = e^{-c} \frac{c^k}{k!}.$$

(13.5)

Then the corresponding generating function would be

$$g(z) = e^{-c} \sum_{k=0}^{\infty} \frac{(cz)^k}{k!} = e^{c(z-1)}.$$

(13.6)

Alternatively, suppose that k follows an exponential distribution of the form

$$p_k = C e^{-\lambda k},$$

(13.7)

with $\lambda > 0$. The normalizing constant is fixed by the condition that $\sum_k p_k = 1$, which gives $C = 1 - e^{-\lambda}$ and hence

$$p_k = (1 - e^{-\lambda}) e^{-\lambda k}.$$

(13.8)

Then

$$g(z) = (1 - e^{-\lambda}) \sum_{k=0}^{\infty} (e^{-\lambda} z)^k = \frac{e^{\lambda} - 1}{e^{\lambda} - z},$$

(13.9)

so long as $z < e^{\lambda}$. (If $z \geq e^{\lambda}$ the generating function diverges. Normally, however, we will be interested in generating functions only in the range $0 \leq z \leq 1$ so, given that $\lambda > 0$ and hence $e^{\lambda} > 1$, the divergence at e^{λ} will not be a problem.)

13.1.2 POWER-LAW DISTRIBUTIONS

One special case of particular interest in the study of networks is the power-law distribution. As we saw in Section 8.4, a number of networks, including the World Wide Web, the Internet, and citation networks, have degree distributions that follow power laws quite closely and this turns out to have interesting consequences that set these networks apart from others. To create and solve models of these networks it will be important for us to be able to write down generating functions for power-law distributions.

There are various forms that are used to represent power laws in practice but the simplest choice, which we will use in many of our calculations, is the “pure” power law

$$p_k = C k^{-\alpha},$$

(13.10)

for constant $\alpha > 0$. This expression cannot apply all the way down to $k = 0$, however, or it would diverge. So commonly one stops at $k = 1$. The normalization constant C can then be calculated from the condition that $\sum_k p_k = 1$, which gives

$$C \sum_{k=1}^{\infty} k^{-\alpha} = 1.$$

(13.11)

The sum unfortunately cannot be performed in closed form. It is, however, a common enough sum that it has a name—it is called the *Riemann zeta function*, denoted $\zeta(\alpha)$:

$$\zeta(\alpha) = \sum_{k=1}^{\infty} k^{-\alpha}.$$

(13.12)

Thus we can write $C = 1/\zeta(\alpha)$ and

$$p_k = \begin{cases} 0 & \text{for } k = 0, \\ k^{-\alpha}/\zeta(\alpha) & \text{for } k \geq 1. \end{cases}$$

(13.13)

Although there is no closed-form expression for the zeta function, there exist good numerical methods for calculating its value accurately, and many programming languages and numerical software packages include functions to calculate it.

For this probability distribution the generating function is

$$g(z) = \frac{1}{\zeta(\alpha)} \sum_{k=1}^{\infty} k^{-\alpha} z^k.$$

(13.14)

Again the sum cannot be expressed in closed form, but again it has a name—it is called the *polylogarithm* of z and is denoted $\text{Li}_{\alpha}(z)$:

$$\text{Li}_{\alpha}(z) = \sum_{k=1}^{\infty} k^{-\alpha} z^k.$$

(13.15)

Thus we can write

$$g(z) = \frac{\text{Li}_{\alpha}(z)}{\zeta(\alpha)}.$$

(13.16)

This is not completely satisfactory. We would certainly prefer a closed-form expression as in the case of the Poisson and exponential distributions of Eqs. (13.6) and (13.9). But we can live with it. Enough properties of the polylogarithm and zeta functions are known that we can carry out useful manipulations of the generating function. In particular, since derivatives of our generating functions will be important to us, we note the following useful relation:

$$\frac{\partial \text{Li}_{\alpha}(z)}{\partial z} = \frac{\partial}{\partial z} \sum_{k=1}^{\infty} k^{-\alpha} z^k = \sum_{k=1}^{\infty} k^{-(\alpha-1)} z^{k-1} = \frac{\text{Li}_{\alpha-1}(z)}{z}.$$

(13.17)

We should note also that in real-world networks the degree distribution does not usually follow a power law over its whole range—the distribution is not a “pure” power law in the sense above. Instead, it typically obeys a power law reasonably closely for values of k above some minimum value $k_{\min}$ but below that point it has some other behavior. In this case the generating function will take the form

$$g(z) = Q_{k_{\min}-1}(z) + C \sum_{k=k_{\min}}^{\infty} k^{-\alpha} z^k,$$

(13.18)

where $Q_n(z) = \sum_{k=0}^n p_k z^k$ is a polynomial in z of degree n and C is a normalizing constant. The sum in Eq. (13.18) also has its own name: it is called the *Lerch transcendent*.^{[186](#)} In the calculations in this book we will stick to the pure power law, since it illustrates nicely the interesting properties of power-law degree distributions and is relatively simple to deal with, but for serious modeling one might sometimes have to use the cut-off form, Eq. (13.18).

13.1.3 NORMALIZATION AND MOMENTS

Let us now look briefly at some of the properties of generating functions that will be useful to us. First of all, note that if we set $z = 1$ in the definition of the generating function, $g(z) = \sum_k p_k z^k$ (Eq. (13.1)), we get

$$g(1) = \sum_{k=0}^{\infty} p_k.$$

(13.19)

If the probability distribution is normalized to unity, $\sum_k p_k = 1$, as are all the examples above, then this immediately implies that

$$g(1) = 1.$$

(13.20)

For most of the generating functions we will look at, this will be true, but not all. As a counter-example, consider the generating function for the sizes of the small components in the Poisson random graph defined in Eq. (12.26). The probabilities π_s appearing in this generating function were the probabilities that a randomly chosen vertex belongs to a small component of size s . If we are in the regime where there is a giant component in the network then not all vertices belong to a small component, and hence the probabilities π_s do not add up to one. In fact, their sum is equal to the fraction of vertices not in the giant component.

The derivative of the generating function $g(z)$ of Eq. (13.1) is

$$g'(z) = \sum_{k=0}^{\infty} k p_k z^{k-1}.$$

(13.21)

(We will use the primed notation $g'(z)$ for derivatives of generating functions extensively in this chapter, as it proves much less cumbersome than the more common notation dg/dz .)

If we set $z = 1$ in Eq. (13.21) we get

$$g'(1) = \sum_{k=0}^{\infty} k p_k = \langle k \rangle,$$

(13.22)

which is just the average value of k . Thus, for example, if p_k is a degree distribution, we can calculate the average degree directly from the generating function by differentiating. This is a very convenient trick. In many cases we will calculate a probability distribution of interest by calculating first its generating function. In principle, we can then extract the distribution itself by applying Eq. (13.2) and so derive any other quantities we want such as averages. But Eq. (13.22) shows us that we don't always have to do this. Some of the quantities we will be interested in can be calculated directly from the generating function without going through any intermediate steps.

In fact, this result generalizes to higher moments of the probability distribution as well. For instance, note that

$$z \frac{d}{dz} \left(z \frac{dg}{dz} \right) = \sum_{k=0}^{\infty} k^2 p_k z^k,$$

(13.23)

and hence, setting $z = 1$, we can write

$$\langle k^2 \rangle = \left[\left(z \frac{d}{dz} \right)^2 g(z) \right]_{z=1}.$$

(13.24)

It is not hard to show that this result generalizes to all higher moments as well:

$$\langle k^m \rangle = \left[\left(z \frac{d}{dz} \right)^m g(z) \right]_{z=1}.$$

(13.25)

This result can also be written as

$$\langle k^m \rangle = \frac{d^m g}{d(\ln z)^m} \Big|_{z=1}.$$

(13.26)

13.1.4 POWERS OF GENERATING FUNCTIONS

Perhaps the most useful property of generating functions—and the one that makes them important for the study of networks—is the following. Suppose we are given a distribution p_k with generating function $g(z)$. And suppose we have m integers k_i , $i = 1 \dots m$, which are independent random numbers drawn from this distribution. For instance, they could be the degrees of m randomly chosen vertices in a network with degree distribution p_k . Then the probability distribution of the sum $\sum_{i=1}^m k_i$ of those m integers has generating function $[g(z)]^m$. This is a very powerful result and it is worth taking a moment to see how it arises and what it means.

Given that our integers are independently drawn from the distribution p_k , the probability that they take a particular set of values $\{k_i\}$ is simply $\prod_i p_{k_i}$ and the probability π_s that the values drawn add up to a specific sum s is the sum of these probabilities over all sets $\{k_i\}$ that add up to s :

$$\pi_s = \sum_{k_1=0}^{\infty} \dots \sum_{k_m=0}^{\infty} \delta(s, \sum_i k_i) \prod_{i=1}^m p_{k_i},$$

(13.27)

where $\delta(a, b)$ is the Kronecker delta. Then the generating function $h(z)$ for the distribution π_s is

$$\begin{aligned} h(z) &= \sum_{s=0}^{\infty} \pi_s z^s \\ &= \sum_{s=0}^{\infty} z^s \sum_{k_1=0}^{\infty} \dots \sum_{k_m=0}^{\infty} \delta(s, \sum_i k_i) \prod_{i=1}^m p_{k_i} \\ &= \sum_{k_1=0}^{\infty} \dots \sum_{k_m=0}^{\infty} z^{\sum_i k_i} \prod_{i=1}^m p_{k_i} \\ &= \sum_{k_1=0}^{\infty} \dots \sum_{k_m=0}^{\infty} \prod_{i=1}^m p_{k_i} z^{k_i} = \left[\sum_{k=0}^{\infty} p_k z^k \right]^m \\ &= [g(z)]^m. \end{aligned}$$

(13.28)

Thus, for example, if we know the degree distribution of a network, it is a straightforward matter to calculate the probability distribution of the sum of the degrees of m randomly chosen vertices from that network. This will turn out to be important in the developments that follow.

13.2 THE CONFIGURATION MODEL

Let us turn now to the main topic of this chapter, the development of the theory of random graphs with general degree distributions.

We can turn the random graph of Chapter 12 into a much more flexible model for networks by modifying it so that the degrees of its vertices are no longer restricted to having a Poisson distribution, and in fact it is possible to modify the model so as to give the network any degree distribution we please. Just as with the Poisson random graph, which can be defined in several slightly different ways, there is more than one way to define random graphs with general degree distributions. Here we describe two of them, which are roughly the equivalent of the $G(n, m)$ and $G(n, p)$ random graphs of Section 12.1.

The most widely studied of the generalized random graph models is the *configuration model*. The configuration model is actually a model of a random graph with a given degree *sequence*, rather than degree distribution. That is, the exact degree of each individual vertex in the network is fixed, rather than merely the probability distribution from which those degrees are chosen. This in turn fixes the number of edges in the network, since the number of edges is given by Eq. (6.21) to be $m = \frac{1}{2} \sum_i k_i$. Thus this model is in some ways analogous to $G(n, m)$, which also fixes the number of edges. (It is quite simple, however, to modify the model for cases where only the degree distribution is known and not the exact degree sequence. We describe how this is done at the end of this section.)

See Section 8.3 for a discussion of the distinction between degree sequences and degree distributions.

Suppose then that we specify the degree k_i that each vertex $i = 1 \dots n$ in our network is to take. We can create a random network with these degrees as follows. We give each vertex i a total of k_i “stubs” of edges as depicted in Fig. 13.1. There are $\sum_i k_i = 2m$ stubs in total, where m is the total number of edges. Then we choose two of the stubs uniformly at random and we create an edge by connecting them to one another, as indicated by the dashed line in the figure. Then we choose another pair from the remaining $2m - 2$ stubs, connect those, and so on until all the stubs are used up. The end result is a network in which every vertex has exactly the desired degree.

More specifically the end result is a particular *matching* of the stubs, a particular set of pairings of stubs with other stubs. The process above generates each possible matching of stubs with equal probability. Technically the configuration model is defined as the ensemble in which each matching with the chosen degree sequence appears with the same probability (those with any other degree sequence having probability zero), and the process above is a process for drawing networks from the configuration model ensemble.




Figure 13.1: The configuration model. Each vertex is given a number of “stubs” of edges equal to its desired degree. Then pairs of stubs are chosen at random and connected together to form edges (dotted line).

The uniform distribution over matchings in the configuration model has the important consequence that any stub in a configuration model network is equally likely to be connected to any other. This, as we will see, is the crucial property that makes the model solvable for many of its properties.

There are a couple of minor catches with the network generation process described here. First, there must be an even number of stubs overall if we want to end up with a network consisting only of vertices and edges, with no dangling stubs left over. This means that the sum $\sum_i k_i$ of the degrees must add up to an even number. We will assume that the degrees we have chosen satisfy this condition, otherwise it is clearly not possible to create a graph with the given degree sequence.

A second issue is that the network may contain self-edges or multiedges, or both. There is nothing in the network generation process that prevents us from creating an edge that connects a vertex to itself or that connects two vertices that are already connected by another edge. One might imagine that one could avoid this by rejecting the creation of any such edges during the process, but it turns out that this is not a good idea. A network so generated is no longer drawn uniformly from the set of possible matchings, which means that properties of the model can no longer be calculated analytically, at least by any means currently known. It can also mean that the network creation process breaks down completely. Suppose, for example, that we come to the end of the process, when there are just two stubs left to be joined, and find that those two both belong to the same vertex so that joining them would create a self-edge. Then either we create the self-edge or the network generation process fails.

In practice, therefore, it makes more sense to allow the creation of both multiedges and self-edges in our networks and the standard configuration model does so. Although some real-world networks have self-edges or multiedges in them, most do not, and to some extent this makes the configuration model less satisfactory as a network model. However, as shown below, the average number of self-edges and multiedges in the configuration model is a constant as the network becomes large, which means that the density of self-edges and multiedges tends to zero in this limit. This means, to all intents and purposes, that we can ignore the self-edges and multiedges in the large size limit.¹⁸⁷

A further issue with the configuration model is that, while all matchings of stubs appear with equal probability in the model, that does not mean that all *networks* appear with equal probability because more than one matching can correspond to the same network, i.e., the same topological connections between vertices. If we label the stubs to keep track of which is which, then there are typically many different ways we can join up pairs of labeled stubs to create the same final configuration of edges. Figure 13.2 shows an example of a set of eight matchings that all correspond to the same three-vertex network.




Figure 13.2: Eight stub matchings that all give the same network. This small network is composed of three vertices of degree two and hence having two stubs each. The stubs are lettered to identify them and there are two distinct permutations of the stubs at each vertex for a total of eight permutations overall. Each permutation gives rise to a different matching of stub to stub but all matchings correspond to the same topological configuration of edges, and hence there are eight ways in which this particular configuration can be generated by the stub matching process.

In general, one can generate all the matchings that correspond to a given network by taking any one matching for that network and permuting the stubs at each vertex in every possible way. Since the number of permutations of the k_i stubs at a vertex i is $k_i!$, this implies that the number of matchings corresponding to each network is $N(\{k_i\}) = \prod_i k_i!$, which takes the same value for all networks, since the degrees are fixed. This implies that in fact networks occur with equal probability in the configuration model: if there are $\Omega(\{k_i\})$ matchings, each occurring with the same probability, then each *network* occurs with probability N/Ω .

However, this is not completely correct. If a network contains self-edges or multiedges then not all permutations of the stubs in the network result in a new matching of stubs. Consider Fig. 13.3. Panel (a) shows a network with the same degree sequence as those of Fig. 13.2, but a different configuration of edges, having one self-edge and a multiedge consisting of two parallel single edges. In panel (b) we have permuted the stubs a and b at the ends of the self-edge but, as we can see, this has not resulted in a new matching of the stubs themselves. Stubs a and b are still connected to one another just as they were before. (The network is *drawn* differently now, but in terms of the matching and the topology of the edges nothing has changed from panel (a).) In panel (c) we have identically permuted the stubs at both ends of the multiedge. Again this has no effect on which stubs are matched with which others.




Figure 13.3: Permutations that do not produce new matchings. (a) The network shown here has the same degree sequence as those of Fig. 13.2 but a different configuration of edges, having one self-loop and a multiedge consisting of two parallel edges. (b) If we permute the stubs a and b

of the self-edge we do not generate a new matching, because a is still matched with b , just as before. (c) If we permute the stubs at either end of a multiedge in exactly the same way we do not generate a new matching, since each stub at one end of the multiedge is still matched with the same stub at the other end.

In general, for each multiedge in a network a permutation of the stubs at one end fails to generate a new matching if we simultaneously permute the stubs at the other end in the same way. This means that the total number of matchings is reduced by a factor of $A_{ij}!$, since A_{ij} is equal to the multiplicity of the edge between i and j . Indeed, this expression is correct even for vertex pairs not connected by a multiedge, if we adopt the convention that $0! = 1$. For self-edges there is a further factor of two because the interchange of the two ends of the edge does not generate a new matching. Combining these results, the number of matchings corresponding to a network turns out to be

$$N = \frac{\prod_i k_i!}{\prod_{i < j} A_{ij}! \prod_i A_{ii}!!},$$

(13.29)

where $n!! = n(n-2)(n-4) \dots 2$ with n even is the so-called double factorial of n . Then the total probability of a particular network within the configuration model ensemble is N/Ω as before. Since the denominator in Eq. (13.29) depends not only on the degree sequence but also on the structure of the network itself, different networks do appear with different probabilities.

As we mentioned, however, the average densities of self-edges and multiedges in the configuration model vanish as n becomes large, so that the variation in probabilities is relatively small in the large- n limit, but it nonetheless does occasionally assume some importance and is therefore worth bearing in mind (see, for instance, Ref. [220]).

As discussed above, we are sometimes (indeed often) interested in the case where it is the degree distribution of the network that is specified rather than the degree sequence. That is, we specify the probability distribution p_k from which the degree sequence is drawn rather than the sequence itself. We can define an obvious extension of the configuration model to this case: we draw a degree sequence from the specified distribution and then generate a network with that degree sequence using the technique described above. More precisely, we define an ensemble in which each degree sequence $\{k_i\}$ appears with probability $\prod_i p_{k_i}$. Then if we can calculate an average value $X(\{k_i\})$ for some quantity of interest X in the standard configuration model, the average value in the extended model is given by

$$\langle X \rangle = \sum_{k_1=0}^{\infty} \dots \sum_{k_n=0}^{\infty} X(\{k_i\}) \prod_{i=0}^n p_{k_i}.$$

(13.30)

In practice the difference between the two models is not actually very great. As we will see, the crucial parameter that enters into most of our configuration model calculations is the fraction of vertices that have each possible degree k . In the extended model above, this fraction is, by definition, equal to p_k in the limit of large n . If, on the other hand, the degree sequence is fixed then

we simply calculate the fraction from the degree sequence and then use those numbers. In either the case the formulas for calculated quantities are the same.

13.2.1 EDGE PROBABILITY IN THE CONFIGURATION MODEL

A central property of the configuration model is the probability p_{ij} of the occurrence of an edge between two specified vertices, i and j . Obviously if either vertex i or vertex j has degree zero then the probability of an edge is zero, so let us assume that $k_i, k_j > 0$. Now consider any one of the stubs that emerges from vertex i . What is the probability that this stub is connected by an edge to any of the stubs of vertex j ? There are $2m$ stubs in total, or $2m - 1$ excluding the one connected to i that we are currently looking at. Of those $2m - 1$, exactly k_j of them are attached to vertex j . So, given that any stub in the network is equally likely to be connected to any other, the probability that our particular stub is connected to any of those around vertex j is $k_j/(2m - 1)$. But there are k_i stubs around vertex i , so the total probability of a connection between i and j is

$$p_{ij} = \frac{k_i k_j}{2m - 1}.$$

(13.31)

Technically, since we have added the probabilities of independent events, this is really the average number of edges between i and j , rather than the probability of having an edge at all. But in the limit of large m , this number becomes small (for given k_i, k_j), and the average number of edges and the probability of an edge become equal. Also in the limit of large m we can ignore the $- 1$ in the denominator and hence we can write

$$p_{ij} = \frac{k_i k_j}{2m}.$$

(13.32)

Note that, even though we assumed $k_i, k_j > 0$, this expression also gives the right result if either degree is zero, namely that in that case the probability of connection is zero.

We can use this result, for example, to calculate the probability of having two edges between the same pair of vertices. The probability of having one edge between vertices i and j is p_{ij} as above. Once we have one edge between the vertices the number of available stubs at each is reduced by one, and hence the probability of having a second edge is given by Eq. (13.32) but with k_i and k_j each reduced by one: $(k_i - 1)(k_j - 1)/2m$. Thus the probability of having (at least) two edges, i.e., of having a multiedge between i and j , is $k_i k_j (k_i - 1)(k_j - 1)/(2m)^2$ and, summing this probability over all vertices and dividing by two (to avoid double counting of vertex pairs), we find that the expected total number of multiedges in the network is

$$\begin{aligned}\frac{1}{2(2m)^2} \sum_{ij} k_i k_j (k_i - 1)(k_j - 1) &= \frac{1}{2\langle k \rangle^2 n^2} \sum_i k_i (k_i - 1) \sum_j k_j (k_j - 1) \\ &= \frac{1}{2} \left[\frac{\langle k^2 \rangle - \langle k \rangle^2}{\langle k \rangle} \right],\end{aligned}\quad (13.33)$$

where

$$\langle k \rangle = \frac{1}{n} \sum_i k_i, \quad \langle k^2 \rangle = \frac{1}{n} \sum_i k_i^2,$$

(13.34)

and we have used $2m = \langle k \rangle n$ (see Eq. (6.23)). Thus the expected number of multiedges remains constant as the network grows larger, so long as $\langle k^2 \rangle$ is constant and finite, and the density of multiedges—the number per vertex—vanishes as $1/n$. We used this result in a number of our earlier arguments.¹⁸⁸

Another way to derive the expression in Eq. (13.32) is to observe that there are $k_i k_j$ possible edges we could form between vertices i and j , while the total number of possible edges in the whole graph is the number of ways of choosing a pair of stubs from the $2m$ total stubs, or $\binom{2m}{2} = m(2m - 1)$. The probability that any particular edge falls between i and j is thus given by the ratio $k_i k_j / m(2m - 1)$, and if we make a total of m edges then the expected total number of edges between i and j is m times this quantity, which gives us Eq. (13.31) again.

The only case in which this derivation is not quite right is for self-edges. In that case the number of pairs of stubs is not $k_i k_j$ but instead is $\binom{k_i}{2} = \frac{1}{2} k_i (k_i - 1)$ and hence the probability of a self-edge from vertex i to itself is

$$p_{ii} = \frac{k_i(k_i - 1)}{4m}.$$

(13.35)

We can use this result to calculate the expected number of self-edges in the network, which is given by the sum over all vertices i :

$$\sum_i p_{ii} = \sum_i \frac{k_i(k_i - 1)}{4m} = \frac{\langle k^2 \rangle - \langle k \rangle^2}{2\langle k \rangle},$$

(13.36)

This expression remains constant as $n \rightarrow \infty$ provided $\langle k^2 \rangle$ remains constant, and hence, as with the multiedges, the density of self-edges in the network vanishes as $1/n$ in the limit of large network size.

We can use Eqs. (13.32) and (13.35) to calculate a number of other properties of vertices in the configuration model. For instance, we can calculate the expected number n_{ij} of common neighbors that vertices i and j share. The probability that i is connected to another vertex l is p_{il} and the probability that j is connected to the same vertex would likewise normally be p_{jl} . However, as with the calculation of multiedges above, if we already know that i is connected to l , then the number of available stubs at vertex l is reduced by one and, rather than being given by the normal expression (13.32), the probability of a connection between j and l is $k_j (k_l - 1)/2m$. Multiplying the probabilities for the two edges and summing over l , we then get our expression for the expected number of common neighbors of i and j :

$$\begin{aligned} n_{ij} &= \sum_l \frac{k_i k_l}{2m} \frac{k_j (k_l - 1)}{2m} = \frac{k_i k_j}{2m} \frac{\sum_l k_l (k_l - 1)}{n \langle k \rangle} \\ &= p_{ij} \frac{\langle k^2 \rangle - \langle k \rangle}{\langle k \rangle}. \end{aligned}$$

(13.37)

Thus the probability of sharing a common neighbor is equal to the probability $p_{ij} = k_i k_j / 2m$ of having a direct connection times a multiplicative factor that depends only on the mean and variance of the degree distribution but not on the properties of the vertices i and j themselves.

In this calculation we have ignored the fact that the probability of self-edges, Eq. (13.35), is different from the probability for other edges. As we have seen, however, the density of self-edges in the configuration model tends to zero as $n \rightarrow \infty$, so in that limit it is usually safe to make the approximation that Eq. (13.32) applies for all i and j .

13.2.2 RANDOM GRAPHS WITH GIVEN EXPECTED DEGREE

The configuration model of the previous section is, as we have said, similar in some ways to the standard random graph $G(n, m)$ described in Section 12.1, in which we distribute a fixed number m of edges at random between n vertices. In the configuration model the total number of edges is again fixed, having value $m = \frac{1}{2} \sum_i k_i$, but in addition we now also fix the individual degree of every vertex as well.

It is natural to ask whether there is also an equivalent of $G(n, p)$ —the model in which only the probability of edges is fixed and not their number—and indeed there is. We simply place an edge between each pair of vertices i, j with independent probabilities taking the form of Eq. (13.32). We define a parameter c_i for each vertex and then place an edge between vertices i and j with probability $p_{ij} = c_i c_j / 2m$. As with the configuration model, we must allow self-edges if the model is to be tractable, and again self-edges have to be treated a little differently from ordinary edges. It turns out that the most satisfactory definition of the edge probability is

$$p_{ij} = \begin{cases} c_i c_j / 2m & \text{for } i \neq j, \\ c_i^2 / 4m & \text{for } i = j, \end{cases}$$

(13.38)

where m is now defined by

$$\sum_i c_i = 2m.$$

(13.39)

With this choice the average number of edges in the network is

$$\sum_{i \leq j} p_{ij} = \sum_{i < j} \frac{c_i c_j}{2m} + \sum_i \frac{c_i^2}{4m} = \sum_{ij} \frac{c_i c_j}{4m} = m,$$

(13.40)

as before. We can also calculate the average number of ends of edges connected to a vertex i , i.e., its average degree $\langle k_i \rangle$. Allowing for the fact that a self-edge contributes two ends of edges to the degree, we get

$$\langle k_i \rangle = 2p_{ii} + \sum_{j(\neq i)} p_{ij} = \frac{c_i^2}{2m} + \sum_{j(\neq i)} \frac{c_i c_j}{2m} = \sum_j \frac{c_i c_j}{2m} = c_i.$$

(13.41)

In other words the parameters c_i appearing in the definition of p_{ij} , Eq. (13.38), are the average or expected degrees in this model, just as the parameter c in $G(n, p)$ is the average degree of a vertex. The *actual* degree of a vertex could in principle take almost any value, depending on the luck of the draw about which edges happen to get randomly created and which do not. In fact one can show that the degree of vertex i will have a Poisson distribution with mean c_i , meaning that in practice it will be quite narrowly distributed about c_i , but there will certainly be some variation, unless c_i is zero.¹⁹¹ Note that c_i does not have to be an integer, unlike the degrees k_i appearing in the configuration model.

Thus in this model we specify the expected number of edges m and the expected degree sequence $\{c_i\}$ of the network but not the actual number of edges and actual degree sequence. This is again analogous to $G(n, p)$, in which we specify only the expected number of edges and not the actual number. Unfortunately, this means we usually cannot choose the degree *distribution* of our network, because the distribution of the actual degrees k_i is not the same as the distribution of the expected degrees c_i . This is a substantial disadvantage of the model since the degree distribution is widely considered to be a crucial property of networks.¹⁹²

This is unfortunate, because this model is in other respects a very nice one. It is straightforward to treat analytically and many of the derivations are substantially simpler for this model than for the configuration model. Nonetheless, because we place such a premium on being able to choose the degree distribution, this model is in fact hardly ever used in real calculations of the properties of networks. Instead, most calculations are made using the configuration model and this is the direction that we will take in this book as well. In the following sections, we describe how one can make use of the machinery of generating functions to calculate many of the properties of the configuration model exactly in the limit of large network size.

13.3 EXCESS DEGREE DISTRIBUTION

In the remainder of this chapter we describe the calculation of a variety of properties of the configuration model. We begin our discussion with some fundamental observations about the model—and networks in general—that will prove central to later developments.

Consider a configuration model with degree distribution p_k , meaning that a fraction p_k of the vertices have degree k . (We can consider either the standard version of the model in which the degree sequence is fixed, as in Section 13.2, or the version of Eq. (13.30) in which only the distribution is fixed but not the exact degree sequence.) The distribution p_k tells us the probability that a vertex chosen uniformly at random from our network has degree k . But suppose instead that we take a vertex (randomly chosen or not) and follow one of its edges (assuming it has at least one) to the vertex at the other end. What is the probability that this vertex will have degree k ?

The answer cannot just be p_k . For instance, there is no way to reach a vertex with degree zero by following an edge in this way, because a vertex with degree zero has no edges. So the probability of finding a vertex of degree zero is itself zero, and not p_0 .

In fact, the correct probability for general k is not hard to calculate. We know that an edge emerging from a vertex in a configuration model network has equal chance of terminating at any “stub” of an edge anywhere else in the network (see Section 13.2). Since there are $\sum_i k_i = 2m$ stubs in total, or $2m - 1$ excluding the one at the beginning of our edge, and k of them are attached to any particular vertex with degree k , our edge has probability $k/(2m - 1)$ of ending at any particular vertex of degree k . In the limit of large network size, where m becomes large (assuming the degree distribution, and hence the average degree, remain constant), we can ignore the -1 and just write this as $k/2m$.

Given that p_k is the total fraction of vertices in the network with degree k , the total number of such vertices is np_k , and hence the probability of our edge attaching to *any* vertex with degree k is

$$\frac{k}{2m} \times np_k = \frac{kp_k}{\langle k \rangle},$$

(13.42)

where $\langle k \rangle$ is the average degree over the whole network and we have made use of the fact that $2m = n\langle k \rangle$, Eq. (6.23).

Thus the probability that we reach a vertex of degree k upon following an edge in this way is proportional not to p_k but to kp_k . To put that another way, the vertex you reach by following an edge is not a typical vertex in the network. It is more likely to have high degree than a typical vertex. Physically, the reasoning behind this observation is that a vertex with degree k has k edges attached to it, and you can reach that vertex by following any one of them. Thus if we choose an edge and follow it you have k times the chance of reaching a vertex with degree k that you have of reaching a vertex with degree 1.

It is important to recognize that this is a property specifically of the configuration model (or similar random graph models). In the real world, the degrees of adjacent vertices in networks are

often correlated (see Section 7.13) and hence the probability of reaching a vertex of degree k when we follow an edge depends on what vertex we are coming from.¹⁹³ Nonetheless, it is found to apply approximately to many real-world networks, which is one of the reasons why insights gained from the configuration model are useful for understanding the world around us.

Equation (13.42) has some strange and counter-intuitive consequences. As an example, consider a randomly chosen vertex in the configuration model and let us calculate the average degree of a neighbor of that vertex. If we were using the configuration model to model a friendship network, for instance, the average degree of an individual's network neighbor would correspond to the average number of friends their friend has. This number is the average of the distribution in Eq. (13.42), which we get by multiplying by k and then summing over k thus:

$$\text{average degree of a neighbor} = \sum_k k \frac{k p_k}{\langle k \rangle} = \frac{\langle k^2 \rangle}{\langle k \rangle}.$$

(13.43)

Note that the average degree of a neighbor is thus different from the average degree $\langle k \rangle$ of a typical vertex in the network. In fact, it is in general larger, as we can show by calculating the difference

$$\frac{\langle k^2 \rangle}{\langle k \rangle} - \langle k \rangle = \frac{1}{\langle k \rangle} (\langle k^2 \rangle - \langle k \rangle^2) = \frac{\sigma_k^2}{\langle k \rangle},$$

(13.44)

where $\sigma_k^2 = \langle k^2 \rangle - \langle k \rangle^2$ is the variance of the degree distribution. The variance, which is the square of the standard deviation, is necessarily non-negative and indeed is strictly positive unless every single vertex in the network has the same degree. Let us assume that there is some variation in the degrees so that σ_k^2 is greater than zero. The average degree $\langle k \rangle$ is also greater than zero, unless all vertices have degree zero. Thus Eq. (13.44) implies that $\langle k^2 \rangle / \langle k \rangle - \langle k \rangle > 0$, or

$$\frac{\langle k^2 \rangle}{\langle k \rangle} > \langle k \rangle.$$

(13.45)

In other words, the average degree of the neighbor of a vertex is greater than the average degree of a vertex. In colloquial terms, "Your friends have more friends than you do."

At first sight, this appears to be a very strange result. Certainly it seems likely that there will be some vertices in the network with higher degree than the average. But there will also be some who have lower degree and when you average over all neighbors of all vertices surely the two should

cancel out. Surely the average degree of a neighbor should be the same as the average degree in the network as a whole. Yet Eq. (13.45) tells us that this is not so. And the equation really is correct. You can create a configuration model network on a computer and average the degrees of the neighbors of every vertex, and you'll find that the formula works to very high accuracy. Even more remarkably, as first shown by Feld [113], you can do the same thing with real networks and, although the configuration model formula doesn't apply exactly to these networks, the basic principle still seems to hold. Here, for instance, are some measurements for two academic collaboration networks, in which scientists are connected together by edges if they have coauthored scientific papers, and for a recent snapshot of the structure of the Internet at the autonomous system level:

Network	n	Average degree	Average neighbor degree	$\frac{\langle k^2 \rangle}{\langle k \rangle}$
Biologists	1 520 252	15.5	68.4	130.2
Mathematicians	253 339	3.9	9.5	13.2
Internet	22 963	4.2	224.3	261.5

According to these results a biologist's collaborators have, on average, more than four times as many collaborators as they do themselves. On the Internet, a node's neighbors have more than 50 times the average degree! Note that in each of the cases in the table the configuration model value of $\langle k^2 \rangle / \langle k \rangle$ overestimates the real average neighbor degree, in some cases by a substantial margin.¹⁹⁵ This is typical of calculations using simplified network models: they can give you a feel for the types of effect one might expect to see, or the general directions of changes in quantities. But they usually don't give quantitatively accurate predictions for the behavior of real networks.

The fundamental reason for the result, Eq. (13.45), is that when you go through the vertices of a network and average the degrees of the neighbors of each one, many of those neighbors appear in more than one average. In fact, a vertex with degree k will appear as one of the neighbors of exactly k other vertices, and hence appear in k of the averages. This means that high-degree vertices are over-represented in the calculations compared with low-degree ones and it is this bias that pushes up the overall average value.

In most of the calculations that follow, we will be interested not in the total degree of the vertex at the end of an edge but in the number of edges attached to that vertex *other* than the one we arrived along. For instance, if we want to calculate the size of the component to which a vertex i belongs then we will want to know first of all how many neighbors i has, and then how many neighbors those neighbors have, *other than* i , and so on.

The number of edges attached to a vertex other than the edge we arrived along is called the *excess degree* of the vertex and it is just one less than the total degree. Since the vertex at the end of an edge always has degree at least 1 (because of that edge) the minimum value of the excess degree is zero.

We can calculate the probability distribution of the excess degree from Eq. (13.43). The probability q_k of having excess degree k is simply the probability of having total degree $k + 1$ and, putting $k \rightarrow k + 1$ in Eq. (13.43), we get

$$q_k = \frac{(k+1)p_{k+1}}{\langle k \rangle}.$$

(Note that the denominator is still just $\sum k$, and not $\sum k+1$, as you can verify for yourself by checking that Eq. (13.46) is correctly normalized so that $\sum_{k=0}^{\infty} q_k = 1$.)

The distribution q_k is called the *excess degree distribution* and it will come up repeatedly in the sections that follow. It is the probability distribution, for a vertex reached by following an edge, of the number of other edges attached to that vertex.

13.4 CLUSTERING COEFFICIENT

As a simple application of the excess degree distribution, let us calculate the clustering coefficient for the configuration model. Recall that the clustering coefficient is the average probability that two neighbors of a vertex are neighbors of each other.

Consider then a vertex v that has at least two neighbors, which we will denote i and j . Being neighbors of v , i and j are both at the ends of edges from v , and hence the number of other edges connected to them, k_i and k_j are distributed according to the excess degree distribution, Eq. (13.46). The probability of an edge between i and j is then $k_i k_j / 2m$ (see Eq. (13.32)) and, averaging both k_i and k_j over the distribution q_k , we get an expression for the clustering coefficient thus:

$$\begin{aligned} C &= \sum_{k_i, k_j=0}^{\infty} q_{k_i} q_{k_j} \frac{k_i k_j}{2m} = \frac{1}{2m} \left[\sum_{k=0}^{\infty} k q_k \right]^2 \\ &= \frac{1}{2m \langle k \rangle^2} \left[\sum_{k=0}^{\infty} k(k+1) p_{k+1} \right]^2 \\ &= \frac{1}{2m \langle k \rangle^2} \left[\sum_{k=0}^{\infty} (k-1) k p_k \right]^2 \\ &= \frac{1}{n} \frac{[\langle k^2 \rangle - \langle k \rangle]^2}{\langle k \rangle^3}, \end{aligned}$$

(13.47)

where we have made use of $2m = n \langle k \rangle$, Eq. (6.23).

Like the clustering coefficient of the Poisson random graph, Eq. (12.11), this expression goes as n^{-1} for fixed degree distribution, and so vanishes in the limit of large system size. Hence, like the Poisson random graph, the configuration model appears to be an unpromising model for real-world networks with high clustering. Note, however, that Eq. (13.47) contains the second moment $\langle k^2 \rangle$ of the degree distribution in its numerator which can become large, for instance in networks with power-law degree distributions (see Section 8.4.2). This can result in surprisingly large values of C in the configuration model. For further discussion of this point see Section 8.6.

13.5 GENERATING FUNCTIONS FOR DEGREE DISTRIBUTIONS

In the calculations that follow, we will make heavy use of the generating functions for the degree distribution and the excess degree distribution of a network. We will denote these generating functions by $g_0(z)$ and $g_1(z)$ respectively. They are defined by

$$g_0(z) = \sum_{k=0}^{\infty} p_k z^k,$$

(13.48)

$$g_1(z) = \sum_{k=0}^{\infty} q_k z^k.$$

(13.49)

Although it will be convenient to have separate notations for these two commonly occurring functions, they are not really independent, since the excess degree distribution is itself defined in terms of the ordinary degree distribution via Eq. (13.46). Using Eq. (13.46) we can write $g_1(z)$ as

$$\begin{aligned} g_1(z) &= \frac{1}{\langle k \rangle} \sum_{k=0}^{\infty} (k+1) p_{k+1} z^k = \frac{1}{\langle k \rangle} \sum_{k=0}^{\infty} k p_k z^{k-1} \\ &= \frac{1}{\langle k \rangle} \frac{dg_0}{dz}. \end{aligned}$$

(13.50)

But Eq. (13.22) tells us that the average vertex degree is $\langle k \rangle = g'_0(1)$, so

$$g_1(z) = \frac{g'_0(z)}{g'_0(1)}.$$

(13.51)

Thus if we can find $g_0(z)$, we can also find $g_1(z)$ directly from it, without the need to calculate the excess degree distribution explicitly.

For example, suppose our degree distribution is a Poisson distribution with mean c :

$$p_k = e^{-c} \frac{c^k}{k!}.$$

(13.52)

Then its generating function is given by Eq. (13.6) to be

$$g_0(z) = e^{c(z-1)}.$$

(13.53)

Applying Eq. (13.51), we then find that

$$g_1(z) = e^{c(z-1)}.$$

(13.54)

In other words, $g_0(z)$ and $g_1(z)$ are identical in this case. (This is one reason why calculations are relatively straightforward for the Poisson random graph—there is no difference between the degree distribution and the excess degree distribution in that case, a fact you can easily demonstrate for yourself by substituting Eq. (13.52) directly into Eq. (13.46).)

A more complicated example is the power-law distribution, Eq. (13.10), which has a generating function given by Eq. (13.16) to be

$$g_0(z) = \frac{\text{Li}_\alpha(z)}{\zeta(\alpha)},$$

(13.55)

where $\text{Li}_\alpha(z)$ is the polylogarithm function and α is the exponent of the power law. Substituting this result into Eq. (13.51) and making use of Eq. (13.17) gives

$$g_1(z) = \frac{\text{Li}_{\alpha-1}(z)}{z \text{Li}_{\alpha-1}(1)} = \frac{\text{Li}_{\alpha-1}(z)}{z \zeta(\alpha-1)},$$

(13.56)

where we have made use of the fact that $\text{Li}_\alpha(1) = \zeta(\alpha)$ (see Eqs. (13.12) and (13.15)).

13.6 NUMBER OF SECOND NEIGHBORS OF A VERTEX

Armed with these results, we are now in a position to make some more detailed calculations of the properties of the configuration model. The first question we will address is a relatively simple one: what is the probability $p_k^{(2)}$ that a vertex has exactly k second neighbors in the network?

Let us break this probability down by writing it in the form

$$p_k^{(2)} = \sum_{m=0}^{\infty} p_m P^{(2)}(k|m),$$

(13.57)

where $P^{(2)}(k|m)$ is the probability of having k second neighbors given that we have m first neighbors and p_m is the ordinary degree distribution. Equation (13.57) says that the total probability of having k second neighbors is the probability of having k second neighbors given that we have m first neighbors, averaged over all possible values of m . We assume that we are given the degree distribution p_m ; we need to find $P^{(2)}(k|m)$ and then complete the sum.




Figure 13.4: Calculation of the number of second neighbors of a vertex. The number of second neighbors of a vertex (top) is equal to the sum of the excess degrees of the first neighbors.

As illustrated in Fig. 13.4, the number of second neighbors of a vertex is equal to the sum of the excess degrees of the first neighbors. And as discussed in the previous section, the excess degrees are distributed according to the distribution q_k , Eq. (13.46), so that the probability that the excess degrees of our m first neighbors take the values $j_1 \dots j_m$ is $\prod_{r=1}^m q_{j_r}$. Summing over all sets of values $j_1 \dots j_m$, the probability that the excess degrees sum to k and hence that we have k second neighbors is

$$P^{(2)}(k|m) = \sum_{j_1=0}^{\infty} \dots \sum_{j_m=0}^{\infty} \delta(k, \sum_{r=1}^m j_r) \prod_{r=1}^m q_{j_r}.$$

(13.58)

Substituting this expression into (13.57), we find that

$$p_k^{(2)} = \sum_{m=0}^{\infty} p_m \sum_{j_1=0}^{\infty} \cdots \sum_{j_m=0}^{\infty} \delta(k, \sum_{r=1}^m j_r) \prod_{r=1}^m q_{j_r}.$$

(13.59)

By now, you may be starting to find sums of this type familiar. We saw them previously in Eqs. (12.25) and (13.27), for example. We can handle this one by the same trick we used before: instead of trying to calculate $p_k^{(2)}$ directly, we calculate instead its generating function $g^{(2)}(z)$ thus:

$$\begin{aligned} g^{(2)}(z) &= \sum_{k=0}^{\infty} p_k^{(2)} z^k \\ &= \sum_{k=0}^{\infty} z^k \sum_{m=0}^{\infty} p_m \sum_{j_1=0}^{\infty} \cdots \sum_{j_m=0}^{\infty} \delta(k, \sum_{r=1}^m j_r) \prod_{r=1}^m q_{j_r} \\ &= \sum_{m=0}^{\infty} p_m \sum_{j_1=0}^{\infty} \cdots \sum_{j_m=0}^{\infty} z^{\sum_{r=1}^m j_r} \prod_{r=1}^m q_{j_r} \\ &= \sum_{m=0}^{\infty} p_m \sum_{j_1=0}^{\infty} \cdots \sum_{j_m=0}^{\infty} \prod_{r=1}^m q_{j_r} z^{j_r} \\ &= \sum_{m=0}^{\infty} p_m \left[\sum_{j=0}^{\infty} q_j z^j \right]^m. \end{aligned}$$

(13.60)

But now we notice an interesting thing: the sum in square brackets in the last line is none other than the generating function $g_1(z)$ for the excess degree distribution, Eq. (13.49). Thus Eq. (13.60) can be written as

$$g^{(2)}(z) = \sum_{m=0}^{\infty} p_m [g_1(z)]^m = g_0(g_1(z)),$$

(13.61)

where $g_0(z)$ is the generating function for the ordinary degree distribution, defined in Eq. (13.48). So once we know the generating functions for our two basic degree distributions the generating function for the distribution of the second neighbors is very simple to calculate.

In fact, there was no need to go through this lengthy calculation to reach Eq. (13.61). We can derive the same result much more quickly by making use of the “powers” property of generating functions that we derived in Section 13.1.4. There we showed (Eq. (13.28)) that, given a quantity k distributed according to a distribution with generating function $g(z)$, m independent quantities drawn from the same distribution have a sum whose distribution is given by the generating function $[g(z)]^m$. We can apply this result here, by noting that the m excess degrees of the first neighbors of our vertex are just such a set of independent quantities. Given that $g_1(z)$ is the generating function for the distribution of a single one of them (Eq. (13.49)), the distribution $P^{(2)}(k|m)$ of their sum—which is the number of second neighbors—has generating function $[g_1(z)]^m$. That is,

$$\sum_{k=0}^{\infty} P^{(2)}(k|m) z^k = [g_1(z)]^m. \quad (13.62)$$

Now, using Eq. (13.57), the generating function for $p_k^{(2)}$ is

$$\begin{aligned} g^{(2)}(z) &= \sum_{k=0}^{\infty} p_k^{(2)} z^k = \sum_{k=0}^{\infty} \sum_{m=0}^{\infty} p_m P^{(2)}(k|m) z^k \\ &= \sum_{m=0}^{\infty} p_m \sum_{k=0}^{\infty} P^{(2)}(k|m) z^k = \sum_{m=0}^{\infty} p_m [g_1(z)]^m \\ &= g_0(g_1(z)). \end{aligned} \quad (13.63)$$

In future calculations, we will repeatedly make use of this shortcut to get our results, rather than taking the long route exemplified in Eq. (13.60).

We can also use similar methods to calculate the probability distribution of the number of third neighbors. The number of third neighbors is the sum of the excess degrees of each of the second neighbors. Thus, if there are m second neighbors, then the probability distribution $P^{(3)}(k|m)$ of the number of third neighbors has generating function $[g_1(z)]^m$ and the overall probability of having k third neighbors is exactly analogous to Eq. (13.63):

$$\begin{aligned} g^{(3)}(z) &= \sum_{k=0}^{\infty} \sum_{m=0}^{\infty} p_m^{(2)} P^{(3)}(k|m) z^k = \sum_{m=0}^{\infty} p_m^{(2)} \sum_{k=0}^{\infty} P^{(3)}(k|m) z^k \\ &= \sum_{m=0}^{\infty} p_m^{(2)} [g_1(z)]^m = g^{(2)}(g_1(z)) \\ &= g_0(g_1(g_1(z))). \end{aligned} \quad (13.64)$$

Indeed, the generating function for the number of neighbors at any distance d can be expressed this way as

$$\begin{aligned} g^{(d)}(z) &= \sum_{k=0}^{\infty} \sum_{m=0}^{\infty} p_m^{(d-1)} P^{(d)}(k|m) z^k \\ &= \sum_{m=0}^{\infty} p_m^{(d-1)} \sum_{k=0}^{\infty} P^{(d)}(k|m) z^k = \sum_{m=0}^{\infty} p_m^{(d-1)} [g_1(z)]^m \\ &= g^{(d-1)}(g_1(z)). \end{aligned}$$

(13.65)

In other words $g^{(d)}(z) = g_0(g_1(\dots g_1(z)\dots))$, with $d - 1$ copies of g_1 nested inside a single g_0 . This expression is correct at arbitrary distances on an infinite network. On a finite network it will break down if d becomes large enough but will be accurate for small values of d .

These results are all very good, but what use are they? Even given the generating function $g^{(2)}(z)$ it is typically quite difficult to extract explicit probabilities for numbers of second neighbors in the network. For instance, if our degree distribution were Poisson with mean c then $g_0(z) = g_1(z) = e^{c(z-1)}$ as in Eqs. (13.53) and (13.54) and

$$g^{(2)}(z) = e^{c(e^{c(z-1)}-1)}.$$

(13.66)

But to find the actual probabilities we have to apply Eq. (13.2), which involves calculating derivatives of $g^{(2)}(z)$. One can, with a little work, calculate the first few derivatives, but finding a general formula for the n th derivative is hard.¹⁹⁶

What we can do, however, is calculate the average number of neighbors at distance d . The average of a distribution is given by the first derivative of its generating function evaluated at $z = 1$ (see Eq. (13.22)) and the derivative of Eq. (13.63) is

$$\frac{dg^{(2)}}{dz} = g'_0(g_1(z)) g'_1(z).$$

(13.67)

Setting $z = 1$ and recalling that $g_1(1) = 1$ (Eq. (13.20)), we find that the average number c_2 of second neighbors is

$$c_2 = g'_0(1)g'_1(1).$$

(13.68)

But $g'_0(1) = \langle k \rangle$ and

$$\begin{aligned} g'_1(1) &= \sum_{k=0}^{\infty} k q_k \\ &= \frac{1}{\langle k \rangle} \sum_{k=0}^{\infty} k(k+1) p_{k+1} = \frac{1}{\langle k \rangle} \sum_{k=0}^{\infty} (k-1) k p_k \\ &= \frac{1}{\langle k \rangle} (\langle k^2 \rangle - \langle k \rangle). \end{aligned}$$

(13.69)

where we have used Eq. (13.46). Thus the mean number of second neighbors can also be written

$$c_2 = \langle k^2 \rangle - \langle k \rangle.$$

(13.70)

We can take this approach further and calculate the mean number c_d of neighbors at any distance d . Differentiating Eq. (13.65) we get

$$\frac{d g^{(d)}}{dz} = g^{(d-1)'}(g_1(z)) g'_1(z),$$

(13.71)

and setting $z = 1$ we get

$$c_d = g^{(d-1)'}(1) g'_1(1) = c_{d-1} g'_1(1).$$

(13.72)

Making use of Eq. (13.68) to write $g'_1(1) = c_2/c_1$ where $c_1 = \langle k \rangle$, this can be expressed in the simple form

$$c_d = c_{d-1} \frac{c_2}{c_1},$$

(13.73)

which implies that

$$c_d = \left(\frac{c_2}{c_1} \right)^{d-1} c_1.$$

(13.74)

In other words, once we know the mean numbers of first and second neighbors, c_1 and c_2 , we know everything. What's more, the average number of neighbors at distance d either grows or falls off exponentially, depending on whether c_2 is greater or less than c_1 . This observation is strongly reminiscent of the argument we made in Section 12.5 for the appearance of a giant component in a random graph. There we argued that if the number of vertices you can reach within a certain distance is increasing with that distance (on average) then you must have a giant component in the network, while if it is decreasing there can be no giant component. Applying the same reasoning here, we conclude that the configuration model has a giant component if and only if we have

$$c_2 > c_1.$$

(13.75)

Using Eq. (13.70) for c_2 and putting $c_1 = \langle k \rangle$, we can also write this condition as $\langle k^2 \rangle - \langle k \rangle > \langle k \rangle$ or

$$\langle k^2 \rangle - 2\langle k \rangle > 0.$$

(13.76)

This condition for the existence of a giant component in the configuration model was first given by Molloy and Reed [224] in 1995.[197](#)

13.7 GENERATING FUNCTIONS FOR THE SMALL COMPONENTS

In this section and the following one we examine the sizes of components in the configuration model. As we will see, the situation is qualitatively similar to that for the Poisson random graph in that a configuration model network generally has at most one giant component, plus a large number of small components. We will approach the calculation of component sizes by a route different from the one we took for the Poisson random graph and examine first the properties of the small components. We will see that it is possible to calculate the distribution of the sizes of the small components by a method similar to the one we used in the Poisson case. Then we can use these results to get at the properties of the giant component: once we have the sizes of the small components, we can subtract them from the size of the graph as a whole and whatever is left, if anything, must be the giant component.

Let π_s be the probability that a randomly chosen vertex belongs to a small (non-giant) component of size s . We will calculate π_s by first calculating its generating function

$$h_0(z) = \sum_{s=1}^{\infty} \pi_s z^s.$$

(13.77)

Note that the minimum value of s is 1, since every vertex belongs to a component of size at least one (namely itself).

By an argument exactly analogous to that of Section 12.6.1 we can show that the small components in the configuration model are trees (in the limit of large n , provided the degree distribution is held constant as we go to the limit). We can use this fact to derive an expression for the distribution of small component sizes as follows.

Consider Fig. 13.5 (which is actually the same as the figure for the Poisson random graph in the previous chapter (Fig. 12.3), but it works just as well as an illustration of the configuration model). If vertex i is a member of a small component then that component is necessarily a tree. Just as in the Poisson case, this implies that the sets of vertices reachable along each of its edges (shaded areas in Fig. 13.5a) are not connected, other than via vertex i , since if they were connected there would be a loop in the component and hence it would not be a tree.

Now, taking a hint from our argument in the Poisson case, let us remove vertex i from the network along with all its edges—see Fig. 13.5b. The shaded areas in the figure are now not connected to one another at all and hence are each now separate components in their own right. And the size of the component to which vertex i belongs on the original network is equal to the sum of the sizes of these new components, plus one for vertex i itself.

A crucial point to notice, however, is that the neighbors $n_1, n_2, \dots$ of vertex i are, by definition, reached by following an edge. Hence, as we have discussed, these are not typical network vertices, being more likely to have high degree than the typical vertex. Thus the components that they belong to in Fig. 13.5b—the shaded regions in the figure—are not distributed according to π_s . Instead they must have some other distribution. Let us denote this distribution by ρ_s . More specifically, let ρ_s be the probability that the vertex at the end of an edge belongs to a small

component of size s after that edge is removed. Let us also define the generating function for this distribution to be

$$h_1(z) = \sum_{s=0}^{\infty} \rho_s z^s.$$

(13.78)




Figure 13.5: The size of one of the small components in the configuration model. (a) The size of the component to which a vertex i belongs is the sum of the number of vertices in each of the subcomponents (shaded regions) reachable via i 's neighbors n_1 , n_2 , n_3 , plus one for i itself. (b) If vertex i is removed the subcomponents become components in their own right.

We don't yet know the value of ρ_s or its generating function and we will have to calculate them later, but for the moment let us proceed with the information we have.

Suppose that vertex i on the original network has degree k and let us denote by $P(s|k)$ the probability that, after i is removed, its k neighbors belong to small components of sizes summing to exactly s . Alternatively, $P(s-1|k)$ is the probability that i itself belongs to a small component of size s given that its degree is k . Then the total probability π_s that i belongs to a small component of size s is this probability averaged over k thus:

$$\pi_s = \sum_{k=0}^{\infty} p_k P(s-1|k).$$

(13.79)

Substituting this expression into Eq. (13.77) we then get an expression for the generating function for π_s as follows:

$$\begin{aligned}
h_0(z) &= \sum_{s=1}^{\infty} \sum_{k=0}^{\infty} p_k P(s-1|k) z^s = z \sum_{k=0}^{\infty} p_k \sum_{s=1}^{\infty} P(s-1|k) z^{s-1} \\
&= z \sum_{k=0}^{\infty} p_k \sum_{s=0}^{\infty} P(s|k) z^s.
\end{aligned}$$

(13.80)

The final sum in this expression is the generating function for the probability that the k neighbors belong to small components whose size sums to s . But the sizes of the small components are independent of one another and hence we can use the “powers” property of generating functions (Section 13.1.4), which tells us that the generating function we want is just equal to the generating function for the size of the component any single neighbor belongs to—the function that we denoted $h_1(z)$ above—raised to the k th power. Thus

$$h_0(z) = z \sum_{k=0}^{\infty} p_k [h_1(z)]^k = z g_0(h_1(z)).$$

(13.81)

We still don’t know the generating function $h_1(z)$ but we can derive it now quite easily. We consider the network in which vertex i is removed and ask what is the probability ρ_s that one of the neighbors of i belongs to a component of size s in this network. In the limit of large network size, the removal of the single vertex i will have no effect on the degree distribution, so the network still has the same distribution as before, which means that if the neighbor has degree k then its probability of belonging to a component of size s is $P(s-1|k)$, just as before. Note, however, that the degree k does not follow the ordinary degree distribution. Since the neighbor was reached by following an edge from i , its degree, discounting the edge to i that has been removed, follows the excess degree distribution q_k defined in Eq. (13.46), rather than the ordinary degree distribution. Thus

$$\rho_s = \sum_{k=0}^{\infty} q_k P(s-1|k),$$

(13.82)

and, substituting this expression into Eq. (13.78), we have

$$h_1(z) = \sum_{s=1}^{\infty} \sum_{k=0}^{\infty} q_k P(s-1|k) z^s = z \sum_{k=0}^{\infty} q_k \sum_{s=0}^{\infty} P(s|k) z^s.$$

(13.83)

As before, the last sum is the generating function for $P(s|k)$, which is equal to $[h_1(z)]^k$, and hence

$$h_1(z) = z \sum_{k=0}^{\infty} q_k [h_1(z)]^k = z g_1(h_1(z)).$$

(13.84)

Collecting together our results, the generating functions for π_s and ρ_s thus satisfy

$$h_0(z) = z g_0(h_1(z)),$$

(13.85)

$$h_1(z) = z g_1(h_1(z)).$$

(13.86)

If we can solve the second of these equations for $h_1(z)$ then we can substitute the result into the first equation and we have our answer for $h_0(z)$. In practice, it is often not easy to solve for $h_1(z)$, and, even if it is, extracting the actual component size distribution from the generating function can be difficult. But that does not mean that these results are useless. On the contrary, there are many useful things we can deduce from them. One important quantity we can calculate is the size of the giant component.

13.8 GIANT COMPONENT

Given the definition $h_0(z) = \sum_s \pi_s z^s$, where π_s is the probability that a randomly chosen vertex belongs to a small component of size s , we have $h_0(1) = \sum_s \pi_s$, which is the total probability that a randomly chosen vertex belongs to a small component. Unlike most generating functions, it is not necessarily the case that $h(1) = 1$ because there may be a giant component in the network. If there is a giant component then some of the vertices do not belong to any small component and $\sum_s \pi_s$ will be less than 1. In fact, $\sum_s \pi_s$ will be simply the fraction of vertices that belong to small components and hence the fraction S of vertices belonging to the giant component is

$$S = 1 - \sum_{s=0}^{\infty} \pi_s = 1 - h_0(1) = 1 - g_0(h_1(1)),$$

(13.87)

where we have used Eq. (13.85). The value of $h_1(1)$ we can get from Eq. (13.86):

$$h_1(1) = g_1(h_1(1)).$$

(13.88)

The quantity $h_1(1)$ will occur frequently in subsequent developments, so for convenience let us define the shorthand notation

$$u = h_1(1),$$

(13.89)

in which case Eqs. (13.87) and (13.88) can be written

$$S = 1 - g_0(u),$$

(13.90)

$$u = g_1(u).$$

(13.91)

In other words, u is a fixed point of the function $g_1(z)$ —a point where the function is equal to its own argument—and if we can find this fixed point then we need only substitute the result into Eq. (13.90) and we have the size of the giant component.

Since $g_1(1) = 1$ (see Eq. (13.20) and the discussion that precedes it), there is always a fixed point of g_1 at $u = 1$, but this solution gives $S = 1 - g_0(1) = 0$ and hence no giant component. If there is to be a giant component there must be at least one other non-trivial solution to Eq. (13.91). We will see some examples of such solutions shortly.

The quantity $u = h_1(1)$ has a simple physical interpretation. Recall that $h_1(z) = \sum_s \rho_s z^s$ is the generating function for the probability ρ_s that the vertex reached by following an edge belongs to a small component of size s if that edge is removed. Thus $h_1(1) = \sum_s \rho_s$ is the total probability that such a vertex belongs to a small component of any size, or equivalently the probability that it doesn't belong to the giant component.

This observation suggests an alternative and simpler derivation of Eqs. (13.90) and (13.91) for the size of a giant component, as follows. To belong to the giant component, a vertex A must be connected to the giant component via at least one of its neighbors. Or equivalently, A does not belong to the giant component if (and only if) it is not connected to the giant component via any of its neighbors. Let us *define* u to be the average probability that a vertex is not connected to the giant component via its connection to some particular neighboring vertex. If vertex A has k neighbors, then the probability that it is not connected to the giant component via any of them is thus u^k . And the average of this probability over the whole network is $\sum_k p_k u^k = g_0(u)$, which is the average probability that a vertex is not in the giant component. But this probability is also, by definition, equal to $1 - S$, where S is the fraction of the graph occupied by the giant component and hence $1 - S = g_0(u)$ or

$$S = 1 - g_0(u),$$

(13.92)

which is Eq. (13.90) again.

Now let us ask what the value of u is. The probability that you are not connected to the giant component via a particular neighboring vertex is equal to the probability that *that* vertex is not connected to the giant component via any of its other neighbors. If there are k of those other neighbors, then that probability is again u^k . But because we are talking about a neighboring vertex, k is now distributed according to the excess degree distribution q_k , Eq. (13.46), and hence taking the average, we find that $u = \sum_k q_k u^k$ or

$$u = g_1(u),$$

(13.93)

which is Eq. (13.91) again. Thus we have rederived our two equations for the size of the giant component, but by a much shorter route. The main disadvantage of this method is that it only gives the size of the giant component and not the complete generating function for all the other components as well, and this is the reason why we took the time to go through the longer derivation. There are many further results we can derive by knowing the entire generating function, as we show in the next section.

13.8.1 EXAMPLE

Let's take a look at a concrete example and see how calculations for the configuration model work out in practice. Consider a network like that of the first example in Section 13.1.1 that has vertices only of degree 0, 1, 2 and 3, and no vertices of higher degree. Then the generating functions $g_0(z)$ and $g_1(z)$ take the form

$$g_0(z) = p_0 + p_1z + p_2z^2 + p_3z^3,$$

(13.94)

$$\begin{aligned} g_1(z) &= \frac{g'_0(z)}{g'_0(1)} = \frac{p_1 + 2p_2z + 3p_3z^2}{p_1 + 2p_2 + 3p_3} \\ &= q_0 + q_1z + q_2z^2. \end{aligned}$$

(13.95)

Equation (13.91) is thus quadratic in this case, $u = q_0 + q_1u + q_2u^2$, which has the solutions

$$u = \frac{1 - q_1 \pm \sqrt{(1 - q_1)^2 - 4q_0q_2}}{2q_2}.$$

(13.96)

However, we know that $\sum_k q_k = 1$, and hence in this case $1 - q_1 = q_0 + q_2$. Using this result to eliminate q_1 we get

$$\begin{aligned} u &= \frac{(q_0 + q_2) \pm \sqrt{(q_0 + q_2)^2 - 4q_0q_2}}{2q_2} \\ &= \frac{(q_0 + q_2) \pm (q_0 - q_2)}{2q_2} \\ &= 1 \quad \text{or} \quad \frac{q_0}{q_2}. \end{aligned}$$

(13.97)

Thus, as expected we have a solution $u = 1$, but we also have another non-trivial solution which *might* imply that we have a giant component.

If $q_2 < q_0$ then this non-trivial solution gives $u > 1$. Since u is a probability it cannot be greater than 1, so in this case we definitely do not have a giant component. On the other hand, if $q_2 > q_0$ we have a viable non-trivial solution $u < 1$ equal to

$$u = \frac{q_0}{q_2} = \frac{p_1}{3p_3},$$

(13.98)

where we have extracted values of q_0 and q_2 from Eq. (13.95). We can also write the condition $q_2 > q_0$ in terms of the p_k as

$$p_3 > \frac{1}{3}p_1.$$

(13.99)

In other words, there can be a giant component if the number of vertices of degree three exceeds one third the number of degree one. This is a remarkable result. It says that the number of vertices of degree zero and degree two don't matter at all (except to the extent that their absence makes room for more vertices of the other degrees). As we will see, this is actually a general result—the values of p_0 and p_2 never make any difference to the presence or absence of a giant component. On the other hand, the size of the giant component for the current example is given by Eq. (13.90) to be

$$S = 1 - g_0(u) = 1 - p_0 - \frac{p_1^2}{3p_3} - \frac{p_1^2 p_2}{9p_3^2} - \frac{p_1^3}{27p_3^2}.$$

(13.100)

Thus the size of the giant component does depend on p_0 and p_2 , even though its presence or absence does not.

We have not, however, yet proved that a giant component actually does exist. In the regime where we have two solutions for u , one with $u = 1$ (no giant component) and one with $u < 1$ (there is a giant component) it is unclear which of these solutions we should believe. In Section 13.6, however, we showed that there is a giant component in the network when the degree sequence

satisfies a specific condition, Eq. (13.76). In the next section, we show that in fact this condition is always satisfied whenever a non-trivial solution $u < 1$ exists, and hence that there is always a giant component when we have such a solution.

13.8.2 GRAPHICAL SOLUTIONS AND THE EXISTENCE OF THE GIANT COMPONENT

The example given in the last section is unusual in that we can solve the fixed-point equation (13.91) exactly for the crucial parameter u . In most other cases exact solutions are not possible, but we can nonetheless get a good idea of the behavior of u by graphical means. The derivatives of $g_1(z)$ are proportional to the probabilities ρ_s and hence are all non-negative. That means that for $z \geq 0$, $g_1(z)$ is in general positive, an increasing function of its argument, and upward concave. It also takes the value 1 when $z = 1$. Thus it must look qualitatively like one of the curves in Fig. 13.6. The solution of the fixed-point equation $u = g_1(u)$ is then given by the intercept of the curve $y = g_1(u)$ with the line $y = u$ (the dotted line in the figure).




Figure 13.6: Graphical solution of Eq. (13.91). The solution of the equation $u = g_1(u)$ is given by the point at which the curve $y = g_1(u)$ intercepts the line $y = u$.

As we already know, there is always a trivial solution at $u = 1$ (top right in the figure). But now we can see that there can be just one other solution with $u < 1$ and only if the curve takes the right form. In particular, we have a non-trivial solution at $u < 1$ if the slope $g'_1(1)$ of the curve at $u = 1$ is greater than the slope of the dotted line. That is, if

$$g'_1(1) > 1.$$

(13.101)

Using Eq. (13.49) for $g_1(z)$, we have

$$\begin{aligned} g'_1(1) &= \sum_{k=0}^{\infty} k q_k = \frac{1}{\langle k \rangle} \sum_{k=0}^{\infty} k(k+1) p_k = \frac{1}{\langle k \rangle} \sum_{k=0}^{\infty} (k-1) k p_k \\ &= \frac{\langle k^2 \rangle - \langle k \rangle}{\langle k \rangle}. \end{aligned}$$

(13.102)

Thus our condition for the solution at $u < 1$ is

$$\frac{\langle k^2 \rangle - \langle k \rangle}{\langle k \rangle} > 1.$$

(13.103)

or equivalently,

$$\langle k^2 \rangle - 2\langle k \rangle > 0,$$

(13.104)

But this is none other than the condition for the existence of a giant component, Eq. (13.76). In other words, the conditions for the existence of a giant component and the existence of the non-trivial solution to Eq. (13.91) are exactly the same and hence, as promised, there is always a giant component whenever a solution $u < 1$ exists for Eq. (13.91).

Writing $\langle k \rangle = n^{-1} \sum_i k_i$ and $\langle k^2 \rangle = n^{-1} \sum_i k_i^2$, we can also write Eq. (13.104) as

$$\sum_i k_i(k_i - 2) > 0.$$

(13.105)

But note now that, as before, vertices of degree zero and degree two make no contribution to the sum, since terms in which $k_i = 0$ or $k_i = 2$ vanish. Thus we can add as many vertices of degree zero or two to the network as we like (or take them away) and it will make no difference to the existence or not of a giant component. We noted a special case of this phenomenon in Section

13.8.1.

13.9 SIZE DISTRIBUTION FOR SMALL COMPONENTS

Having looked in some detail at the behavior of the giant component in the configuration model, let us return once more to the small components. In Eqs. (13.85) and (13.86) we have—in theory at least—the generating functions that give the entire distribution of sizes of the small components. Unfortunately, it is in most cases impossible to solve these equations exactly, but we can still extract plenty of useful information from them. For example, we can calculate the mean size of the component to which a randomly chosen vertex belongs, which is given by the equivalent of Eq. (12.29) thus:

$$\langle s \rangle = \frac{\sum_s s \pi_s}{\sum_s \pi_s} = \frac{h'_0(1)}{1-S} = \frac{h'_0(1)}{g_0(u)},$$

(13.106)

where we have used Eq. (13.90) in the final equality. Differentiating Eq. (13.85) we get

$$\begin{aligned} h'_0(z) &= g_0(h_1(z)) + z g'_0(h_1(z)) h'_1(z) \\ &= g_0(h_1(z)) + z g'_0(1) g_1(h_1(z)) h'_1(z) \\ &= \frac{h_0(z)}{z} + g'_0(1) h_1(z) h'_1(z), \end{aligned}$$

(13.107)

where we have used Eq. (13.51) in the second equality and Eqs. (13.85) and (13.86) in the third. Setting $z = 1$ we then get

$$h'_0(1) = h_0(1) + g'_0(1) h_1(1) h'_1(1) = 1 - S + g'_0(1) h'_1(1) u,$$

(13.108)

where we have used Eqs. (13.87) and (13.89). To calculate $h'_1(1)$ we differentiate Eq. (13.86) thus:

$$\begin{aligned} h'_1(z) &= g_1(h_1(z)) + z g'_1(h_1(z)) h'_1(z) \\ &= \frac{h_1(z)}{z} + z g'_1(h_1(z)) h'_1(z), \end{aligned}$$

(13.109)

or, rearranging,

$$h_1'(z) = \frac{h_1(z)/z}{1 - zg_1'(h_1(z))}.$$

(13.110)

Setting $z = 1$ in this expression gives

$$h_1'(1) = \frac{u}{1 - g_1'(u)}.$$

(13.111)

Combining Eqs. (13.106), (13.108), and (13.111), we then find that

$$\langle s \rangle = 1 + \frac{g_0'(1)u^2}{g_0(u)[1 - g_1'(u)]}.$$

(13.112)

Using values of S and u from Eqs. (13.90) and (13.91) we can then calculate $\langle s \rangle$ from this equation.

A simple case occurs when we are in the region where there is no giant component. In this region we have $S = 0$ and $u = 1$ by definition and hence

$$\langle s \rangle = 1 + \frac{g_0'(1)}{1 - g_1'(1)}.$$

(13.113)

Thus the average size of the component to which a vertex belongs diverges precisely at the point where $g_1'(1) = 1$, the point at which the curve in Fig. 13.6 is exactly tangent to the dotted line (the

middle curve in the figure). This is, of course, also the point at which the giant component first appears.

Thus the picture we have is similar to that shown in Fig. 12.4 for the Poisson random graph, in which the typical size of the component to which a vertex belongs grows larger and larger until we reach the point, or *phase transition*, where the giant component appears, at which it diverges. Beyond this point the small components shrink in size again, although the overall mean component size, including the giant component, is infinite.

Equation (13.113) can also be expressed in a couple of other forms that may be useful in some circumstances. From Eq. (13.69) we know that $g'_1(1) = (\langle k^2 \rangle - \langle k \rangle) / \langle k \rangle$ and, putting $g'_0(1) = \langle k \rangle$ also, we find that

$$\langle s \rangle = 1 + \frac{\langle k \rangle^2}{2\langle k \rangle - \langle k^2 \rangle}.$$

(13.114)

This expression can be evaluated easily given only a knowledge of the degree sequence and avoids the need to calculate any generating functions. Using the notation introduced earlier in which c_1 and c_2 are the mean number of first and second neighbors of a vertex, with c_2 given by Eq. (13.70), we can also write (13.114) in the form

$$\langle s \rangle = 1 + \frac{c_1^2}{c_1 - c_2},$$

(13.115)

so that the average size of the component a vertex belongs to is dictated entirely by the mean numbers of first and second neighbors.

13.9.1 AVERAGE SIZE OF A SMALL COMPONENT

As with the Poisson random graph, we must be careful about our claims in the previous section. We have calculated the average size $\langle s \rangle$ of the component to which a randomly chosen vertex belongs but this is not the same thing as the average size of a component, since more vertices belong to larger components, which biases the value of $\langle s \rangle$. If we want the true average size R of the small components, we must use Eq. (12.36), which we reproduce here for convenience:

$$R = \frac{1 - S}{\sum_s \pi_s / s}.$$

(13.116)

The sum can be calculated as before using the equivalent of Eq. (12.37):

$$\sum_{s=1}^{\infty} \frac{\pi_s}{s} = \int_0^1 \frac{h_0(z)}{z} dz.$$

(13.117)

Taking $h_0(z)/z$ from Eq. (13.107), we get

$$\begin{aligned} \sum_{s=1}^{\infty} \frac{\pi_s}{s} &= \int_0^1 \frac{dh_0}{dz} dz - g'_0(1) \int_0^1 h_1(z) \frac{dh_1}{dz} dz \\ &= \int_0^{1-S} dh_0 - \langle k \rangle \int_0^u h_1 dh_1 \\ &= 1 - S - \frac{1}{2} \langle k \rangle u^2. \end{aligned}$$

(13.118)

Then

$$R = \frac{2}{2 - \langle k \rangle u^2 / (1 - S)}.$$

(13.119)

Note that the value of this average at the transition point where $S = 0$ and $u = 1$ is just $2/(2 - \kappa)$, which is normally perfectly finite.¹⁹⁸ Thus the average component size does not normally diverge at the transition (unlike s).

13.9.2 COMPLETE DISTRIBUTION OF SMALL COMPONENT SIZES

One of the most surprising results concerning the configuration model is that it is possible to derive an expression not just for the average size of the component to which a vertex belongs, but for the exact probability that it belongs to a component of any specific size—the probability that it belongs to a component of size ten, or a hundred, or a million. The derivation of this result is similar to the derivation given in Section 12.6.3 for the corresponding quantity for the Poisson random graph.

Since a component cannot have size zero, the generating function for the probabilities π_s has the form

$$h_0(z) = \sum_{s=1}^{\infty} \pi_s z^s,$$

(13.120)

with the sum starting at 1. Dividing by z and differentiating $s - 1$ times, we then find that

$$\pi_s = \frac{1}{(s-1)!} \left[\frac{d^{s-1}}{dz^{s-1}} \left(\frac{h_0(z)}{z} \right) \right]_{z=0},$$

(13.121)

(which is just a minor variation on the standard formula, Eq. (13.2)). Using Eq. (13.85), this can also be written

$$\begin{aligned} \pi_s &= \frac{1}{(s-1)!} \left[\frac{d^{s-1}}{dz^{s-1}} g_0(h_1(z)) \right]_{z=0} \\ &= \frac{1}{(s-1)!} \left[\frac{d^{s-2}}{dz^{s-2}} [g'_0(h_1(z)) h'_1(z)] \right]_{z=0}. \end{aligned}$$

(13.122)

Now we make use of the Cauchy formula for the n derivative of a function, which says that

$$\left. \frac{d^n f}{dz^n} \right|_{z=z_0} = \frac{n!}{2\pi i} \oint \frac{f(z)}{(z-z_0)^{n+1}} dz,$$

(13.123)

where the integral is around a contour that encloses z_0 in the complex plane but encloses no poles in $f(z)$. Applying this formula to Eq. (13.122) with $z_0 = 0$ we get

$$\pi_s = \frac{1}{2\pi i(s-1)} \oint \frac{g'_0(h_1(z))}{z^{s-1}} \frac{dh_1}{dz} dz.$$

(13.124)

For our contour, we choose an infinitesimal circle around the origin. Changing the integration variable to h_1 , we can also write this as

$$\pi_s = \frac{1}{2\pi i(s-1)} \oint \frac{g'_0(h_1)}{z^{s-1}} dh_1.$$

(13.125)

Here we are regarding z now as a function of h_1 , rather than the other way around. Furthermore, since $h_1(z)$ goes to zero as $z \rightarrow 0$, the contour in h_1 surrounds the origin too. (The proof is the same as for Eq. (12.46).)

Now we make use of Eq. (13.86) to eliminate z and write

$$\begin{aligned} \pi_s &= \frac{1}{2\pi i(s-1)} \oint \frac{[g_1(h_1)]^{s-1} g'_0(h_1)}{h_1^{s-1}} dh_1 \\ &= \frac{g'_0(1)}{2\pi i(s-1)} \oint \frac{[g_1(h_1)]^s}{h_1^{s-1}} dh_1, \end{aligned}$$

(13.126)

where we have made use of Eq. (13.51) in the second line. Given that the contour surrounds the origin, this integral is now in the form of Eq. (13.123) again, and hence

$$\pi_s = \frac{\langle k \rangle}{(s-1)!} \left[\frac{d^{s-2}}{dz^{s-2}} [g_1(z)]^s \right]_{z=0},$$

(13.127)

where we have written $g'_0(1) = \langle k \rangle$.

The only exception to this formula is for the case $s = 1$, for which Eq. (13.124) gives 0/0 and is therefore clearly incorrect. However, since the only way to belong to a component of size 1 is to have no connections to any other vertices, the probability π_1 is trivially equal to the probability of having degree zero:

$$\pi_1 = p_0.$$

(13.128)

Equations (13.127) and (13.128) give the probability that a randomly chosen vertex belongs to a component of size s in terms of the degree distribution. In principle if we know p_k we can calculate π_s . It is not always easy to perform the derivatives in practice and in some cases we may not even know the generating function $g_1(z)$ in closed form, but at least in some cases the calculations are possible. As an example, consider a network with the exponential degree distribution

$$p_k = (1 - e^{-\lambda})e^{-\lambda k},$$

(13.129)

with exponential parameter $\lambda > 0$. From Eqs. (13.9) and (13.51) the generating functions $g_0(z)$ and $g_1(z)$ are given by

$$g_0(z) = \frac{e^\lambda - 1}{e^\lambda - z}, \quad g_1(z) = \left(\frac{e^\lambda - 1}{e^\lambda - z} \right)^2.$$

(13.130)

Then it is not hard to show that

$$\frac{d^n}{dz^n} [g_1(z)]^s = \frac{(2s-1+n)!}{(2s-1)!} \frac{[g_1(z)]^s}{(e^\lambda - z)^n},$$

(13.131)

and hence

$$\pi_s = \frac{(3s-3)!}{(s-1)!(2s-1)!} e^{-\lambda(s-1)} (1 - e^{-\lambda})^{2s-1}.$$

(13.132)




Figure 13.7: The distribution of component sizes in a configuration model. The probability π_s that a vertex belongs to a component of size s for the configuration model with an exponential degree distribution of the form (13.129) for $\lambda = 1.2$. The solid lines represent the exact formula, Eq. (13.132), for the $n \rightarrow \infty$ limit and the points are measurements of π_s averaged over 100 computer-generated networks with $n = 10^7$ vertices each.

Figure 13.7 shows a comparison of this formula with the results of numerical simulations for $\lambda = 1.2$ and, as we can see, the agreement between formula and simulations is good—our calculations seem to describe the simulated random graph well even though the graph is necessarily finite in size while the calculations are performed in the limit of large n .

13.10 POWER-LAW DEGREE DISTRIBUTIONS

As we saw in Section 8.4, a number of networks have degree distributions that approximately obey a power law. As an example of the application of the machinery developed in this chapter, let us look at the properties of a random graph with a power-law degree distribution.

Suppose we have a network with a “pure” power-law degree distribution of the form

$$p_k = \begin{cases} 0 & \text{for } k = 0, \\ k^{-\alpha} / \zeta(\alpha) & \text{for } k \geq 1. \end{cases}$$

(13.133)

(See Eq. (13.13).) Here $\alpha > 0$ is a constant exponent and $\zeta(\alpha)$ is the Riemann zeta function:

$$\zeta(\alpha) = \sum_{k=1}^{\infty} k^{-\alpha}.$$

(13.134)

Using the results of the previous sections we can, for instance, say whether there is a giant component in this network or not. Equation (13.76) tells us that there will be a giant component if and only if

$$\langle k^2 \rangle - 2\langle k \rangle > 0.$$

(13.135)

In the present case

$$\langle k \rangle = \sum_{k=0}^{\infty} k p_k = \frac{1}{\zeta(\alpha)} \sum_{k=1}^{\infty} k^{-\alpha+1} = \frac{\zeta(\alpha-1)}{\zeta(\alpha)},$$

(13.136)

and

$$\langle k^2 \rangle = \sum_{k=0}^{\infty} k^2 p_k = \frac{1}{\zeta(\alpha)} \sum_{k=1}^{\infty} k^{-\alpha+2} = \frac{\zeta(\alpha-2)}{\zeta(\alpha)}.$$

(13.137)

Thus there is a giant component if

$$\zeta(\alpha-2) > 2\zeta(\alpha-1).$$

(13.138)

Figure 13.8 shows this inequality in graphical form. The two curves in the figure show the values of $\zeta(\alpha-2)$ and $2\zeta(\alpha-1)$ as functions of α and, as we can see, the inequality (13.138) is satisfied only for sufficiently low values of α , below the dotted line in the figure. In fact a numerical solution of the equation $\zeta(\alpha-2) = 2\zeta(\alpha-1)$ indicates that the network will have a giant component only for $\alpha < 3.4788 \dots$, a result first given by Aiello *et al.* [9] in 2000.

In practice this result is of only limited utility because it applies only for the pure power law. In general, other distributions with power-law tails but different behavior for low k will have different thresholds at which the giant component appears. There is however a general result we can derive that applies to all distributions with power-law tails. In Section 8.4.2 we noted that the second moment $\langle k^2 \rangle$ diverges for any distribution with a power-law tail with exponent $\alpha \leq 3$, while the first moment $\langle k \rangle$ remains finite so long as $\alpha > 2$. This means that Eq. (13.135) is always satisfied for any configuration model with a power-law tail to its degree distribution so long as α lies in the range $2 < \alpha \leq 3$, and hence there will always be a giant component no matter what else the distribution does. For $\alpha > 3$, on the other hand, there may or may not be a giant component, depending on the precise functional form of the degree distribution. (For $\alpha \leq 2$ it turns out that there is always a giant component, although more work is needed to demonstrate this.) Note that, as discussed in Section 8.4, most observed values of α for real-world networks lie in the range $2 < \alpha \leq 3$ and hence we tentatively expect such networks to have a giant component, although we must also bear in mind that the configuration model is a simplified model of a network and is not necessarily a good representation of any specific real-world network.




Figure 13.8: Graphical solution of Eq. (13.138). The configuration model with a pure power-law degree distribution (Eq. (13.133)) has a giant component if $\zeta(\alpha - 2) > 2\zeta(\alpha - 1)$. This happens for values of α below the crossing point of the two curves.

Returning to the pure power law let us calculate the size S of the giant component, when there is one. The fundamental generating functions $g_0(z)$ and $g_1(z)$ for the power-law distribution are given by Eqs. (13.55) and (13.56), which we repeat here for convenience:

$$g_0(z) = \frac{\text{Li}_\alpha(z)}{\zeta(\alpha)}, \quad g_1(z) = \frac{\text{Li}_{\alpha-1}(z)}{z\zeta(\alpha-1)}.$$

(13.139)

Here $\zeta(\alpha)$ is the Riemann zeta function again and $\text{Li}_\alpha(z)$ is the polylogarithm

$$\text{Li}_\alpha(z) = \sum_{k=1}^{\infty} k^{-\alpha} z^k.$$

(13.140)

(See Eq. (13.15).) Now the crucial equation (13.91) for the probability $u = h_1(1)$ reads

$$u = \frac{\text{Li}_{\alpha-1}(u)}{u\zeta(\alpha-1)} = \frac{\sum_{k=1}^{\infty} k^{-\alpha+1} u^k}{u\zeta(\alpha-1)} = \frac{\sum_{k=0}^{\infty} (k+1)^{-\alpha+1} u^k}{\zeta(\alpha-1)},$$

(13.141)

where we have used the explicit definition of the polylogarithm for clarity.

In general there is no closed-form solution for this equation, but we do notice some interesting points. In particular, note that the sum in the numerator is strictly positive for $u \geq 0$, which means that if $\zeta(\alpha - 1)$ diverges we will get a solution $u = 0$. And indeed $\zeta(\alpha - 1)$ does diverge. It diverges at $\alpha = 2$ and all values below, as one can readily verify from the definition, Eq. (13.134).¹⁹⁹ Thus for $\alpha \leq 2$ we have $u = 0$ and Eq. (13.90) then tells us that the giant component has size $S = 1 - g_0(0) = 1 - p_0$. However, for our particular choice of degree distribution, Eq. (13.133), there are no vertices with degree zero, and hence $p_0 = 0$ and $S = 1$. That is, the giant component fills the entire network and there are no small components at all!

Technically, this statement is not quite correct. There is always some chance that, for instance, a vertex of degree 1 will connect to another vertex of degree 1, forming a small component. What we have shown is that the probability that a randomly chosen vertex belongs to a small component is zero in the limit of large n , i.e., that what small components there are fill a fraction of the network that vanishes as $n \rightarrow \infty$. In the language used by mathematicians, a randomly chosen vertex “almost surely” belongs to the giant component, meaning it is technically possible to observe another outcome, but the probability is vanishingly small.

Thus our picture of the pure power-law configuration model is one in which there is a giant component for values of $\alpha < 3.4788 \dots$ and that giant component fills essentially the entire network when $\alpha \leq 2$. In the region between $\alpha = 2$ and $\alpha = 3.4788$ there is a giant component but it does not fill the whole network and some portion of the network consists of small component. If $\alpha > 3.4788 \dots$ there are only small components. As a confirmation of this picture, Fig. 13.9 shows the size of the giant component extracted from a numerical solution of Eq. (13.141).²⁰⁰ As we can see it fits nicely with the picture described above.

We could in principle take our calculations further, calculating, for instance, the mean size of the small components in the region $\alpha > 2$ using Eq. (13.112), or the entire distribution of their sizes using Eq. (13.127).

13.11 DIRECTED RANDOM GRAPHS

In this chapter we have studied random graph models that go a step beyond the Poisson random graph of Chapter 12 by allowing us to choose the degree distribution of our model network. This introduces an additional level of realism to the model that makes it substantially more informative. It is, however, only a first step. There are many other features we can add to the model to make it more realistic still. We can for instance create random graph models of networks with assortative (or disassortative) mixing [237], bipartite structure [253], or clustering [247]. All of these models are still exactly solvable in the limit of large system size, although the solutions are more complicated than for the models we have seen in this chapter. For instance, in the case of the random graph with assortative mixing the fundamental generating function $g_1(z)$ becomes a vector, the corresponding equation (13.86) for the distribution of component sizes becomes a vector equation, and the condition for the existence of a giant component, Eq. (13.76), becomes a condition on the determinant of a matrix.




Figure 13.9: Size of the giant component for the configuration model with a power-law degree distribution. This plot shows the fraction of the network filled by the giant component as a function of the exponent α of the power law, calculated by numerical solution of Eqs. (13.91) and (13.141). The dotted lines mark the value $\alpha = 2$ below which the giant component has size 1 and the value $\alpha = 3.4788$ above which there is no giant component.

We will not go into detail on all of the many random graph models that have been proposed and studied, but in this section we take a look at one case, that of the directed random graph, as an example of the types of calculation that are possible.

13.11.1 GENERATING FUNCTIONS FOR DIRECTED GRAPHS

As discussed in Section 6.4, many networks, including the World Wide Web, metabolic networks, food webs, and others, are directed. The configuration model can be generalized to directed networks in a straightforward fashion, although the generalization displays some new behaviors not seen in the undirected case. Our presentation follows that of Refs. [100] and [253].

To create a directed equivalent of the configuration model, we must specify a double degree sequence, consisting of an in-degree j_i and an out-degree k_i for each vertex i . We can think of these as specifying the numbers of ingoing and outgoing stubs of edges at each vertex. Then we create a network by repeatedly choosing pairs of stubs—one ingoing and one outgoing—uniformly at random and connecting them to make directed edges, until no unused stubs remain. The result is a matching of the stubs drawn uniformly at random from the set of all possible matchings, just as in the configuration model, and the model itself is defined to be the ensemble of such directed networks in which each matching appears with equal probability. (The only small catch is that we must make sure that the total number of ingoing and outgoing stubs is the same, so that none are left over at the end of the process. We will assume this to be the case in the following developments.)

The probability that a particular outgoing stub at vertex w attaches to one of the j_v ingoing stubs at vertex v is

$$\frac{j_v}{\sum_i j_i} = \frac{j_v}{m},$$

(13.142)

where m is the total number of edges and we have made use of Eq. (6.26). Since the total number of outgoing stubs at w is k_w , the total expected number of directed edges from vertex w to vertex v is then $j_v k_w / m$, which is also the probability of an edge from w to v in the limit of large network size, provided the network is sparse. This is similar to the corresponding result, Eq. (13.32), for the undirected configuration model, but not identical—notice that there is no factor of two now in the denominator.

As in the undirected case we can, if we prefer, work with the degree distribution, rather than the degree sequence. As discussed in Section 8.3, the most correct way to describe the degree distribution of a directed network is by a joint distribution: we define p_{jk} to be the fraction of vertices in the network that have in-degree j and out-degree k . This allows for the possibility that the in- and out-degrees of vertices are correlated. For instance, it would allow us to represent a network in which the in- and out-degrees of each vertex were exactly equal to one another.²⁰¹ (This is rather an extreme example, but it demonstrates the point.)

The joint degree distribution can be captured in generating function form by defining a *double generating function* $g_{00}(x, y)$ thus:

$$g_{00}(x, y) = \sum_{j,k=0}^{\infty} p_{jk} x^j y^k.$$

(13.143)

(The two subscript zeros are the equivalent for the double generating function of the subscript zero in our previous generating function $g_0(z)$ for the undirected network.) As in the undirected case, the generating function $g_{00}(x, y)$ captures all the information contained in the degree distribution. Given the generating function we can reconstruct the degree distribution by differentiating:

$$p_{jk} = \frac{1}{j!k!} \left. \frac{\partial^j \partial^k g_{00}}{\partial x^j \partial y^k} \right|_{x,y=0}.$$

(13.144)

This is the equivalent for the directed case of Eq. (13.2) in the undirected case.

Just as in the undirected case the generating function satisfies certain conditions. First, since the degree distribution must be normalized according to $\sum_{jk} p_{jk} = 1$, the generating function satisfies

$$g_{00}(1, 1) = 1.$$

(13.145)

Second, the average in- and out-degrees are given by

$$\langle j \rangle = \sum_{jk=0}^{\infty} j p_{jk} = \left. \frac{\partial g_{00}}{\partial x} \right|_{x,y=1},$$

(13.146)

$$\langle k \rangle = \sum_{jk=0}^{\infty} k p_{jk} = \left. \frac{\partial g_{00}}{\partial y} \right|_{x,y=1}.$$

(13.147)

In a directed graph, however, the average in- and out-degrees are equal—see Eq. (6.27)—so $\langle j \rangle =$

$\square k \square$ and

$$\left. \frac{\partial g_{00}}{\partial x} \right|_{x,y=1} = \left. \frac{\partial g_{00}}{\partial y} \right|_{x,y=1}.$$

(13.148)

For convenience we will denote the average in-degree and out-degree by c in the equations that follow. Thus $\square j \square = \square k \square = c$.

We can also write down generating functions for the excess degree distribution of vertices reached by following an edge in the network. There are two different ways of following a directed edge—either forward or backward. Consider first the forward case. If we follow an edge forward to the vertex it points to, then the probability of reaching a particular vertex will be proportional to the number of edges pointing to that vertex, i.e., to its in-degree. Thus the joint degree distribution of such a vertex is proportional not to p_{jk} but to $j p_{jk}$. As before, we will be interested primarily in the number of edges entering and leaving a vertex other than the one we arrived along. If j and k denote these numbers then the total in-degree is $j + 1$ and the total out-degree is just k , so the distribution we want is proportional to $(j + 1) p_{j+1,k}$ or, correctly normalized, $(j + 1) p_{j+1,k} / c$. The double generating function for this excess degree distribution is then

$$\begin{aligned} g_{10}(x, y) &= \frac{\sum_{jk} (j + 1) p_{j+1,k} x^j y^k}{\sum_{jk} (j + 1) p_{j+1,k}} = \frac{\sum_{jk} j p_{jk} x^{j-1} y^k}{\sum_{jk} j p_{jk}} \\ &= \frac{1}{c} \frac{\partial g_{00}}{\partial x}. \end{aligned}$$

(13.149)

The backward case is similar. The appropriate excess degree distribution for the vertex from which an edge originates is $(k + 1) p_{j,k+1} / c$ and has generating function


$$g_{01}(x, y) = \frac{\sum_{jk} (k + 1) p_{j,k+1} x^j y^k}{\sum_{jk} (k + 1) p_{j,k+1}} = \frac{1}{c} \frac{\partial g_{00}}{\partial y}.$$

(13.150)

13.11.2 GIANT COMPONENTS

A directed graph has various different types of component, as discussed in Sections 6.11.1 and 8.1.1, including strongly and weakly connected components, in-components, and out-components. (Take a look at the “bow tie” diagram, Fig. 8.2 on page 240, for a reminder of the definitions of the components.) In general there can be both small and giant components of each of these types. Let us look at the giant components.

A strongly connected component is a set of vertices in which every vertex is reachable by a directed path from every other in the set. To put that a different way, for a vertex to belong to a strongly connected component at least one of its outgoing edges must lead to another vertex from which there is a path to the strongly connected component, and at least one of its ingoing edges must lead from a vertex to which there is path from the strongly connected component (see figure).



A vertex belongs to a strongly connected component if it has a directed path to the component and another from the component.

Let v be the probability that the vertex to which a randomly chosen edge in our graph leads has no directed path to the giant strongly connected component. For this to happen, it must be that none of the other outgoing edges from that vertex themselves have such a path. If the vertex has out-degree k , this happens with probability v^k . But j and k are distributed according to the excess degree distribution $(j+1)p_{j+1,k}/c$ and hence, averaging over both, we find that

$$v = \frac{1}{c} \sum_{j,k=0}^{\infty} (j+1)p_{j+1,k}v^k = g_{10}(1, v).$$

(13.151)

Similarly, consider the vertex from which a randomly chosen edge originates and let u be the probability that there is no path from the giant strongly connected component to that vertex. Then u is the solution to

$$u = g_{01}(u, 1).$$

(13.152)

Now consider a vertex with in-degree j and out-degree k . The probability that there is no path to the vertex from the giant strongly connected component via any of the vertex's j ingoing edges is u^j and the probability that there *is* such a path is $1 - u^j$. Similarly the probability that there is a path from the vertex to the giant strongly connected component is $1 - v^k$. And the probability that there are both—and hence that the vertex itself belongs to the giant strongly connected component—is the product of these two, or $(1 - u^j)(1 - v^k)$. Averaging this expression over the joint distribution of j and k we then find that the average probability S_s that a vertex lies in the giant strongly connected component, which is also the size of the giant strongly connected component measured as a fraction of the network size, is

$$\begin{aligned} S_s &= \sum_{j,k=0}^{\infty} p_{jk}(1 - u^j)(1 - v^k) \\ &= \sum_{j,k=0}^{\infty} p_{jk} - \sum_{j,k=0}^{\infty} p_{jk}u^j - \sum_{j,k=0}^{\infty} p_{jk}v^k + \sum_{j,k=0}^{\infty} p_{jk}u^jv^k \\ &= 1 - g_{00}(u, 1) - g_{00}(1, v) + g_{00}(u, v), \end{aligned}$$

(13.153)

with u and v given by Eqs. (13.151) and (13.152).

As discussed in Section 6.11, each strongly connected component in a network also has an in-component and an out-component associated with it—the sets of vertices from which it can be reached, and which can be reached from it. The in- and out-components of the giant strongly connected component are usually called the giant in- and out-components. By their definition, both are supersets of the giant strongly connected component itself, and we can calculate the size of both for our directed random graph. In fact, we have performed most of the calculation already.

A vertex with out-degree k fails to belong to the giant in-component only if none of its outgoing edges leads to a vertex that has a path to the strongly connected component. This happens with probability v^k , where v is as above. Averaging over j and k , we find the probability S_i that the vertex does belong to the giant in-component (which is also the size of the giant in-component) to be

$$S_i = 1 - \sum_{j,k=0}^{\infty} p_{jk}v^k = 1 - g_{00}(1, v).$$

(13.154)

Similarly the size of the giant out-component is given by

$$S_o = 1 - g_{00}(u, 1).$$

(13.155)

Using these results we can also write an expression for the combined size of the giant strongly connected component and its in- and out-components—the entire “bow tie” in Fig. 8.2. Since the giant in- and out-components both include the giant strongly connected component as a subset, their sum is equal to the size of the whole bow tie except that it counts the strongly connected part twice. Subtracting S_s to allow for this overcounting we then find the size of the bow tie to be and the fraction of the network not in the bow tie is just $g_{00}(u, v)$. (We could have derived this result by more direct means, just by noting that a vertex not in the bow tie has a path neither to nor from the giant strongly connected component.)

$$S_i + S_o - S_s = 1 - g_{00}(u, v),$$

(13.156)

And what about the giant weakly connected component? A weakly connected component in a directed graph is a normal graph component of connected vertices in which we ignore the directions of all the edges. At first glance one might imagine that the size of the giant weakly connected component was just equal to the combined size $S_i + S_o - S_s$ of the in-, out-, and strongly connected components calculated above. This, however, is not correct because the definition of the giant weakly connected component includes some vertices that are not in the in-, out-, or strongly connected components. An example would be any vertex that is reachable from the giant in-component but that does not itself have a path to the strongly connected component and hence is not in the giant in-component. Thus the size of the giant weakly connected component is, in general, larger than $S_i + S_o - S_s$. Nonetheless, we can still calculate the size of the giant weakly connected component by an argument quite similar to the ones we have already seen.

A vertex belongs to the giant weakly connected component if any of its edges, ingoing or outgoing, are connected to a vertex in that component. Let u now be the probability that a vertex is not connected to the giant weakly connected component via the vertex at the other end of one of its ingoing edges and let v be the equivalent probability for an outgoing edge. Then the probability that a vertex with in-degree j and out-degree k is not in the giant weakly connected component is $u^j v^k$ and the probability that it is in the giant weakly connected component is $1 - u^j v^k$. Averaging over the joint distribution p_{jk} of the two degrees we then find that the size S_w of the giant weakly connected component is

$$S_w = \sum_{jk} p_{jk} - \sum_{jk} p_{jk} u^j v^k = 1 - g_{00}(u, v).$$

(13.157)

We can derive the value of u by noting that the vertex at the end of an ingoing edge is not in the giant weakly connected component with probability $u^j v^k$ again, but with j and k being the numbers of edges excluding the edge we followed to reach the vertex. These numbers are distributed according to the appropriate excess degree distribution and, performing the average, we find that

$$u = g_{01}(u, v),$$

(13.158)

Similarly we can show that

$$v = g_{10}(u, v),$$

(13.159)

and Eqs. (13.157) to (13.159) between them give us our solution for the size of the giant weakly connected component.

13.11.3 THE APPEARANCE OF THE GIANT COMPONENTS

As in the undirected random graph, there may or may not be giant components in the direct random graph, depending on the degree distribution. We can derive conditions for the existence of the giant components using the machinery developed above. The calculation is easiest for the giant in- and out-components. Their size is given by Eqs. (13.154) and (13.155). Given that $g_{00}(1, 1) = 1$ (Eq. (13.145)), these equations give a non-zero size only if u or v is less than 1. Looking at Eq. (13.151) for the value of v we see a similar situation to that depicted in Fig. 13.6: we can have a solution with $v < 1$, and hence a giant in-component, only if

$$\left. \frac{\partial g_{10}}{\partial y} \right|_{x,y=1} > 1,$$

(13.160)

or equivalently if

$$\left. \frac{\partial^2 g_{00}}{\partial x \partial y} \right|_{x,y=1} > c,$$

(13.161)

where we have made use of Eqs. (13.148) and (13.149). Similarly we can have a giant out-component only if

$$\left. \frac{\partial g_{01}}{\partial x} \right|_{x,y=1} > 1,$$

(13.162)

or equivalently,

$$\left. \frac{\partial^2 g_{00}}{\partial x \partial y} \right|_{x,y=1} > c.$$

(13.163)

Interestingly, Eqs. (13.161) and (13.163) are identical, meaning that the conditions for the giant in–and out-components to appear are the same. If there is a phase transition at which one appears, the other also appears at the exact same moment.²⁰²

We can express (13.161) directly in terms of the degree distribution if we want. Substituting from Eq. (13.143) we find that

$$\left. \frac{\partial^2 g_{00}}{\partial x \partial y} \right|_{x,y=1} = \sum_{j,k=0}^{\infty} jk p_{jk}$$

(13.164)

and hence the giant in–and out-components appear if

$$\sum_{j,k=0}^{\infty} jk p_{jk} > c.$$

(13.165)

If we prefer we can write $c = \sum_j j p_{jk} = \sum_j k p_{jk}$ to give the alternative form

$$\sum_{j,k=0}^{\infty} (2jk - j - k) p_{jk} > 0.$$

(13.166)

This result is the equivalent for a directed network of Eq. (13.76) for the undirected case.

The calculation for the giant strongly connected component is similar. From Eq. (13.153) we see that $S_s = 0$ unless at least one of u and v is non-zero, so the condition for the existence of a giant strongly connected component is the same as for the in–and out-components, Eq. (13.166). In other words, the giant in-, out-, and strongly connected components all appear or disappear simultaneously.

The giant weakly connected component, however, is different. It is possible for there to be a giant weakly connected component in a network but no giant strongly connected component and hence the condition for the existence of a giant weakly connected component must be different from that for the other giant components. For instance, a network in which all vertices have either only ingoing edges or only outgoing edges can have a giant weakly connected component but trivially has no strongly connected component of size greater than one, since there are only paths to or from each vertex, but not both. Weakly connected components, however, are generally of less interest than strongly connected ones, and the calculation of the condition for the existence of the

giant weakly connected component is non-trivial, so we leave it as an exercise for the motivated reader and move on to other things.

13.11.4 SMALL COMPONENTS

We can also calculate the distribution of small components in a directed random graph. In fact the distribution of small strongly connected components is trivial: there aren't any. Or more properly the probability that a randomly chosen vertex belongs to a strongly connected component of size greater than one other than the giant strongly connected component is zero in the limit of large network size. To see this, recall that the small components in the undirected configuration model take the form of trees (see Section 13.7). If we consider a small strongly connected component in a directed network and ignore the directions of its edges, then the same argument we used before indicates that the resulting subgraph will also be a tree. But a tree has no loops in it, which leads to a contradiction because a strongly connected component must have loops—the paths in either direction between any pair of vertices form a loop. Thus, we conclude, there cannot be any small strongly connected components of size greater than one in the network.

In fact, this is not precisely true. In a random network there is always some chance that, for example, two vertices will each have a directed edge to the other, forming a strongly connected component of two vertices. In the limit of large n , however, the probability that a randomly chosen vertex belongs to such a component tends to zero. A detailed calculation shows that on average there is only a constant number of short loops in the network and their density vanishes as $1/n$ in the limit of large network size.

There can however be small in- and out-components. In a directed network, each strongly connected component has its own in- and out-components. In the present model, as we have said, we have no small strongly connected components, other than single vertices, so the component structure consists of the giant in- and out-components and then a large number of small in- and out-components for single vertices. Let us ask what the probability is that a randomly chosen vertex has a small out-component of size s , i.e., that there are s vertices including itself that can be reached by directed paths starting from the vertex. We can calculate the distribution of sizes by the same method we used for the undirected case. We define a generating function $h_1(y)$ for the distribution of the size of the out-component of a vertex reached by following an edge in the forward direction, which then satisfies an equation of the form of Eq. (13.86), except that the generating function g_1 for the excess degree distribution is replaced by the corresponding generating function for the directed network, Eq. (13.149), giving

$$h_1(y) = yg_{10}(1, h_1(y)).$$

(13.167)

And the generating function $h_0(y)$ for the size of the out-component to which a randomly chosen vertex belongs is then

$$h_0(y) = yg_{00}(1, h_1(y)).$$

(13.168)

We can write similar equations for in-components too and, armed with these equations, we can find the average size of the in–or out-component to which a vertex belongs, or even find the entire distribution of component sizes using the equivalent of Eq. (13.127).

PROBLEMS

13.1 Consider the binomial probability distribution $p_k = \binom{n}{k} p^k (1-p)^{n-k}$.

- a. Show that the distribution has probability generating function $g(z) = (pz + 1 - p)^n$.
- b. Find the first and second moments of the distribution from Eq. (13.25) and hence show that the variance of the distribution is $\sigma^2 = np(1-p)$.
- c. Show that the sum of two numbers drawn independently from the same binomial distribution is distributed according to $\binom{2n}{k} p^k (1-p)^{2n-k}$.

13.2 Consider a configuration model in which every vertex has the same degree k .

- a. What is the degree distribution p_k ? What are the generating functions g_0 and g_1 for the degree distribution and the excess degree distribution?
- b. Show that the giant component fills the whole network for all $k \geq 3$.
- c. What happens when $k = 1$?
- d. When $k = 2$ show that in the limit of large n the probability π_s that a vertex belongs to a component of size s is given by $\pi_s = 1 / [2\sqrt{n(n-s)}]$.

13.3 Consider the configuration model with exponential degree distribution $p_k = (1 - e^{-\lambda})e^{-\lambda k}$ with $\lambda > 0$, so that the generating functions $g_0(z)$ and $g_1(z)$ are given by Eq. (13.130).

- a. Show that the probability u of Eq. (13.91) satisfies the cubic equation

$$u^3 - 2e^\lambda u^2 + e^{2\lambda} u - (e^\lambda - 1)^2 = 0.$$

- b. Noting that $u = 1$ is always a trivial solution of this equation, show that the non-trivial solution corresponding to the existence of a giant component satisfies the quadratic equation $u^2 - (2e^\lambda - 1)u + (e^\lambda - 1)^2 = 0$, and hence that the size of the giant component, if there is one, is

$$S = \frac{3}{2} - \sqrt{e^\lambda - \frac{3}{4}}.$$

- c. Show that the giant component exists only if $\lambda < \ln 3$.

13.4 Equation (13.74) tells us on average how many vertices are a distance d away from a given vertex.

- a. Assuming that this expression works for all values of d (which is only a rough approximation to the truth), at what value of d is this average number of vertices equal to the number n in the whole network?

- b. Hence derive a rough expression for the diameter of the network in terms of c_1 and c_2 , and so argue that configuration model networks display the small-world effect in the sense that typical geodesic distances between vertices are $O(\log n)$.

13.5 Consider a network model in which edges are placed independently between each pair of vertices i, j with probability $p_{ij} = Kf_i f_j$, where K is a constant and f_i is a number assigned to vertex i . Show that the expected degree c_i of vertex i within the model is proportional to f_i , and hence that the only possible choice of probability with this form is $p_{ij} = c_i c_j / 2m$, as in the model of Section 13.2.2.

13.6 As described in Section 13.2, the configuration model can be thought of as the ensemble of all possible matchings of edge stubs, where vertex i has k_i stubs. Show that for a given degree sequence the number Ω of matchings is

$$\Omega = \frac{(2m)!}{2^m m!},$$

which is independent of the degree sequence.

13.7 Consider the example model discussed in Section 13.8.1, a configuration model with vertices of degree three and less only and generating functions given by Eqs. (13.94) and (13.95).

- a. In the regime in which there is no giant component, show that the average size of the component to which a randomly chosen vertex belongs is

$$\langle s \rangle = 1 + \frac{(p_1 + 2p_2 + 3p_3)^2}{p_1 - 3p_3}.$$

- b. In the same regime find the probability that such a vertex belongs to components of size 1, 2, and 3.

13.8 Consider a directed random graph of the kind discussed in Section 13.11.

- a. If the in- and out-degrees of vertices are uncorrelated, i.e., if the joint in/out-degree distribution p_{jk} is a product of separate functions of j and k , show that a giant strongly connected component exists in the graph if and only if $c(c - 1) > 0$, where c is the mean degree, either in or out.
- b. In real directed graphs the degrees are usually correlated (or anti-correlated). The correlation can be quantified by the covariance ρ of in- and out-degrees. Show that in the presence of correlations, the condition above for the existence of a giant strongly connected component generalizes to $c(c - 1) + \rho > 0$.
- c. In the World Wide Web the in- and out-degrees of the vertices have a measured covariance of about $\rho = 180$. The mean degree is around $c = 4.6$. On the basis of these numbers, do we expect the Web to have a giant strongly connected component?