


PC1–PC2 of the frozen embedding, QM7 molecules (seed 0; R² is out-of-fold ridge, not the picture)

unsupervised
(PC1+PC2 = 39% of variance)

MolWt
trained
2PC R²=0.02 · full R²=0.84




MolLogP
trained
2PC R²=0.03 · full R²=0.94



TPSA
trained
2PC R²=0.06 · full R²=0.94



Asphericity
NOT trained (2% w/ MolWt)
2PC R²=0.06 · full R²=0.64



RadiusOfGyration
NOT trained (8% w/ MolWt)
2PC R²=0.18 · full R²=0.81




label
NOT trained (9% w/ MolWt)
2PC R²=0.02 · full R²=0.15




supervised, desc
(PC1+PC2 = 35% of variance)


MolWt
trained
2PC R²=0.03 · full R²=0.92




MolLogP
trained
2PC R²=−0.00 · full R²=0.97




TPSA
trained
2PC R²=0.02 · full R²=0.96



Asphericity
NOT trained (2% w/ MolWt)
2PC R²=0.14 · full R²=0.64



RadiusOfGyration
NOT trained (8% w/ MolWt)
2PC R²=0.20 · full R²=0.84



label
NOT trained (9% w/ MolWt)
2PC R²=0.02 · full R²=0.18

