


PC1–PC2 of the frozen embedding, QM7 molecules (seed 0; R^2 is out-of-fold ridge, not the picture)

unsupervised
(PC1+PC2 = 39% of variance)

MolWt
trained
2PC $R^2=0.02$ · full $R^2=0.84$




MolLogP
trained
2PC $R^2=0.03$ · full $R^2=0.94$



TPSA
trained
2PC $R^2=0.06$ · full $R^2=0.94$




Asphericity
NOT trained (2% w/ MolWt)
2PC $R^2=0.06$ · full $R^2=0.64$



RadiusOfGyration
NOT trained (8% w/ MolWt)
2PC $R^2=0.18$ · full $R^2=0.81$




label
NOT trained (9% w/ MolWt)
2PC $R^2=0.02$ · full $R^2=0.15$



supervised, desc
(PC1+PC2 = 35% of variance)


MolWt
trained
2PC $R^2=0.03$ · full $R^2=0.92$




MolLogP
trained
2PC $R^2=-0.00$ · full $R^2=0.97$




TPSA
trained
2PC $R^2=0.02$ · full $R^2=0.96$



Asphericity
NOT trained (2% w/ MolWt)
2PC $R^2=0.14$ · full $R^2=0.64$



RadiusOfGyration
NOT trained (8% w/ MolWt)
2PC $R^2=0.20$ · full $R^2=0.84$



label
NOT trained (9% w/ MolWt)
2PC $R^2=0.02$ · full $R^2=0.18$

