

Jupiter Science in the Next Decade: From Juno to JUICE

juice

The 1st
European mission
to Jupiter

@LeighFletcher
(w/thanks to D. Grassi, M. Galand, T. Cavalie, R. Hueso, O. Witasse & JUICE Science Working Team)



UNIVERSITY OF
LEICESTER



European Research Council
Established by the European Commission
Supporting top researchers
from anywhere in the world




**THE ROYAL
SOCIETY**



The 1st
European mission
to Jupiter

Ariane 5 launch in 2022

A
7.5-year
journey



Europa
Callisto




Exploring **three**
potentially ocean-bearing
icy moons

and
3.5-year tour
of the Jovian system

1st orbiter of an icy moon:
Ganymede

5.3 tonne spacecraft



UNIVERSITY OF
LEICESTER


@LeighFletcher, ISSI Planetary Atmospheres Workshop, November 2018

How Did We Get Here?



- ESA **Cosmic Vision** 2007 – Laplace
- JGO+JEO=EJSM (Europa-Jupiter System Mission) 2008-2011.
- NASA Withdrawal led JGO->JUICE 2011-2012.
- Officially selected as L1 in May 2012.
- Phase A/B1 (prelim definition) to formal adoption Nov 2014.
- Airbus D&S selected July 2015.
- Passed PDR and into Phase C (detailed definition) March 2017, prototype units under construction.
- Upcoming CDR to move into Phase D (2.5yrs, Q2 2019).
- Launch May-June 2022.

JUICE Science Themes



- *Emergence of habitable worlds around gas giants*
- *Jupiter system as an archetype for gas giants*

Ganymede as a planetary object and possible habitat

- Largest satellite in the solar system
- Ocean between icy layers
- Internal dynamo
- Richest crater morphologies
- Archetype of waterworlds

Europa's recently active zones

- Impactor history
- Enigmatic differentiation
- Witness of early ages

Callisto as a remnant of the early Jovian system

- An active world?
- Ocean in contact with silicates

JUICE Science Themes



- *Emergence of habitable worlds around gas giants*
- *Jupiter system as an archetype for gas giants*

Jovian atmosphere

- Archetype for giant planets
- Fluid dynamics, chemistry, meteorology,...
- Formational history of planetary system

Jovian magnetosphere


- Largest object in our Solar System
- Astrophysical mechanisms at work
- Giant particle accelerator

Jovian satellite and ring systems


- Tidal forces: Laplace resonance
- Electromagnetic interactions to magnetosphere and upper atmosphere of Jupiter

Context: Why Jupiter?



- Previous exploration provided **snapshots**:
 - Flybys from New Horizons, Cassini, Voyagers, etc.
 - Limited data return from Galileo remote sensing
 - In situ sampling of unusual meteorological site.
- Literature reveals the need for:
 - **Continuity of data coverage** over long baselines to (i) define a mean atmospheric state and (ii) study departures from it.
 - **Broad, simultaneous spectral coverage** to relate visible changes to environmental conditions.
 - **Novel instrumental techniques** to reveal poorly-explored atmospheric regions and their coupling processes.






JUICE Jupiter Science Questions







JUICE Jupiter Science Questions





JUICE Jupiter Science Questions






JUICE Jupiter Science Questions




JUICE Jupiter Science Questions



JUICE Jupiter Science Questions



JUICE Jupiter Science Questions



- Require remote sensing across **broad wavelength range** to probe vertical structure from the thermosphere to the cloud decks and below.
- Require **high spatial resolutions and global coverage** of Jupiter's atmosphere at multiple atmospheric levels.
- Require **long baseline of observations** across a broad wavelength range to monitor variability on multiple timescales from hours to years.

JANUS: Visible Camera System

PI: Pasquale Palumbo, Parthenope University, Italy.
Co-PI: Ralf Jaumann, DLR, Germany


- $\geq 7.5\text{m/pixel}$
- Multiband imaging, 380 - 1080 nm
- Icy moon geology
- Io activity monitoring and other moons
- Jovian atmosphere dynamics



MAJIS: Imaging VIS-NIR/IR Spectrograph

PI: Yves Langevin, IAS, France
Co-PI: Giuseppe Piccioni, INAF, Italy

- 0.9-1.9 μm and 1.5-5.7 μm
- $\geq 62.5\text{ m/pixel}$
- Surface composition
- Jovian atmosphere



UVS: UV Imaging Spectrograph

PI: Randy Gladstone, SwRI, USA


- 55-210 nm
- 0.04°-0.16°
- Aurora and Airglow
- Surface albedos
- Stellar and Solar Occultation



PRIDE: Planetary Radio Interferometer & Doppler Experiment

PI: Leonid Gurvits, JIVE, EU/The Netherlands


- VLBI Radio Tracking
- S/C state vector
- Ephemerides
- bi-static and radio occultation experiments



SWI: Sub-mm Wave Instrument

PI: Paul Hartogh, MPS, Germany

- 600 GHz and 1200 GHz
- Jovian Stratosphere
- Moon atmosphere
- Atmospheric isotopes



JMAG: JUICE Magnetometer

PI: Michele Dougherty, Imperial, UK

- Dual Fluxgate and Scalar mag
- $\pm 8000\text{ nT}$ range, 0.2 nT accuracy
- Moon interior through induction
- Dynamical plasma processes



PEP: Particle Environment Package

PI: Stas Barabash, IRF-K, Sweden
Co-PI: Peter Wurz, UBe, Switzerland

- Six sensor suite
- Ions, electrons, neutral gas (in-situ)
- Remote ENA imaging of plasma and torus



GALA: Laser Altimeter

PI: Hauke Hussmann, DLR, Germany

- $\geq 40\text{ m}$ spot size
- $\geq 0.1\text{ m}$ accuracy
- Shape and rotational state
- Tidal deformation
- Slopes, roughness, albedo



RIME: Ice Penetrating Radar

PI: Lorenzo Bruzzone, Trento, Italy
Co-PI: Jeff Plaut, JPL, USA

- $\geq 40\text{ m}$ spot size
- $\geq 0.1\text{ m}$ accuracy
- Shape and rotational state
- Tidal deformation
- Slopes, roughness, albedo



RPWI: Radio and Plasma Wave Investigation

PI: Jan-Erik Wahlund, IRF-U, Sweden

- Langmuir Probes
- Search Coil Magnetometer
- Tri-axial dipole antenna
- E and B-fields
- Ion, electron and charged dust parameters



3GM: Gravity, Geophysics, Galilean Moons

PI: Luciano Iess, Rome, Italy

Co-PI: David J. Stevenson, CalTech, USA

Co-PI: Yohai Kaspi, Israel


- Ranging by radio tracking + USO
- 2 $\mu\text{m/s}$ range rate
- 20 cm range accuracy
- Gravity fields and tidal deformation



JUICE Payload

JANUS: Visible Camera System

Filter id.	λ / width [nm] (tbc)	Note
FPAN	650/500	Panchromatic – monochromatic imaging
FBLUE	450/60	Blue – satellite colours
FGREEN	530/60	Green, background for Na – satellite colours
FRED	656/60	Red, background for Ha – satellite colours
CMT medium	750/20	Continuum for strong Methane band on Jupiter, geology
Na	590/10	Sodium D-lines in exospheres
MT strong	889/20	Strong Methane band on Jupiter
CMT strong	940/20	Continuum for medium Methane band on Jupiter, Fe ²⁺ on satellites
MT medium	727/10	Medium Methane band on Jupiter
Violet	410/80	UV slope of satellites surfaces
NIR 1	910/80	Fe ²⁺ , Io lava spots
NIR 2	1000/150	Fe ²⁺ , Io lava spots
Ha	656/10	Ha-line for aurorae and lightnings



JANUS





Figure 9 Example of JANUS per Jove operations. Angular diameter of Jupiter (black-line, left axis) and phase angle (red line, right axis) surrounding P2 and 2G2.

Day-side: Global maps
wind noise measurement
level: $\sim 1 \text{ m/s}$

NEVER ACHIEVED BEFORE

Limb views at 12 km/pix
(low data required)
NEVER ACHIEVED BEFORE



Juno high-clouds
from views on the terminator
(60 km)

Feature tracking wind noise
measurement level: $\sim 0.3 \text{ m/s}$
Moderate DV required
NEVER ACHIEVED BEFORE

Resolution 22 km/pix
Size of Io footprint depend on
magnetic projection of Io
200-800 km (10-33 resolution
elements)


Lightning flashes size depends on its
depth. For lightning in water
condensation clouds 5 pixels width
lightning strokes are expected.

NEVER ACHIEVED BEFORE




MAJIS: Vis/NIR Spectrograph

- Pushbroom – 400-pixel long line, spectrum in each.
- **Two channels**
 - VIS-NIR: range 0.5-2.35 μm with a spectral sampling of 3.64 nm
 - IR: range 2.25-5.54 μm with a spectral sampling of 6.48 nm
- **Compared to previous (VIMS, NIMS, JIRAM):**
 - Longer wavelengths (better water/ NH_3 discrimination)
 - Shorter wavelengths (CH_4 bands for cloud tomography).
 - Higher resolution (break degeneracies, measure H_3^+).






From JIRAM to MAJIS




UVS: Imaging Spectrograph

- Updated design compared to Juno/UVS.
- EUV and FUV: 55 – 210 nm (e.g., emissions: H₂ bands, H-Lyman series; absorption: H₂, hydrocarbons, NH₃)
- **Primary modes:**
 - UV atmospheric emissions (airglow/aurora) & scattered sunlight
 - UV transmission in stellar & solar occultations.




Advantages of JUICE/UVS



- Juno focus on aurora, JUICE focus on middle and upper atmosphere globally.
- Juno narrow observing windows near perijove; JUICE extensive long-term monitoring throughout the orbital tour.
- Juno orbits constraints time of day (dawn/dusk), JUICE samples at a broad range of local times.
- JUICE capabilities for stellar and solar occultations.



SWI: Sub-mm Wave Instrument

- Never done before:
 - Direct measurement of middle atmospheric winds.
 - Temperature measurements from CH4.
 - Tracing composition from exogenic sources (H₂O, HCN, CO)
 - Water isotopologues and para-H₂ fraction for origins.
- Two passively cooled (140 K) Schottky receivers
 - 530 – 625 GHz (600-900 K DSB)
 - 1080 – 1275 GHz (3000-4000 K DSB)
 - IF-range: 3.5 – 8.5 GHz



SWI: Observing Strategy

- Scan single point over disc, 30 cm aperture elliptic antenna, movable ± 72 and ± 4.3 deg (antenna and rocker mechanisms)
- Continuous operations 16h/day, using moveable mirror to perform scans and rasters.
- CH_4/HCN for T retrieval; H_2O for composition; Doppler shift for direct winds.





3GM: Jupiter Radio Occultations

- **Three components:**

- High performance dual-frequency radio links (X and Ka-band)
- Steerable medium gain antenna (MGA),
- Ultra-stable oscillator (USO).


- **Radio occultations of neutral atmosphere and ionosphere:**

- Retrieve vertical profiles of density, pressure, temperature, and NH₃.
- Measure ionospheric electron densities in auroral to study auroral structure and energy transport.
- Study the relation between vertically propagating waves and the heating mechanisms for the thermosphere.
- Inference of zonal winds.



JUICE Tour


Date	Event or phase
June 2022	Launch from Kourou with Ariane 5
May 2023	Earth flyby #1
October 2023	Venus flyby
September 2024	Earth flyby #2
February 2025	Mars flyby
November 2026	Earth flyby #3
October 2029	Jupiter orbit insertion
October 2029-October 2030	Energy reduction phase
October 2030	2 Europa flybys
October 2030-August 2031	Jupiter inclined phase - Callisto flybys
September 2031-November 2032	Phase "transfer to Ganymede"
December 2032	Ganymede orbit insertion
December 2032-September 2033	Elliptical and circular orbits (5000/500 km)
September 2033	End of mission



1. Pre-JOI and first ellipse, from 7 April 2029 to 27 May 2030 (PJ1)
2. Energy reduction phase, from 27 May 2030 to 28 September 2030 (PJ2-5)
3. Europa flybys, from 28 September 2030 to 26 October 2030 (PJ6-7)
4. High latitude, from 26 October 2030 to 25 June 2031 (PJ8-20)
5. Low energy phase, from 24 June 2031 to 24 August 2032 (PJ21-51)


Current Status (Phase C)

- Four working groups assessing tour opportunities.
- Segmentation of tour based on science priorities (Level-0 planning).
- Level-1 planning in the coming years.




JUICE in Wider Context

- **Juno** extension to ~2021.
- **JWST** launch in ~2021 (0.6-28.5 micron):
 - Near Infrared Camera (NIRCam)
 - Near Infrared Spectrograph (NIRSpec)
 - Mid-Infrared Instrument (MIRI)
 - GTO programme for Jupiter, Saturn, Uranus, Neptune (Hammel, Fletcher)
 - ERS programme for Jupiter (de Pater, Fouchet)
- **ELT** first light ~2024.
 - HARMONI IFS 0.5-2.5 μm
 - METIS Mid-IR spectrograph 3-20 μm
 - MICADO Near-IR imager/spectrograph 0.8-2.4 μm
- **Europa Clipper** ~2023
 - Launch on SLS?
 - Focused on Europa, but could be persuaded to look at some “calibration” objects....



Summary & Perspectives

- **JUICE 2.5-year tour** achieves:
 - Multi-wave remote sensing to probe atmosphere in 3D from clouds to thermosphere.
 - Long baseline of observations.
 - Novel instrument techniques.
 - Simultaneity of data to understand coupling processes.
- Tour and payload specifics continue to evolve.
- Complements continued **Earth- and space-based** programmes.
- On course for **2022 launch** on one of the last Ariane 5s.
 - ~180 weeks away....



First European mission to Jupiter.

Three ocean-bearing icy worlds.

First orbiter of an icy moon.

First mission to the largest planetary satellite in the Solar System.