UNIVERSITÀ DEGLI STUDI DI MILANO

FACOLTÀ DI SCIENZE E TECNOLOGIE

FACOLTÀ DI SCIENZE POLITICHE,
ECONOMICHE E SOCIALI

*Master Degree in Data Science for Economics*

# THESIS TITLE

**Supervisor:** Prof. Luca ROSSINI

**Co-supervisors:** Prof. Maria Fernanda PINTADO SERRANO

Thesis by:
Leila CIELOK
Student ID: 65522A

Academic Year 2026/2027

*<exergo - dedicate>*

# Aknowledgments

ACKNOWLEDGMENTS

# Contents

**Aknowledgments** **iii**

**1 Introduction** **2**

1.1 Research Question . . . . . 2

1.2 Contributions . . . . . 2

1.3 Thesis structure . . . . . 3

**2 Literature review** **4**

2.1 Section 1 . . . . . 4

2.2 Section 2 . . . . . 4

2.2.1 Subsection 2.1 . . . . . 4

**3 Data and network construction** **5**

3.1 Data sources . . . . . 5

3.1.1 LNG trade and infrastructure data . . . . . 6

3.1.2 Maritime routes and chokepoint data . . . . . 8

3.2 Data preparation . . . . . 12

3.2.1 Terminal infrastructure characteristics . . . . . 13

3.2.2 Temporal aggregation . . . . . 14

3.3 Dynamic network construction . . . . . 14

3.3.1 Nodes and edges . . . . . 15

3.3.2 Node activity . . . . . 16

3.4 Construction of node-level measures . . . . . 17

3.4.1 Terminal criticality measures . . . . . 17

3.4.2 Chokepoint criticality measures . . . 21
3.4.3 Common structural indicators . . . 23
3.5 Construction of the model-ready panels . . . 25
3.5.1 Node activity panel . . . 26
3.5.2 Terminal criticality panel . . . 27
3.5.3 Chokepoint criticality panel . . . 27
3.6 Data and methodological limitations . . . 28

**4 Exploratory network analysis 30**
4.1 Network evolution . . . 31
4.2 Node activity and temporal sparsity . . . 32
4.3 Terminal network position . . . 34
4.3.1 Terminal roles and trade activity . . . 34
4.3.2 Centrality, concentration and dependence . . . 35
4.4 Chokepoint network position . . . 40
4.4.1 Flow exposure . . . 40
4.4.2 Centrality measures . . . 41
4.5 Structural position of nodes . . . 43
4.6 Associations and modelling implications . . . 46

**5 Bayesian Reduced-Rank model for network measures 49**
5.1 General modelling framework . . . 49
5.1.1 Estimation samples, responses and covariates . . . 49
5.1.2 Response likelihoods . . . 51
5.1.3 Prior specification . . . 52
5.2 Model specifications . . . 54
5.2.1 Baseline rank-one specification . . . 54
5.2.2 Random-intercept rank-one specification . . . 60
5.2.3 Rank-two specification . . . 64
5.3 Posterior computation . . . 66

5.3.1 NUTS sampling algorithm and implementation . . . . . 66
5.3.2 Sampling diagnostics . . . . . 69

**6 Results and discussion** **71**
6.1 Rank-one model results . . . . . 71
6.1.1 Terminal posterior analysis . . . . . 71
6.1.2 Chokepoint posterior analysis . . . . . 78
6.1.3 Sampling diagnostics . . . . . 83
6.2 Node-specific intercept model results . . . . . 86
6.2.1 Terminal posterior analysis . . . . . 86
6.2.2 Chokepoint posterior analysis . . . . . 91
6.2.3 Sampling diagnostics . . . . . 96
6.3 Rank-two model results . . . . . 97
6.4 Prior sensitivity results . . . . . 140
6.4.1 Alternative prior specifications . . . . . 140
6.4.2 Prior predictive assessment . . . . . 141
6.4.3 Posterior sensitivity and predictive robustness . . . . . 142
6.5 Comparative discussion and limitations . . . . . 144
6.5.1 Model limitations and implications . . . . . 144

**7 Conclusions and future work** **147**

**A Chokepoint geometry registry** **148**

**B Sensitivity analysis of chokepoint assignments** **151**
B.1 Purpose and method . . . . . 151
B.2 Sensitivity results . . . . . 151

**C Model-ready panels variables** **154**
C.1 Node activity panel . . . . . 154
C.2 Terminal criticality panel . . . . . 156

C.3 Chokepoint criticality panel . . . . . 159

**D Full conditional posterior distributions** **162**

D.1 Baseline rank-one specification . . . . . 162

D.1.1 Factor-regression coefficients . . . . . 162

D.1.2 Response loadings . . . . . 163

D.1.3 Response intercepts . . . . . 168

D.1.4 Beta precision parameters . . . . . 171

D.1.5 Latent factors . . . . . 172

D.2 Random-intercept rank-one specification . . . . . 174

D.2.1 Factor-regression coefficients . . . . . 174

D.2.2 Latent factors . . . . . 175

D.2.3 Node-specific intercepts . . . . . 176

D.2.4 Node-effect scale parameter . . . . . 177

**E Additional posterior results** **179**

E.1 Baseline rank-one specification . . . . . 179

E.1.1 Terminals . . . . . 179

E.1.2 Chokepoints . . . . . 184

E.2 Node-intercept rank-one specification . . . . . 184

E.2.1 Terminals . . . . . 184

E.2.2 Chokepoints . . . . . 188

**F Model diagnostics** **190**

F.1 Prior sensitivity analysis . . . . . 190

F.1.1 Prior predictive checks . . . . . 190

F.1.2 MCMC sampling diagnostics . . . . . 190

F.2 Rank-one MCMC convergence diagnostics . . . . . 191

F.2.1 R-hat and Effective Sample Sizes . . . . . 191

F.2.2 Geweke diagnostic . . . . . 191

F.2.3 Heidelberger–Welch diagnostic . . . . . 192
F.2.4 Trace plots . . . . . 193
F.2.5 Autocorrelation diagnostics . . . . . 205
F.3 Random-intercept rank-one MCMC convergence diagnostics . . . . . 205

# Abstract

ABSTRACT.

# Chapter 1

# Introduction

Background and Motivation

## 1.1 Research Question

Research problem: identifying how global lng maritime network nodes' criticality evolves in time, distinguishing the probability that a node is active from its criticality conditioned on being active.

1. How will the structure of the global LNG maritime network evolve between 2020 and 2024?

2. What characteristics determine the probability that a terminal or chokepoint will be operational in a given month?

3. Which latent dimensions collectively describe the various measures of criticality?

4. How do structural, infrastructural and dynamic characteristics influence the criticality of nodes?

5. How do the mechanisms of criticality differ between terminals and chokepoints?

## 1.2 Contributions

Empirical (new monthly lng maritime dataset), methodology (separation: activity vs conditional criticality), theoretical (multidimensionality of criticality).

## 1.3 Thesis structure

This thesis is structured as follows...

# Chapter 2

# Literature review

## 2.1 Section 1

Global LNG Maritime trade networks: LNG organization, description of nodes' roles and routes.

## 2.2 Section 2

Potential subsections:

Criticality and Vulnerability in network analysis: comparison of importance, centrality, criticality, vulnerability, network science measures' roles.

Dynamic and temporal networks: temporal dependence, nodes inactivity and persistence.

Statistical models for sparse network processes: hurdle models

Bayesian reduced-rank models: multivariate bayesian models, shrinkage...

Research gap: combine dynamicity, multidimensional criticality, terminals and chokepoints analysis together

### 2.2.1 Subsection 2.1

# Chapter 3

# Data and network construction

This chapter describes the data sources and processing steps used to construct the dynamic LNG maritime network analysed in this thesis. The final dataset combines observed LNG tanker voyages, import and export terminals, reconstructed maritime routes, and the strategic chokepoints traversed by those routes. Together, these components represent the global LNG transportation system as a directed and weighted network evolving at a monthly frequency from January 2020 to December 2024.

The construction process involved harmonising the source data, matching observed voyages to their departure and arrival terminals, reconstructing the corresponding maritime routes, identifying the chokepoints crossed by each route, and aggregating trade flows and network relationships at the monthly level. The resulting dataset contains the nodes, connections, flow volumes, route characteristics, and infrastructure attributes required for the descriptive and Bayesian analyses developed in the following chapters.

## 3.1 Data sources

Despite the growing importance of LNG for international energy trade and energy security, publicly available data with high spatial and temporal resolution remain limited. Official and industry sources generally report LNG trade at the country level and at monthly or annual frequencies, whereas more granular information is often available only through commercial databases. Moreover, to the best of my knowledge, no single

open-access dataset jointly represents individual LNG voyages, terminal infrastructure, reconstructed maritime routes, and the strategic chokepoints traversed by those routes.

The dataset used in this thesis was therefore constructed by integrating voyage-level LNG trade records and terminal infrastructure data with reconstructed maritime routes and additional geographic information on maritime chokepoints. The following subsections describe the four main components of this integrated dataset.

### 3.1.1 LNG trade and infrastructure data

The main data source is the Liquefied Natural Gas from Tanker, Terminal, and Trade (LNG-T3) dataset developed by Zhou et al. (2026). LNG-T3 is an open-access dataset providing global LNG infrastructure and trade information at daily resolution for the period from January 2020 to December 2024. It was constructed by combining Automatic Identification System (AIS) vessel-position and draught reports, historical port-call records, LNG tanker specifications, and terminal metadata. The complete dataset contains 17,592 individual LNG tanker voyage events and provides information on tanker movements, terminal activity, and country-to-country trade flows.

Two components of LNG-T3 were used directly in this study: the voyage-level records and the terminal inventory. The voyage data form the empirical basis for identifying directed LNG trade connections and measuring the volumes transported between terminals. The terminal inventory defines the infrastructure nodes of the network and provides the geographic and structural attributes used to characterise them. These two components were linked through the terminal identifiers included in the original dataset.

LNG-T3 was preferred to aggregate country-level trade statistics because its voyage-level resolution makes it possible to identify the specific origin and destination terminals associated with each LNG shipment. Its 2020–2024 coverage also provides a consistent five-year observation window that captures both established trade patterns and major changes in the global LNG market, including those following the disruption

of European gas supplies in 2022.

### LNG voyages

The LNG-T3 voyage data contain individual movements of LNG tankers between departure and arrival terminals. For each voyage, the available information includes the tanker identifier, the origin and destination terminals, the departure and arrival dates, and the estimated volume of LNG transported. These records were used to identify the direction, timing, and magnitude of LNG flows within the network.

Each voyage was represented as a directed movement from its departure terminal to its arrival terminal. The voyage date was then used to assign the observation to a monthly period between January 2020 and December 2024. When multiple voyages occurred between the same origin–destination pair during the same month, their transported volumes were aggregated. The resulting monthly observations define the directed and weighted trade connections between terminal nodes.

Voyage identifiers were also used to retain information on the number of shipments associated with each connection. This makes it possible to distinguish between connections characterised by a small number of high-volume voyages and those supported by more frequent tanker movements. Once the corresponding maritime routes had been reconstructed, route distance and chokepoint crossings were linked back to the individual voyage records and subsequently aggregated at the monthly level.

### LNG terminals

The LNG-T3 terminal inventory provides the identifiers, names, geographic coordinates, countries, and infrastructure characteristics of LNG facilities worldwide. Terminals constitute the origin and destination nodes of the trade network and were linked to the voyage records using their unique identifiers.

The infrastructure attributes retained for the analysis include the terminal's country, infrastructure type, processing capacity, number of infrastructure units, and start year. Infrastructure type distinguishes facilities according to their import or export

function, while capacity provides an indicator of their potential scale of operation. The number of infrastructure units supplies an additional measure of facility size and complexity. The start year was used to construct terminal age as a time-varying variable, calculated for each monthly observation as the difference between the observation year and the year in which the terminal began operating.

Terminal coordinates were used both to position terminal nodes geographically and to reconstruct the maritime route associated with each observed origin–destination pair. Before route reconstruction, terminal identifiers and coordinates were checked for consistency across the voyage and infrastructure data. This harmonisation ensured that every retained voyage could be linked to a valid departure terminal and arrival terminal.

### 3.1.2 Maritime routes and chokepoint data

While the LNG-T3 voyage records identify the origin and destination of each shipment, they do not provide the actual maritime path connecting the two terminals. For this reason, maritime routes were reconstructed from the geographic coordinates of the departure and arrival terminals using the `SeaRoute` routing software (Eurostat, 2021).

#### SeaRoute software

SeaRoute applies Dijkstra's algorithm (Dijkstra, 1959) to identify the shortest path between two locations on a predefined network representing frequently used shipping lanes, which accounts for coastlines, navigable waterways, and supported maritime passages. Specifically, the 5-km resolution of the maritime network produced through the `marnet` module was used: edges shorter than this scale are progressively contracted and geometrically similar connections are removed, reducing network complexity while preserving the principal maritime routes. Additionally, in line with the thesis' objectives, transit through both the Suez and Panama canals was enabled, allowing the routing algorithm to include these passages when they formed part of the shortest available maritime path.

A route was generated for each unique origin–destination terminal pair and linked to all voyages associated with that pair. Afterwards, the reconstructed routes were merged with the chokepoints locations to identify the strategic maritime passages traversed by each trade connection.

### Chokepoints geometries

The initial set of 28 maritime chokepoints, together with their names, identifiers, and reference coordinates, was obtained from the IMF PortWatch chokepoint database (International Monetary Fund, 2026), but because the PortWatch locations are represented primarily as point features, they could not be used directly to determine whether a reconstructed maritime route effectively crossed a passage. A geometry registry was therefore compiled to associate each chokepoint with an appropriate spatial representation. The registry, including the source and construction method for each geometry, is provided in Appendix A.

The final geometries were assembled from several sources. Published polygons for the Suez Canal, Bab el-Mandeb Strait, and Cape of Good Hope were directly obtained from the World Bank Development Data Partnership's Red Sea Monitoring project (World Bank Data Lab, 2025). Detailed IHO or SeaVoX geometries for the Malacca, Hormuz, Gibraltar, Dover, and Makassar straits were obtained through the Marine Regions Gazetteer and its geospatial services (Flanders Marine Institute, nd). For the Lombok, Sunda, Torres, and Bering straits, transit geometries were reconstructed from the WGS84 coordinates reported in official International Maritime Organization routeing documents (International Maritime Organization, 2019, 2014; Russian Federation and United States, 2017).

Where no suitable published polygon was available, operational geometries were constructed specifically using either a buffered centreline, as in the Panama Canal and Bosporus Strait, or an oriented crossing gate positioned across the relevant navigable passage. The placement of these geometries was informed by the reference coordinates provided by PortWatch (International Monetary Fund, 2026), geographic information

from Marine Regions (Flanders Marine Institute, nd), and navigational references from relevant IMO documents, the NGA GeoNames database, and NOAA/NCEI hydrographic survey records (International Maritime Organization, 1998; National Geospatial-Intelligence Agency, nd; National Centers for Environmental Information, nd).

Figure 1 shows the locations of the 28 candidate chokepoints and representative examples of the four types of geometry used to identify route crossings: a published polygon, a polygon reconstructed from IMO routeing coordinates, an operational crossing gate, and a buffered centreline.

All geometries were transformed into a common geographic coordinate system and stored in a harmonised geospatial format. To accommodate small discrepancies between the reconstructed `SeaRoute` paths and the chokepoint geometries, a uniform spatial-tolerance buffer of 25 km was applied around each geometry. A reconstructed origin–destination route was classified as traversing a chokepoint when it intersected the corresponding buffered geometry.

Of the 28 candidate chokepoints, 24 were traversed by at least one reconstructed route under the final 25-km specification and were therefore retained as nodes in the monthly network. The robustness of the route–chokepoint assignments was assessed by comparing this specification with exact geometric intersection and with alternative buffers of 5 and 10 km. The complete sensitivity analysis is reported in Appendix B.

Table 1 summarises the principal data sources, their units of observation, the main variables retained, and their role in the construction of the network.




Figure 1: PortWatch reference locations of the 28 candidate chokepoints and representative route-crossing geometries. The enlarged panels show the Suez Canal (published polygon), Lombok Strait (geometry reconstructed from IMO routeing coordinates), Taiwan Strait (operational crossing gate), and Panama Canal (buffered operational centreline).

Table 1: Summary of data sources

| **Source** | **Unit of observation** | **Main variables** | **Role in the analysis** |
|---|---|---|---|
| LNG-T3 tanker voyages | Individual voyage | Departure and arrival terminals, voyage dates, tanker identifier, estimated LNG volume | Identification of directed LNG trade connections and monthly flow volumes from January 2020 to December 2024 |
| LNG-T3 terminal inventory | LNG terminal | Terminal identifier, name, coordinates, country, infrastructure type, capacity, unit count, start year | Definition and characterisation of terminal nodes |
| `searoute` | Unique origin–destination pair | Reconstructed maritime path, route distance | Reconstruction of the geographic route associated with each trade connection |
| Chokepoint geographic data | Maritime chokepoint | Name, type, and geographic location | Definition of chokepoint nodes, identification of route crossings |

## 3.2 Data preparation

Before constructing the network, the voyage and terminal datasets were harmonised to ensure that each observed shipment could be associated with a unique origin and destination node.

As an initial validation step, voyage dates, LNG volumes, terminal capacities, and unit counts were parsed into consistent formats. No invalid or missing values were found in these fields.

The LNG-T3 terminal inventory contains 545 records corresponding to 471 unique terminal names. Sixty-five names occur more than once, generally because the same facility has multiple infrastructure-status records. Whereas most repeated records have identical coordinates, three terminal names have two coordinate variants each.

Records were first distinguished by name and coordinates. One record was then

selected to represent each terminal name in the final terminal map, based on the infrastructure status reported for that record in the LNG-T3 inventory. The selection prioritised records classified as *operating*, followed by those classified as *under construction*, *idle*, *mothballed*, and *proposed*. Each of the 471 terminals in the final map was then assigned a unique node identifier.

Voyage origins and destinations were matched to the consolidated terminal map using the terminal names reported in the LNG-T3 voyage data. The corresponding node identifiers and coordinates were attached to each voyage record, and the matching status of the two endpoints was recorded separately. All origin and destination values in the 17,592 voyage records were successfully matched to the terminal map.

The original voyage table contains 8,645 laden export voyages and 8,947 return movements. Since the latter take place after delivery and therefore report zero transported LNG volume, only the laden export voyages were retained to construct the LNG trade network. The retained voyages were also required to have a valid departure date, matched origin and destination identifiers, and a strictly positive LNG volume. All export records satisfied these conditions, meaning that these checks did not produce any further exclusions.

Potential duplicate export observations were identified using the departure and arrival dates, tanker IMO number, voyage type, origin and destination terminals, and transported volume. After removing three duplicate entries, the final dataset used to construct the network contains 8,642 laden LNG voyages.

### 3.2.1 Terminal infrastructure characteristics

Terminal characteristics were obtained from the LNG-T3 infrastructure inventory and linked to the consolidated terminal identifiers. The retained attributes include country, region, geographic coordinates, infrastructure type, operational status, annual processing capacity, number of infrastructure units, and start year. No missing values were found for these attributes among the terminals retained in the final network.

Additionally, terminal age was constructed as a time-varying characteristic by subtracting the terminal's reported start year from the year of each monthly observation:

$$\mathrm{Age}_{i,t} = \mathrm{Year}_t - \mathrm{StartYear}_i \,. \tag{1}$$

### 3.2.2 Temporal aggregation

Each retained voyage was assigned to a monthly period according to its departure date. The resulting observation window extends from January 2020 to December 2024, corresponding to the complete five-year coverage of the LNG-T3 dataset and yielding 60 monthly network observations.

To preserve months in which a node had no observed activity, complete node–month panels were subsequently constructed by combining every retained node with all 60 monthly periods. Flow and voyage measures for node–month observations with no activity were assigned a value of zero to distinguish these structural zeros from missing or non-applicable node characteristics, which remained coded as unavailable.

## 3.3 Dynamic network construction

The global LNG maritime network was represented as a sequence of monthly directed and weighted graphs,

$$G_t = (V, E_t, W_t), \qquad t = 1, \ldots, 60, \tag{2}$$

where $V$ is the time-invariant set of terminal and chokepoint nodes, $E_t$ is the set of directed connections observed in month $t$, and $W_t = [w_{ij,t}]$ is the corresponding weighted adjacency matrix. An element $w_{ij,t} > 0$ represents the total LNG volume transported from node $i$ to node $j$ during month $t$, while $w_{ij,t} = 0$ indicates that no such connection was observed.

### 3.3.1 Nodes and edges

The final node set contains 183 nodes and can be written as

$$V = V^T \cup V^C, \qquad |V^T| = 159, \qquad |V^C| = 24, \tag{3}$$

where $V^T$ denotes the set of LNG terminals and $V^C$ the set of maritime chokepoints.

The 159 terminal nodes are the facilities that appear as an origin or destination of at least one retained laden voyage. Of the 28 candidate chokepoints included in the geographic registry, 24 were traversed by at least one reconstructed route and were therefore retained in the observed network.

Each commercial voyage defines an underlying directed movement from an export terminal to an import terminal. To incorporate the maritime infrastructure traversed during transportation, the reconstructed route of voyage $r$ was represented as an ordered path

$$P_r = (v_{r,0}, v_{r,1}, \ldots, v_{r,K_r}), \tag{4}$$

where $v_{r,0}$ is the departure terminal, $v_{r,K_r}$ is the arrival terminal, and any intermediate nodes are the chokepoints crossed by the route. The voyage was then expanded into the consecutive directed connections

$$E(P_r) = \{(v_{r,0}, v_{r,1}), (v_{r,1}, v_{r,2}), \ldots, (v_{r,K_r-1}, v_{r,K_r})\}. \tag{5}$$

A route that crossed no retained chokepoint generated a direct terminal-to-terminal edge. Routes crossing one or more chokepoints generated terminal-to-chokepoint, chokepoint-to-chokepoint, and chokepoint-to-terminal edges according to the order in which those nodes were encountered.

Let $\mathcal{R}_t$ denote the set of voyages during month $t$, and $q_r$ the LNG volume transported by voyage $r$. The edge weight from node $i$ to node $j$ at month $t$ is defined as

$$w_{ij,t} = \sum_{r \in \mathcal{R}_t} q_r \, \mathbb{I}\left\{(i,j) \in \mathcal{E}(P_r)\right\}, \tag{6}$$

where $\mathbb{I}\{\cdot\}$ is the indicator function. Thus, the full volume of a shipment was assigned to every segment traversed by its route. Because shipments crossing multiple segments would consequently be counted more than once in the sum of all edge weights, aggregate monthly trade volume was calculated directly from the original voyage observations.

Voyages sharing the same directed pair of nodes within a month were aggregated into a single edge. In addition to LNG volume, each monthly edge records the number of voyage traversals, the number of distinct origin–destination routes using the connection, and their route identifiers.

### 3.3.2 Node activity

Node activity was defined from the presence of incident flow in each monthly graph. For node $i$ in month $t$, the binary activity indicator was defined as

$$a_{i,t} = \mathbb{I}\left\{\sum_{j \in V} w_{ij,t} + \sum_{j \in V} w_{ji,t} > 0\right\}. \tag{7}$$

A terminal was therefore classified as active when it recorded positive incoming or outgoing LNG flow during the month. Since terminal direction was fully consistent with the infrastructure classification reported in LNG-T3, active export terminals recorded outgoing flow and active import terminals recorded incoming flow.

A chokepoint was classified as active when at least one voyage traversed it during the month, generating positive incident flow on one or more adjacent route segments. Chokepoints not traversed by any voyage in a given month were retained in the balanced node–month panel and assigned an activity value of zero.

## 3.4 Construction of node-level measures

A set of monthly node-level measures was constructed to represent different dimensions of criticality in the LNG transportation network. Since terminals and chokepoints perform different functions, their criticality was evaluated using separate but complementary measures and network representations.

On the one hand, terminals represent the origins and destinations of commercial LNG flows. Their criticality was therefore defined in terms of their importance as suppliers or recipients, the concentration of their trading relationships, and the dependence associated with those relationships. On the other hand, chokepoints represent intermediate passages along maritime routes, and their criticality was consequently defined in terms of their position within the physical transportation network, the volume of LNG passing through them, and their role in connecting otherwise separated parts of the network or in mediating important transportation paths.

Terminal criticality measures were calculated from the original terminal-to-terminal voyage flows, thereby preserving the direct commercial relationship between origins and destinations and avoiding the repeated counting of shipments across multiple route segments. Chokepoint criticality measures were instead calculated on the complete segment-level network, including both terminals and chokepoints, so that the position of each passage within the full sequence of maritime movements could be represented. Additional structural indicators were calculated for both node types.

### 3.4.1 Terminal criticality measures

Terminal criticality was represented through measures of directional network importance, counterparty concentration, trade dependence, and voyage distance. These measures were constructed monthly and, where relevant, adapted to the terminal's observed role as an LNG exporter or importer.

### Terminal role

Let $q_{ij,t}$ denote the total LNG volume transported from terminal $i$ to terminal $j$ during month $t$. Monthly outgoing flow, incoming flow, and throughput were defined as

$$\begin{aligned} O_{i,t} &= \sum_{j} q_{ij,t}, \\ I_{i,t} &= \sum_{j} q_{ji,t}, \\ T_{i,t} &= O_{i,t} + I_{i,t}. \end{aligned} \tag{8}$$

A terminal was classified as active in month $t$ when $T_{i,t} > 0$. Among active terminals, the observed monthly role was classified as exporter when $O_{i,t} > 0$ and $I_{i,t} = 0$, importer when $I_{i,t} > 0$ and $O_{i,t} = 0$, and bidirectional when both flows were positive.

For active terminal–months, export share was defined as

$$s_{i,t}^{\text{out}} = \frac{O_{i,t}}{T_{i,t}}, \tag{9}$$

so that exporters have $s_{i,t}^{\text{out}} = 1$, importers have $s_{i,t}^{\text{out}} = 0$, and bidirectional terminals take values between zero and one.

### Counterparty concentration

The concentration of a terminal's trade across its counterparties was measured using the Herfindahl–Hirschman Index (HHI), a standard measure of concentration based on the squared shares of individual counterparties (Hirschman, 1945).

Let $\mathcal{D}_{i,t} = \{j : q_{ij,t} > 0\}$ be the set of destination terminals receiving LNG from terminal $i$ in month $t$, and let $\mathcal{S}_{i,t} = \{j : q_{ji,t} > 0\}$ be the set of origin terminals supplying it. The outgoing and incoming terminal-level concentration indices were calculated, respectively, as

$$\begin{aligned} H_{i,t}^{\text{out}} &= \sum_{j \in \mathcal{D}_{i,t}} \left( \frac{q_{ij,t}}{O_{i,t}} \right)^2, \\ H_{i,t}^{\text{in}} &= \sum_{j \in \mathcal{S}_{i,t}} \left( \frac{q_{ji,t}}{I_{i,t}} \right)^2. \end{aligned} \tag{10}$$

The same procedure was applied after aggregating counterparties by country, producing corresponding country-level concentration indices. Values approaching one indicate that flows are concentrated in a small number of counterparties, whereas lower values indicate greater diversification.

A role-specific concentration measure was then assigned according to the terminal's observed direction of trade:

$$H_{i,t}^{\text{role}} = \begin{cases} H_{i,t}^{\text{out}}, & \text{if } O_{i,t} > 0 \text{ and } I_{i,t} = 0, \\ H_{i,t}^{\text{in}}, & \text{if } I_{i,t} > 0 \text{ and } O_{i,t} = 0, \\ s_{i,t}^{\text{out}} H_{i,t}^{\text{out}} + \left(1 - s_{i,t}^{\text{out}}\right) H_{i,t}^{\text{in}}, & \text{if } O_{i,t} > 0 \text{ and } I_{i,t} > 0. \end{cases} \tag{11}$$

**Directional PageRank**

Terminal network importance was measured using PageRank, which considers both the magnitude of a terminal's connections and the importance of the terminals to which it is connected (Page et al., 1999). The measure was calculated separately for importing and exporting roles on the monthly directed and flow-weighted terminal network: import PageRank was calculated using the observed direction of LNG flows, whereas export PageRank was calculated after reversing all edges. Intuitively, import PageRank assigns greater importance to terminals receiving flows from other important terminals, while export PageRank captures the corresponding importance of LNG suppliers.

For the observed network, weighted import PageRank satisfies

$$PR_{i,t}^{\text{imp}} = \frac{1-\alpha}{N_t} + \alpha \sum_{\substack{j \\ (j,i) \in E_t}} PR_{j,t}^{\text{imp}} \frac{q_{ji,t}}{\sum_k q_{jk,t}}, \tag{12}$$

where $N_t$ is the number of active terminals in month $t$, and the damping factor was set to $\alpha = 0.85$, following the conventional value used by Brin and Page (1998). In this formulation, $\alpha$ controls the contribution propagated through the network's directed flows, while $1 - \alpha$ determines the uniform redistribution component. Export PageRank, $PR_{i,t}^{\text{exp}}$, was obtained by applying the same calculation to the reversed

monthly network.

The role-specific PageRank measure was defined as

$$PR_{i,t}^{\text{role}} = \begin{cases} PR_{i,t}^{\text{exp}}, & \text{if } O_{i,t} > 0 \text{ and } I_{i,t} = 0, \\ PR_{i,t}^{\text{imp}}, & \text{if } I_{i,t} > 0 \text{ and } O_{i,t} = 0, \\ s_{i,t}^{\text{out}} PR_{i,t}^{\text{exp}} + \left(1 - s_{i,t}^{\text{out}}\right) PR_{i,t}^{\text{imp}}, & \text{if } O_{i,t} > 0 \text{ and } I_{i,t} > 0. \end{cases} \tag{13}$$

**Trade dependence**

For each directed terminal pair, the dependence of an importing terminal $j$ on an exporter $i$ was measured as the share of $j$'s total imports supplied by $i$:

$$d_{ij,t} = \frac{q_{ij,t}}{I_{j,t}}. \tag{14}$$

To summarise the dependence generated by exporter $i$, this bilateral measure was weighted by the share of the exporter's flow directed towards each destination:

$$D_{i,t}^{\text{exp}} = \sum_{j} \left( \frac{q_{ij,t}}{O_{i,t}} \right) d_{ij,t}. \tag{15}$$

The resulting measure is high when an exporter directs a substantial share of its LNG towards terminals that depend strongly on its supply.

For an importing terminal $i$ located in country $c(i)$, dependence was measured as its share of the total LNG volume received by all terminals in that country:

$$D_{i,t}^{\text{imp}} = \frac{I_{i,t}}{\sum_{k:\, c(k)=c(i)} I_{k,t}}. \tag{16}$$

The role-specific dependence measure was subsequently defined as

$$D_{i,t}^{\text{role}} = \begin{cases} D_{i,t}^{\text{exp}}, & \text{if } O_{i,t} > 0 \text{ and } I_{i,t} = 0, \\ D_{i,t}^{\text{imp}}, & \text{if } I_{i,t} > 0 \text{ and } O_{i,t} = 0, \\ s_{i,t}^{\text{out}} D_{i,t}^{\text{exp}} + \left(1 - s_{i,t}^{\text{out}}\right) D_{i,t}^{\text{imp}}, & \text{if } O_{i,t} > 0 \text{ and } I_{i,t} > 0. \end{cases} \tag{17}$$

**Mean voyage distance**

Mean voyage distance was calculated as the unweighted arithmetic mean of the reconstructed distances of voyages associated with each terminal. For an exporting terminal, this corresponds to

$$\overline{L}_{i,t}^{\text{out}} = \frac{1}{n_{i,t}^{\text{out}}} \sum_{r \in \mathcal{R}_{i,t}^{\text{out}}} L_r, \tag{18}$$

where $\mathcal{R}_{i,t}^{\text{out}}$ is the set of voyages departing from terminal $i$ in month $t$, $n_{i,t}^{\text{out}}$ is the cardinality of this set, and $L_r$ is the reconstructed distance of voyage $r$.

For an importing terminal, the corresponding measure was defined as

$$\overline{L}_{i,t}^{\text{in}} = \frac{1}{n_{i,t}^{\text{in}}} \sum_{r \in \mathcal{R}_{i,t}^{\text{in}}} L_r, \tag{19}$$

where $\mathcal{R}_{i,t}^{\text{in}}$ is the set of voyages arriving at terminal $i$ in month $t$ with cardinality $n_{i,t}^{in}$. The role-relevant mean voyage distance was subsequently lagged by one month before being included among the model covariates.

### 3.4.2 Chokepoint criticality measures

Chokepoint criticality was represented through weighted betweenness, PageRank, and the share of monthly network flow. Together, these measures capture complementary dimensions of a passage's importance, including its position along high-flow paths, its centrality in the direction of LNG movement, and the volume of LNG transported through it.

**Weighted betweenness centrality**

Betweenness centrality measures the extent to which a node lies on shortest paths between other nodes (Brandes, 2001). Since edge weights in the LNG network represent transported volumes rather than geographic or economic distances, they could not be used directly as path lengths. A distance measure was therefore assigned to

each directed edge as the inverse of its positive LNG flow:

$$\ell_{ij,t} = \frac{1}{w_{ij,t}}, \qquad w_{ij,t} > 0. \tag{20}$$

where $w_{ij,t}$ is the LNG volume transported from node $i$ to node $j$ during month $t$. Under this transformation, edges carrying larger LNG volumes are treated as stronger and therefore shorter connections.

Normalised weighted betweenness for node $i$ was then calculated as

$$B_{i,t} = \frac{1}{(N_t - 1)(N_t - 2)} \sum_{\substack{j \neq i \neq k \\ j \neq k}} \frac{\sigma_{jk,t}(i)}{\sigma_{jk,t}}, \tag{21}$$

where $N_t = |V_t|$ is the number of active nodes in month $t$ and $\sigma_{jk,t}$ in the denominator is the number of shortest directed paths from node $j$ to node $k$, calculated using the edge distances defined in Equation 20. The term $\sigma_{jk,t}(i)$ denotes the number of these paths that pass through node $i$. A high value of $B_{i,t}$ indicates that a chokepoint lies on a large share of the shortest paths defined by the distribution of LNG flows and therefore occupies an important intermediary position in the monthly transportation network.

**PageRank**

PageRank provides a complementary measure of chokepoint importance by accounting for both the magnitude and direction of LNG flows and the importance of the nodes from which those flows originate. For node $i$, it is defined as

$$PR_{i,t} = \frac{1 - \alpha}{N_t} + \alpha \sum_{\substack{j \\ (j,i) \in E_t}} PR_{j,t} \frac{w_{ji,t}}{\sum_k w_{jk,t}}, \tag{22}$$

where $\alpha = 0.85$ is the damping factor.

**Share of monthly network flow**

For a terminal, monthly node throughput equals the sum of its incoming and outgoing flows. For a chokepoint, the same shipment appears on both an incoming and an outgoing segment as it passes through the node. Let $V^{\mathrm{CP}} \subseteq V$ denote the set of chokepoint nodes. For each $i \in V^{\mathrm{CP}}$, throughput was defined as half of its total incident flow:

$$F_{i,t} = \frac{1}{2}\left(\sum_{j} w_{ji,t} + \sum_{k} w_{ik,t}\right), \qquad i \in V^{\mathrm{CP}}. \tag{23}$$

The share of monthly network flow associated with node $i$ was then calculated as

$$S_{i,t} = \frac{F_{i,t}}{\sum_{n \in V} F_{n,t}}, \qquad i \in V^{\mathrm{CP}}. \tag{24}$$

This measure represents the share of total node throughput accounted for by a chokepoint in a given month.

### 3.4.3 Common structural indicators

In addition to the node-type-specific criticality measures, a common set of structural indicators was calculated for both terminals and chokepoints. These measures describe whether a node or one of its incident connections contributes to maintaining the connectivity of the monthly transportation network.

The indicators were calculated separately for each month using the undirected projection $U_t$ of the active directed network $G_t$. Edge direction was disregarded because articulation points and bridges concern the preservation of connectivity rather than the direction of LNG movement. Self-loops were excluded because they do not contribute to connectivity.

### Articulation points

A node was classified as an articulation point when its removal increased the number of connected components in the monthly undirected network. The corresponding binary indicator was defined as

$$A_{i,t} = \mathbb{I}\left\{\kappa\left(U_t^{(-i)}\right) > \kappa\left(U_t\right)\right\}, \tag{25}$$

where $\kappa(U_t)$ denotes the number of connected components in $U_t$, and $U_t^{(-i)}$ is the graph obtained by removing node $i$ and its incident edges.

A separate indicator identifies nodes directly connected to an articulation point:

$$CA_{i,t} = \mathbb{I}\left\{\exists\, j \in \mathcal{N}_{i,t} \text{ such that } A_{j,t} = 1\right\}, \tag{26}$$

where $\mathcal{N}_{i,t}$ is the set of immediate neighbours of node $i$ in $U_t$. This distinction separates nodes whose own removal could disrupt network connectivity from nodes directly exposed to a structurally critical neighbour.

### Global and local bridges

An edge was classified as a global bridge when its removal increased the number of connected components in the undirected network. Denoting the set of global bridges in month $t$ by $\mathcal{B}_t^G$, the number of global bridges incident to node $i$ was calculated as

$$GBC_{i,t} = \sum_{e \in \delta_t(i)} \mathbb{I}\left\{e \in \mathcal{B}_t^G\right\}, \tag{27}$$

where $\delta_t(i)$ is the set of edges incident to node $i$ in month $t$. The corresponding binary indicator was defined as

$$GB_{i,t} = \mathbb{I}\left\{GBC_{i,t} > 0\right\}. \tag{28}$$

Local bridges were defined as edges whose endpoints had no common neighbour. Because every global bridge also satisfies this condition, global bridges were removed

from the local-bridge set so that the two measures represented distinct structural mechanisms. Denoting the resulting set of local-only bridges by $\mathcal{B}_t^L$, the corresponding node-level count was calculated as

$$LBC_{i,t} = \sum_{e \in \delta_t(i)} \mathbb{I}\left\{e \in \mathcal{B}_t^L\right\}. \tag{29}$$

To account for differences in node degree, the shares of incident edges classified as global and local-only bridges were also calculated:

$$\begin{aligned} GBS_{i,t} &= \frac{GBC_{i,t}}{d_{i,t}}, \\ LBS_{i,t} &= \frac{LBC_{i,t}}{d_{i,t}}, \end{aligned} \qquad d_{i,t} > 0, \tag{30}$$

where $d_{i,t}$ is the degree of node $i$ in the monthly undirected network. Together, these measures distinguish connections whose removal would disconnect part of the network from connections that locally bridge otherwise weakly connected neighbourhoods.

## 3.5 Construction of the model-ready panels

Three model-ready panels were constructed from the monthly network data described above: a node activity panel, a terminal criticality panel, and a chokepoint criticality panel. Each panel has a balanced structure in which every unit is represented in every month from January 2020 to December 2024, including months with no observed activity. This preserves transitions between active and inactive states and provides a consistent temporal basis for the subsequent models.

For each node, one-month lags of activity, throughput, and voyage count were calculated from the corresponding values in the preceding month. Variables with strongly right-skewed distributions were also included in logarithmic form using the transformation $\log(1 + x)$, which retains zero-valued observations. Depending on the panel, these variables comprise lagged throughput, lagged voyage count, terminal processing capacity, and lagged mean voyage distance.

Given the temporal coverage of the data, lagged variables are unavailable for January 2020. The binary indicator `lag_available` identifies observations for which the required preceding-month information can be calculated.

Lagged neighbourhood activity was constructed using an undirected historical neighbourhood: for each month $t$, the neighbours of a node are those connected to it by at least one directed edge observed up to and including month $t-1$. The resulting variables record the number of historical neighbours, the number that were active in $t-1$, and the corresponding active share. Historical neighbourhoods were used because defining a node's neighbourhood exclusively from edges observed in $t-1$ would make every identified neighbour active by construction, causing the active-neighbour share to be mechanically equal to one. The cumulative definition instead retains previously observed neighbours regardless of whether they were active in $t-1$, thereby capturing variation in the recent activity of a node's established connections. Connections first observed after $t-1$ are excluded to prevent the use of future information: neighbourhood-activity measures remain unavailable until a node has acquired at least one historical neighbour.

Finally, the common set of model covariates included seasonal terms, represented through sine and cosine transformations of the calendar month. This cyclical representation allows December and January to be treated as adjacent rather than as the endpoints of a linear sequence.

A complete description of the variables included in each model-ready panel, together with their transformations and availability, is provided in Appendix C.

### 3.5.1 Node activity panel

The node activity panel contains all 183 network nodes, comprising 159 LNG terminals and 24 maritime chokepoints, observed over 60 months, resulting in $183 \times 60 = 10{,}980$ node–month observations.

The response variable is the binary indicator of current node activity. The model

covariates comprise preceding-month activity, throughput, voyage count, historical-neighbour activity, and the seasonal terms defined above. Time-invariant or slowly varying node characteristics were obtained from the consolidated node inventory and include node type and, where applicable, country, infrastructure type, start year, and annual processing capacity. Months without observed participation were retained with `active` equal to zero and the corresponding flow and voyage measures set to zero.

### 3.5.2 Terminal criticality panel

The terminal criticality panel is restricted to the 159 LNG terminals, producing $159 \times 60 = 9{,}540$ terminal–month observations.

Its outcomes comprise the terminal criticality measures defined in Section 3.4.1, namely role-specific PageRank, role-specific counterparty concentration at the terminal and country levels, and role-specific dependence. They also include the common structural indicators defined in Section 3.4.3, which identify whether a terminal is an articulation point, is incident to a global bridge, or is directly connected to an articulation point.

Criticality measures based on realised commercial activity are defined only for active terminal–months and remain unavailable for inactive observations. Structural indicators, by contrast, are retained for every terminal–month and are equal to zero when the terminal does not belong to the active monthly network.

The terminal panel additionally includes the preceding-month terminal role and role-relevant mean voyage distance, together with country, infrastructure type, annual processing capacity, number of infrastructure units, start year, and terminal age. Continuous covariates are retained in both their original and, where appropriate, log-transformed forms.

### 3.5.3 Chokepoint criticality panel

The chokepoint criticality panel contains the 24 chokepoints included in the final network over the same 60-month period, yielding $24 \times 60 = 1{,}440$ chokepoint–month

observations.

The outcomes comprise weighted betweenness centrality, PageRank, and the share of monthly network flow defined in Section 3.4.2, together with the common structural indicators introduced in Section 3.4.3. Additional measures of degree, route exposure, and voyage traversals were retained as supplementary network characteristics.

Finally, the panel includes the common lagged and seasonal covariates described above, together with lagged mean voyage distance for voyages traversing each chokepoint.

## 3.6 Data and methodological limitations

The construction of the LNG maritime network is subject to limitations arising both from the underlying LNG-T3 data and from the procedures used to reconstruct routes and derive network measures.

As discussed by Zhou et al. (2026), LNG-T3 relies extensively on Automatic Identification System messages, which are self-reported by vessels and may contain errors, delays, or incomplete information. AIS coverage also varies geographically according to the availability and density of receiver stations. Coverage is generally more extensive in Europe and North America than in regions such as Southeast Asia, where archipelagic geography and a lower density of receivers may generate spatial blind spots. These differences can introduce geographic variation in data quality.

Incomplete AIS trajectories and missing port-call records may result in partially reconstructed or unobserved voyages, while inaccurate draught reporting may affect the classification of whether a tanker is loaded. To mitigate these sources of uncertainty, LNG-T3 adopts a conservative validation procedure and retains only voyages considered suitable for estimating trade flows. Although this improves the reliability of the individual records, it may result in a systematic underestimation of aggregate LNG trade relative to industry and government statistics (Zhou et al., 2026).

Additional uncertainty concerns the estimation of cargo volumes. LNG-T3 assumes that tankers depart fully loaded, which provides a consistent basis for global

aggregation but does not capture possible partial loadings associated with operational constraints, contractual arrangements, or market conditions. Estimated volumes may therefore overstate the cargo transported on some individual voyages. At the aggregate level, however, this potential upward bias coexists with the potential downward bias arising from incomplete voyage coverage, and their combined effect cannot be determined unambiguously.

A further limitation arises from the reconstruction of maritime routes, as the available voyage records identify departure and arrival terminals but do not contain the complete trajectories followed by individual tankers. The routes used in this study, generated using `SeaRoute`, are modelled approximations rather than documented voyage trajectories and may not reproduce routing decisions influenced by weather, congestion, regulatory restrictions, security conditions, canal availability, or commercial considerations.

Small spatial discrepancies between the reconstructed paths and the operational chokepoint geometries were addressed through a uniform 25-km tolerance. However, the sensitivity analysis reported in Appendix B shows that the assignments for Korea Strait and Mindoro Strait are particularly dependent on the selected tolerance. Chokepoint exposure should therefore be interpreted as an approximation derived from reconstructed routes rather than direct evidence that individual vessels crossed each passage.

As a final clarification of scope, the resulting network represents physical LNG flows observed during 2020–2024 and the transportation paths reconstructed for those flows. Accordingly, the resulting measures characterise realised network activity and estimated route exposure within the study period.

# Chapter 4

# Exploratory network analysis

This chapter examines the structure and evolution of the reconstructed LNG maritime network from January 2020 to December 2024, by describing changes in monthly trade flows and connectivity, patterns of node activity, and the network positions of terminals and chokepoints using the measures introduced in Chapter 3. The analysis provides a descriptive overview of the main temporal and cross-sectional patterns in the data and examines the relationships among the candidate variables, to identify relevant patterns, potential redundancy, and shared variation that may inform the subsequent modelling analysis.

To address these objectives, the chapter is organised as follows. Section 4.1 examines the temporal evolution of the monthly LNG network; Section 4.2 then describes node activity and temporal sparsity over the observation period; Section 4.3 focuses on terminals, considering their observed trade roles and the measures of centrality, concentration, and dependence associated with their network positions; Section 4.4 examines chokepoints through their flow exposure, PageRank, and weighted betweenness; Section 4.5 compares the structural positions of terminals and chokepoints using articulation-point and bridge-based measures; finally, Section 4.6 examines associations among the candidate variables and discusses their implications for the subsequent modelling analysis.

As described in Section 3.2.2, the node–month and terminal–month datasets retain

both active and inactive observations. In this chapter, summary statistics for network-position measures are generally calculated over active observations, while analyses of node activity and temporal sparsity retain inactive months. The relevant sample is specified in the text or table notes whenever necessary.

## 4.1 Network evolution

The observed LNG maritime network changes in size over the study period. The numbers of active nodes and directed edges generally increase from 2021 to late 2023, reaching their respective maxima in December 2023, before declining during the second half of 2024. The ranges and averages of these network characteristics, together with monthly trade activity and directed density, are summarised in Table 2.

Despite this evolution in network size, directed density remains within a relatively narrow range, ranging from 0.0122 to 0.0180. The monthly LNG network is therefore consistently sparse: only a small fraction of all possible directed connections among active nodes is observed in any given month.

Directed density is calculated among active nodes, so changes in its denominator must be considered when interpreting the results: a month with fewer edges can still have higher density if fewer nodes are active. Network expansion or contraction should therefore not be interpreted as a corresponding change in the degree of connectivity among active nodes.

Table 2: Selected monthly network and node-activity summaries, 2020–2024.

| Measure | Minimum | Mean | Maximum |
|---|---|---|---|
| Active nodes | 72 | 96.5 | 126 |
| Active directed edges | 88 | 136.5 | 192 |
| Export voyages | 70 | 144.0 | 237 |
| Exported volume (MCM) | 11.48 | 24.01 | 38.77 |
| Directed density among active nodes | 0.0122 | 0.0148 | 0.0180 |

*Note:* Exported volume is expressed in million cubic metres (MCM).

While Table 2 summarises the monthly networks quantitatively, Figure 2 illustrates the geographic structure of a representative month. June 2021 was selected as an

illustrative snapshot because its 98 active nodes and 138 directed edges are close to the corresponding monthly averages reported in Table 2.




Figure 2: Geographic snapshot of the observed LNG network in June 2021.

*Note:* Points show active terminals and chokepoints; arrows indicate the direction of observed network connections. Line width represents monthly LNG volume on each directed edge; connections are schematic rather than sailed routes.

To examine how the network's size and trade volume evolve over time, Figure 3 compares the monthly evolution of active nodes, directed edges, and observed LNG export volume, with each series indexed to its mean over the study period.

The three measures show broadly similar changes over time, but export volume exhibits substantially greater month-to-month variation, showing that changes in network size are not necessarily proportional to changes in the trade volume carried by the network.

## 4.2 Node activity and temporal sparsity

Using the definition of node activity introduced in Section 3.3.2, of 10,980 potential node-month observations, 5,792 (52.75%) are active and 5,188 (47.25%) inactive.

As shown in Figure 3, the number of active nodes generally increases through 2023 and declines towards the end of 2024. Correspondingly, the monthly share of inactive nodes ranges from 31.15% to 60.66% (Figure 4).




Figure 3: Monthly evolution of network size and observed LNG export volume, 2020–2024.

*Note:* Each series is indexed to its monthly mean over the 2020–2024 period, which is set equal to 100. Export volume is calculated from the observed export voyages.

Terminals account for 9,540 node–month observations, of which 4,552 are active and 4,988 inactive (52.29%). The 24 chokepoints account for 1,440 observations, of which 1,240 are active and 200 inactive (13.89%). This difference is consistent with the nodes' distinct roles in the network: a terminal participates as an origin or destination of a voyage, whereas a chokepoint can be traversed by voyages linking many different terminal pairs. Aggregate inactivity is therefore primarily attributable to terminals, although chokepoint activity is not constant. These different base rates suggest that activity may need to be examined separately by node type.




Figure 4: Monthly inactive-node shares for all nodes and by node type.

Inactivity is substantively relevant to the interpretation of centrality: an inactive node may receive a zero in the balanced node-month dataset even though the criticality measure is chiefly meaningful when it participates in trade. For this reason, the subsequent analysis of node criticality focuses on active node-month observations, avoiding the conflation of inactivity with low criticality.

## 4.3 Terminal network position

This section examines how terminals participate in observed LNG trade and how their positions differ across the monthly networks. It first describes terminal roles and trade activity, then considers centrality, counterparty concentration, and dependence.

### 4.3.1 Terminal roles and trade activity

Terminal participation varies considerably across the 60 monthly networks: of the 4,552 active terminal–month observations, 3,234 are classified as importers and 1,318 as exporters.

Country-level roles were assessed over the full 2020–2024 period using each country's net trade position. Qatar, the United States, and Russia recorded the largest cumulative export volumes, while China, France, and Japan recorded the largest cumulative import volumes. A net importer or exporter role does not imply that a country trades exclusively in one direction. For instance, Malaysia was a net importer despite recording exports, and the United States was a net exporter despite recording imports.

Table 3 compares throughput, voyage counts, and the numbers of terminal counterparties for active terminal–month observations, distinguished by importer and exporter role.

Exporter observations are characterised by higher levels of trade activity across all three measures. Median monthly throughput is 0.505 million cubic metres for exporters, compared with 0.328 million cubic metres for importers, while the corresponding median numbers of voyages are four and two. Exporters also interact with

Table 3: Trade activity of active terminal–month observations by terminal role, 2020–2024.

| | Importer ($n = 3{,}234$) | | Exporter ($n = 1{,}318$) | |
|---|---|---|---|---|
| Measure | Median | 95th percentile | Median | 95th percentile |
| Throughput (million cubic metres) | 0.328 | 1.226 | 0.505 | 3.792 |
| Voyages | 2 | 7 | 4 | 22 |
| Terminal counterparties | 1 | 4 | 3 | 14 |

The analysis is restricted to active terminal–month observations. Throughput values are rounded to three decimal places.

a larger number of terminal counterparties, with a median of three compared with one for importers. These differences are even more pronounced at the 95th percentile, particularly for voyage counts and numbers of counterparties.

More generally, throughput is strongly right-skewed across active terminal–month observations. Its overall median is 346,000 cubic metres, and the 95th percentile is approximately 1.90 million cubic metres, while Rasgas LNG Terminal 3 reaches the observed maximum of 10.91 million cubic metres in October 2023.

### 4.3.2 Centrality, concentration and dependence

The terminal measures discussed below were defined in Section 3.4.1. Here, their distributions are examined across active terminal–month observations to explore different aspects of terminals' positions and trade relationships in the observed LNG network. These pooled distributions are complemented by terminal-level summaries of the monthly measures over the entire period, which distinguish the positions of individual terminals and help assess whether high scores are sustained over time or occur only occasionally.

First, role-specific PageRank has a median of 0.0142 and a 95th percentile of 0.0363. The 95th percentile is approximately 2.6 times the median, indicating substantial variation in the scores assigned to active terminals. The highly right-skewed distribution suggests that a relatively small number of terminal–month observations occupy particularly prominent positions within the monthly trade network, while most receive

lower scores.

These scores should nevertheless be interpreted as relative measures within each monthly network. Because PageRank is normalised separately each month, its values are influenced by both the distribution of trade relationships and the number of active terminals. Comparisons across months therefore describe changes in relative network prominence rather than changes in an absolute measure of importance.

To examine terminal positions over the full period, two summaries are considered: mean PageRank conditional on activity and mean PageRank across all 60 months, assigning zero to inactive months. The first describes a terminal's average prominence when participating in the network; the second combines that prominence with its frequency of participation.

Table 4 reports the three highest full-period mean scores within each terminal role. Rasgas LNG Terminal 3 has the highest mean PageRank among exporters, at 0.0912, followed by Sabine Pass and Yamal, at 0.0449 and 0.0418, respectively. All three terminals are active in every month, so their mean scores reflect sustained network participation over the full observation period.

Among importers, Gate LNG Terminal has the highest full-period mean, at 0.0195, followed by Yung-An and Fos Tonkin. Fos Tonkin has a slightly higher mean during active months than Yung-An, but its lower number of active months reverses their ordering when inactivity is included. The distinction is more pronounced for terminals observed only occasionally: Sepetiba Bay FSRU has an active-month mean of 0.0191, close to Gate's, but is active in only one month. Consequently, its mean PageRank across the full panel is approximately 0.00032, highlighting how a high conditional mean alone should therefore not be interpreted as evidence of sustained prominence.

PageRank is relevant to the assessment of terminal prominence, but does not establish how easily a terminal could be replaced. A well-positioned terminal may trade with partners that have alternative sources or destinations, whereas a less prominent terminal may serve counterparties that rely heavily on its supply or handle a substantial share of its country's LNG receipts. Concentration and dependence are therefore

Table 4: Terminals with the highest mean role-specific PageRank across all 60 months, ranked separately by observed role.

| Terminal | Country | Active months | Mean PageRank | |
|---|---|---|---|---|
| | | | Active months | All months |
| *Exporters* | | | | |
| Rasgas LNG Terminal 3 | Qatar | 60 | 0.0912 | 0.0912 |
| Sabine Pass LNG Terminal | United States | 60 | 0.0449 | 0.0449 |
| Yamal LNG Terminal | Russia | 60 | 0.0418 | 0.0418 |
| *Importers* | | | | |
| Gate LNG Terminal | Netherlands | 60 | 0.0195 | 0.0195 |
| Yung-An LNG Terminal | Taiwan | 57 | 0.0182 | 0.0172 |
| Fos Tonkin LNG Terminal | France | 55 | 0.0187 | 0.0171 |

*Note:* Arithmetic means of monthly PageRank scores. Active-month means exclude inactivity; all-month means assign zero to inactive months.

examined alongside PageRank to assess the trade relationships underlying a terminal's network position.

The counterparty concentration measures show substantial variation among active terminal–month observations. The median HHI is 0.556 when counterparties are individual terminals and 0.674 when they are aggregated by country. These median values correspond to approximately 1.8 and 1.5 equally weighted effective counterparties at the terminal and country levels, respectively. Thus, even when a terminal trades with several partners, its observed monthly flow may be concentrated among relatively few of them.

An HHI of one occurs in 43.5% of active terminal–month observations at the terminal level, indicating that the role-relevant flow involves a single terminal counterparty. At the country level, the corresponding proportion is 47.4%, indicating concentration in a single country, potentially across multiple terminals.

Terminal-level averages reveal different patterns behind these pooled results. As shown in Table 5, Rasgas and Sabine Pass have relatively low mean terminal-level HHIs, at 0.094 and 0.125, respectively, despite their prominent PageRank positions.

Their observed monthly exports are therefore distributed more broadly across counterparties than the pooled distribution might suggest. In contrast, Cameroon FLNG has a mean terminal-level HHI of 0.953 across 32 active months, indicating highly concentrated shipments during its observed activity.

Country-level concentration also distinguishes diversification across terminals from diversification across countries. For instance, the mean HHI of Gate increases from 0.362 at the terminal level to 0.531 at the country level, showing that some diversification across supplying terminals occurs within the same countries.

These findings suggest that an interruption involving a major partner could affect a substantial share of some terminals' existing trade. The measures nevertheless describe observed monthly relationships and do not establish whether alternative suppliers or destinations would be available. Moreover, the terminal-level averages summarise monthly concentration: they do not imply that a terminal trades with the same counterparties throughout the period.

Table 5: Mean counterparty concentration and role-specific dependence during active months for selected terminals, 2020–2024.

| Terminal | Active months | Mean HHI | | Mean dependence |
|---|---|---|---|---|
| | | Terminal | Country | |
| *Exporters* | | | | |
| Rasgas LNG Terminal 3 | 60 | 0.094 | 0.139 | 0.730 |
| Sabine Pass LNG Terminal | 60 | 0.125 | 0.179 | 0.508 |
| Cameroon FLNG Terminal | 32 | 0.953 | 0.969 | 0.350 |
| *Importers* | | | | |
| Gate LNG Terminal | 60 | 0.362 | 0.531 | 0.910 |
| Al Zour LNG Import Terminal | 40 | 0.652 | 0.657 | 1.000 |
| Zeebrugge LNG Terminal | 59 | 0.558 | 0.599 | 1.000 |

*Note:* Terminals are selected to illustrate contrasting relationships, rather than to provide a ranking. All values are unweighted arithmetic means across active months.

Finally, role-specific dependence has a median of 0.406 across active terminal–month observations, but this pooled value combines different interpretations for the two terminal roles. For exporters, which have a median of 0.495, higher values indicate

that a larger share of their flow reaches importing terminals that rely heavily on its supply. An exporter can therefore have high dependence even when its shipments are distributed across several destinations, provided that it supplies a substantial share of those destinations' imports. Rasgas illustrates this combination: its low mean terminal-level HHI of 0.094 coexists with mean dependence of 0.730. Its exports are relatively diversified, yet its receiving terminals tend to rely substantially on its supply when weighted by their shares of its shipments. Cameroon FLNG illustrates the opposite pattern: its highly concentrated exports coexist with lower mean dependence of 0.350. Concentration in an exporter's own destinations therefore does not necessarily imply equally strong reliance by those destinations on its supply.

Among active importer months, the median dependence is 0.333. Here, higher values indicate that the terminal accounts for a larger share of its country's observed LNG receipts. A high value therefore identifies a terminal through which a substantial proportion of national LNG imports passes, making its continued operation potentially relevant to the country's ability to receive LNG.

The terminal-level summaries illustrate the extent of this concentration at individual terminals. Gate accounts for an average of 91.0% of the Netherlands' observed monthly LNG receipts across its 60 active months, while Al Zour and Zeebrugge have dependence equal to one in all their active months, numbering 40 and 59, respectively. In those months, all observed LNG receipts in their respective countries pass through these terminals.

The crucial distinction from the counterparty HHI measure described above lies in whose reliance is being described: HHI quantifies the concentration of a terminal's own trade across its partners, whereas role-specific dependence describes its counterparties' reliance on its supply or its country's reliance on that terminal for observed LNG receipts. An importer may receive all its LNG from one exporter, giving it an HHI of one, while handling only a small share of its country's total imports and therefore having low role-specific dependence. Conversely, a terminal may receive LNG from several suppliers while accounting for a large share of national LNG receipts. Gate

illustrates the latter pattern, combining a mean terminal-level HHI of 0.362 with mean dependence of 0.910.

Concentration and dependence are therefore retained alongside PageRank to capture distinct dimensions of terminal criticality: partner concentration, reliance on individual terminals, and prominence within the trade network. Together, these measures characterise terminal criticality without directly quantifying the consequences of a disruption.

## 4.4 Chokepoint network position

Unlike terminals, which serve as the origins or destinations of LNG voyages, chokepoints act as intermediate nodes along maritime routes. Their position can therefore be examined in terms of both exposure to LNG flows and location within the wider network, captured by PageRank and betweenness.

Table 6 provides summary statistics for the main flow-exposure and centrality measures considered in this section.

Table 6: Summary statistics for flow-exposure and centrality measures across active chokepoint–month observations, 2020–2024.

| Measure | Mean | Median | 95th percentile | Maximum |
|---|---|---|---|---|
| Throughput (MCM) | 2.007 | 1.342 | 5.857 | 10.702 |
| Voyages | 11.08 | 8 | 31 | 55 |
| Route exposure | 8.30 | 7 | 21 | 31 |
| Share of monthly network flow | 0.0225 | 0.0148 | 0.0639 | 0.1132 |
| PageRank | 0.0172 | 0.0163 | 0.0299 | 0.0572 |
| Betweenness | 0.0126 | 0.0072 | 0.0408 | 0.0946 |

*Note:* Statistics are calculated across the 1,240 active chokepoint–month observations. Throughput is reported in million cubic metres.

### 4.4.1 Flow exposure

Flow exposure varies substantially across the 24 chokepoints and over months. As shown in Table 6, the distribution of monthly throughput is strongly right-skewed,

with the observed maximum of 10.70 million cubic metres recorded at the Strait of Hormuz in April 2022.

The same heterogeneity appears when flow is expressed relative to the total flow observed in the monthly network. Among active observations, the share of monthly network flow passing through a chokepoint has a median of approximately 1.48%, while the 95th percentile reaches 6.39%. The maximum is recorded at the Strait of Hormuz in June 2020, when approximately 11.32% of the month's observed LNG trade passed through the chokepoint.

The flow-share measure complements the absolute volume of LNG passing through a chokepoint. Because monthly LNG trade varies over the observation period, a given throughput may represent different levels of exposure depending on the total volume traded in that month. Expressing chokepoint throughput as a share of total monthly network flow therefore provides a relative measure of how much of the observed network flow is routed through each passage.

Beyond flow volume, voyage and route counts provide additional information on the frequency and variety of chokepoint use. Active chokepoints are traversed by a median of eight voyages per month and are exposed to a median of seven distinct routes, while the corresponding 95th percentiles are 31 voyages and 21 routes.

### 4.4.2 Centrality measures

Flow exposure alone does not describe the structural location of a chokepoint within the reconstructed network. Two passages carrying similar LNG volumes may connect different parts of the network and, consequently, differ in their centrality. PageRank and betweenness are therefore considered alongside flow exposure to capture these complementary aspects of chokepoint prominence.

As defined in Section 3.4.2, PageRank accounts for both the magnitude and direction of LNG flows and the importance of the nodes from which those flows originate.

It therefore complements flow exposure by capturing a chokepoint's relative position within each monthly network rather than simply the amount of LNG passing

through it. The distribution of raw PageRank values indicated relatively prominent positions for some chokepoints, with a median of 0.0163 among active observations and values reaching 0.0572. This motivated a closer examination of their magnitude relative to the size of each monthly network. Observed PageRank was therefore compared with a uniform benchmark of $1/N_t$, corresponding to an equal distribution of PageRank across the $N_t$ active nodes.

Among active observations, the median PageRank was 1.54 times this monthly benchmark, and 85.6% of observations exceeded it. Active chokepoints therefore tended to receive a larger share of PageRank than would result from an equal distribution across active nodes.

While PageRank captures relative prominence within the flow network, betweenness focuses on a different aspect of centrality by indicating the extent to which a chokepoint lies on shortest paths connecting other nodes. A chokepoint may therefore receive a relatively high PageRank without occupying a particularly strong intermediary position, while another may have a lower PageRank but lie on a comparatively large number of shortest paths.

As shown in Table 6, its distribution is strongly right-skewed, with high levels of shortest-path intermediation concentrated in a comparatively small subset of active chokepoint–month observations.

Notably, positive activity does not necessarily imply positive betweenness: betweenness is equal to zero in 8 of the 1,240 active chokepoint–month observations. A chokepoint may therefore be traversed by observed LNG routes without lying on any of the shortest paths between other node pairs captured by this measure.

The distinction between the three measures is also visible when their average values are compared across individual chokepoints. Table 7 reports four passages selected to illustrate contrasting combinations of flow exposure, PageRank, and betweenness. All four are active throughout the 60-month observation period, so differences in their averages are not driven by different frequencies of activity.

The Strait of Hormuz has the largest mean flow share among the selected passages,

Table 7: Selected chokepoints illustrating different dimensions of network position, 2020–2024.

| Chokepoint | Active months | Mean flow share | Mean PageRank | Mean betweenness |
|---|---|---|---|---|
| Strait of Hormuz | 60 | 0.0766 | 0.0182 | 0.0181 |
| Gibraltar Strait | 60 | 0.0438 | 0.0263 | 0.0360 |
| Taiwan Strait | 60 | 0.0364 | 0.0309 | 0.0287 |
| Suez Canal | 60 | 0.0327 | 0.0211 | 0.0301 |

*Note:* Chokepoints are selected to illustrate contrasting relationships among the three measures rather than to provide a ranking. Means are calculated over active months.

at 7.66%, but substantially lower mean PageRank and betweenness than some passages carrying smaller shares of monthly network flow. The Taiwan Strait, by contrast, has a mean flow share of 3.64% but the highest mean PageRank of the four, at 0.0309. Gibraltar Strait combines a lower mean flow share than Hormuz, at 4.38%, with the highest mean betweenness, at 0.0360, while the Suez Canal also exhibits relatively high betweenness despite accounting for a mean 3.27% of monthly network flow.

These comparisons show that greater flow exposure does not necessarily correspond to higher PageRank or betweenness, highlighting the distinct dimensions of chokepoint criticality captured by these measures.

## 4.5 Structural position of nodes

The structural indicators defined in Section 3.4.3 complement the node-type-specific measures by describing connectivity in the monthly undirected projection of the reconstructed network. They distinguish nodes whose removal would disconnect parts of the network from nodes that are incident to structurally important edges or adjacent to articulation points. Rather than reflecting the volume of LNG transported through individual connections, these indicators depend only on their presence in the monthly network.

The distributions differ substantially between terminals and chokepoints. Articulation status occurs in 393 of the 4,552 active terminal–month observations (8.6%),

compared with 902 of the 1,240 active chokepoint–month observations (72.7%). Incidence to at least one global bridge is common in both groups, occurring in 64.7% of active terminal months and 72.5% of active chokepoint months. Local-only bridge incidence is more frequent among chokepoints, at 82.8%, than among terminals, at 26.1%.

These differences are consistent with the roles of terminals and chokepoints in the network: terminals are voyage origins or destinations, whereas chokepoints are intermediate nodes connecting successive route segments. Chokepoints consequently have more opportunities to connect branches of the observed network, although their classifications depend on the routes used in each month.

The analysis of articulation points further illustrates this distinction. Among terminals, Vysotsk repeatedly acts as an articulation point, being classified as an articulation point in 49 of its 59 active months. Among chokepoints, Gibraltar Strait, Taiwan Strait and the Strait of Hormuz are articulation points in all 60 months. Removing the node therefore disconnects the monthly undirected network in every observed month.

Table 8: Frequency of structural classifications for selected terminals and chokepoints, 2020–2024.

| Node | Activemonths | Articulationpoint | Globalbridge | Local-onlybridge |
|---|---|---|---|---|
| *Terminals* | | | | |
| Vysotsk LNG Terminal | 59 | 49 | 58 | 0 |
| Dunkirk LNG Terminal | 56 | 0 | 56 | 0 |
| Nigeria LNG Terminal | 59 | 21 | 21 | 59 |
| Sabine Pass LNG Terminal | 60 | 24 | 25 | 58 |
| *Chokepoints* | | | | |
| Gibraltar Strait | 60 | 60 | 60 | 58 |
| Strait of Hormuz | 60 | 60 | 58 | 36 |
| Malacca Strait | 60 | 54 | 54 | 60 |
| Suez Canal | 60 | 46 | 46 | 59 |

*Note:* Entries report the number of active months in which each classification applies. Global and local-only bridge columns indicate incidence to at least one edge of the corresponding type. Classifications can coexist at a node, although global bridges are excluded from the local-only edge set. Examples illustrate contrasting patterns rather than a ranking.

Since a global bridge is an edge whose removal separates its endpoints into different

connected components, incidence to such an edge can indicate a connection between groups of nodes, but it can also describe the attachment of a peripheral node to the rest of the network.

Dunkirk LNG Terminal provides a clear example of the latter case. In all its 56 active months, it has one neighbour in the undirected transport network and its sole incident edge is a global bridge. It is nevertheless never an articulation point: removing Dunkirk does not split the remaining network, whereas removing its incident edge isolates Dunkirk. Its classification therefore describes peripheral attachment rather than a role in connecting other nodes.

Conversely, a node can be an articulation point without being incident to a global bridge. The Strait of Hormuz is an articulation point in all 60 months but is incident to a global bridge in only 58. In the remaining two months, removing the node separates the network even though removing any individual incident edge does not. This distinction illustrates why node-removal and edge-removal indicators provide different information.

Local-only bridges identify another structural feature. Their endpoints have no common neighbour, but removing the edge does not disconnect the component because a longer alternative path remains. They therefore indicate the absence of an immediate two-edge alternative through a shared neighbour, rather than an edge that is indispensable to overall connectivity.

Nigeria LNG Terminal is incident to at least one local-only bridge in all its 59 active months, and Sabine Pass in 58 of 60. Among chokepoints, Malacca Strait has local-only bridge incidence in every month, while Suez Canal has it in 59 of 60 months (Table 8).

These classifications can coexist with articulation status or global bridge incidence because a node may have several incident edges with different structural properties. For instance, Malacca Strait is an articulation point and is incident to a global bridge in 54 months, while also being incident to a local-only bridge in all 60 months. This combination indicates that some connections have longer alternative paths, whereas

other connections, or the node as a whole, maintain connectivity between otherwise separated parts of the network. It is to be noted that the alternative paths identified in the undirected projection are not necessarily feasible substitutes for the original LNG movements: these indicators do not account for direction, navigability or transport capacity.

These results distinguish the frequent intermediate connectivity roles of chokepoints from the more varied structural positions of terminals.

## 4.6 Associations and modelling implications

Spearman rank correlations provide a preliminary assessment of the overlap among variables considered in the subsequent modelling analysis.

Among the terminal covariates, lagged log-transformed throughput is strongly associated with lagged log-transformed voyage count ($\rho_S = 0.989$), whereas its associations with other terminal characteristics are substantially weaker. Its correlation with lagged log-transformed mean voyage distance is 0.223 when the latter is defined, while the corresponding correlations with terminal capacity and terminal age are 0.423 and 0.199, respectively. Neighbour activity is also only weakly associated with these node-specific characteristics. These patterns suggest that, beyond the strong association between throughput and voyage count, the candidate covariates capture aspects of terminal activity that cannot be explained solely by operational scale.

A similar pattern is observed among the chokepoint covariates. Lagged log-transformed throughput is strongly associated with lagged log-transformed voyage count ($\rho_S = 0.992$), whereas its associations with the lagged share of active neighbours ($\rho_S = 0.118$) and lagged log-transformed mean voyage distance ($\rho_S = -0.057$) are substantially weaker. The seasonal terms also show little association with lagged throughput. Beyond the close correspondence between throughput and voyage count, the chokepoint covariates therefore capture distinct aspects of activity, route characteristics, and temporal variation.

The associations among the criticality measures differ between terminals and chokepoints. Among active terminal–month observations, terminal- and country-level counterparty HHIs are strongly correlated ($\rho_S = 0.920$), reflecting their related but distinct measures of trade concentration. Their associations with role-specific PageRank are negative and more moderate in magnitude ($-0.596$ and $-0.545$, respectively), while role-specific dependence is more weakly associated with the other terminal measures.

For active chokepoint–month observations, the three continuous measures show stronger positive associations. Flow share correlates with weighted betweenness at 0.810 and with PageRank at 0.704, while the correlation between PageRank and betweenness is 0.706. Thus, although the individual comparisons in Section 4.4 show that greater flow exposure does not necessarily correspond to higher centrality, the aggregate associations indicate that chokepoints carrying larger shares of observed LNG flows tend to have higher PageRank and betweenness values. The correlations remain below one, consistent with the measures capturing related but distinct dimensions of chokepoint criticality.

Figure 5 summarises the pairwise associations among the terminal and chokepoint criticality measures. Taken together, these patterns provide descriptive evidence of shared variation across multiple dimensions of node criticality while preserving meaningful differences between the individual measures. These findings provide the empirical motivation for investigating whether the measures' joint variation can be represented by a lower-dimensional structure in the modelling framework developed in Chapter 5.




(a) Terminal criticality measures

(b) Chokepoint criticality measures

Figure 5: Spearman rank correlations among the criticality measures for active node–month observations. Panel (a) reports correlations among terminal measures, while panel (b) reports correlations among chokepoint measures.

# Chapter 5

# Bayesian Reduced-Rank model for network measures

This chapter develops a Bayesian reduced-rank framework to investigate common patterns across multiple measures of nodes' positions within the network and their associations with node characteristics and lagged network conditions.

Separate rank-one latent factor models are estimated for terminals and chokepoints, reflecting their different functions and response measures within the LNG maritime network. Each model includes its own set of response variables, through a shared latent factor, modelled as a function of the corresponding set of covariates.

The analysis focuses on node-month observations for which activity is observed in consecutive months, using a static specification to examine associations between network position and the selected covariates. The resulting latent factors provide node-specific summaries of network position. Their interpretation and adequacy are evaluated through posterior inference, predictive checks and a prior sensitivity analysis.

## 5.1 General modelling framework

### 5.1.1 Estimation samples, responses and covariates

The terminal model-ready panel contains 9,540 observations for 159 terminals from January 2020 to December 2024.

For the current specification, an observation is retained when the terminal is active in month $t$, a lag from the preceding calendar month is available, and the terminal

was also active in that month. This restriction allows the analysis to focus on network position among consecutively active terminals, using information from the preceding month without modelling transitions between active and inactive states.

After excluding 4,988 inactive observations, 71 active observations without an available calendar lag, and 916 observations preceded by inactivity the sample contains 3,565 observations active in consecutive months. A further 112 observations with missing terminal age are excluded, resulting in an estimation sample of size 3,453 observations for 137 terminals. Of these observations, 2,325 have lagged importer status and 1,128 have lagged exporter status.

The terminal model includes six responses: three bounded measures (role-specific PageRank, terminal counterparty HHI and role-specific dependence) and three binary indicators (articulation-point status, incident-global-bridge status and connection to an articulation point). These measures are expected to capture different aspects of network position and need not all increase with the same latent dimension.

Seven predictors are included: lagged historical-neighbour activity share, log-transformed lagged mean voyage distance, log-transformed lagged terminal capacity, monthly sine and cosine terms, terminal age, and an indicator for exporter status in the preceding month. Numeric covariates are standardised, while the exporter indicator remains on its zero–one scale. Accordingly, coefficients for numeric predictors correspond to one-standard-deviation differences, whereas the role coefficient compares lagged exporters with lagged importers, holding other covariates fixed.

The bounded responses are modelled on their original scales. PageRank lies strictly within $(0, 1)$, whereas HHI and dependence take values in $(0, 1]$. Neither HHI nor dependence has observed zeros, but HHI equals one in 34.09% of observations and dependence equals one in 20.76%. These features motivate a Beta likelihood for PageRank and one-inflated Beta likelihoods for HHI and dependence, as specified in Section 5.1.2.

The chokepoint panel contains 1,440 observations for 24 nodes. Applying the same calendar-lag and consecutive-activity criteria excludes 200 currently inactive observations, 21 active observations without an available calendar lag, and 56 observations

preceded by inactivity. The resulting estimation sample contains 1,163 observations for all 24 chokepoints.

The chokepoint model includes five responses: PageRank, betweenness, share of monthly network flow, articulation-point status, and connection to an articulation point. The incident-global-bridge indicator is excluded because it differs from articulation-point status in only three observations and the correlation between the two binary indicators is 0.993.

Four predictors are included: lagged historical-neighbour activity share, log-transformed lagged mean voyage distance, and monthly sine and cosine terms. All four predictors are standardised within the chokepoint estimation sample.

PageRank and flow share lie strictly within $(0, 1)$, while betweenness has six exact zeros (0.516%) and otherwise takes values in $(0, 1)$. None of these responses equals one. Consequently, PageRank and flow share are modelled using Beta likelihoods, while betweenness uses a zero-inflated Beta likelihood.

### 5.1.2 Response likelihoods

The following response families are used for the baseline model and provide the likelihood framework for the proposed extensions. The expressions below use a scalar factor $f_{i,t}$ and its response-specific loading, as in Model 1; their latent linear predictors will require extension for the rank-two model. For bounded response $j$, the interior mean and precision are defined as

$$\operatorname{logit}(\mu_{i,t,j}) = \alpha_j + \lambda_j f_{i,t}, \tag{31}$$

$$A_{i,t,j} = \mu_{i,t,j}\phi_j, \qquad D_{i,t,j} = (1 - \mu_{i,t,j})\phi_j. \tag{32}$$

For terminals, $j = 1$ denotes PageRank, $j = 2$ HHI, and $j = 3$ dependence. For $j = 1$, the PageRank follows

$$Y_{i,t,1} \mid f_{i,t} \sim \text{Beta}(A_{i,t,1}, D_{i,t,1}). \tag{33}$$

For HHI and dependence ($j = 2, 3$), the boundary probability has a separate loading:

$$\begin{aligned} \operatorname{logit}(\pi_{i,t,j}) &= c_j^T + d_j^T f_{i,t}, \qquad j = 2, 3, \\ Y_{i,t,j} \mid f_{i,t} &\sim \pi_{i,t,j}\delta_1 + (1 - \pi_{i,t,j}) \operatorname{Beta}(A_{i,t,j}, D_{i,t,j}). \end{aligned} \tag{34}$$

where $\delta_1$ denotes a point mass at one. Interior observations contribute the Beta density multiplied by $1 - \pi_{i,t,j}$, while exact ones contribute by the probability $\pi_{i,t,j}$. Observed values are not clipped or displaced from the boundary. The overall conditional mean is

$$\mathbb{E}(Y_{i,t,j} \mid f_{i,t}) = \pi_{i,t,j} + (1 - \pi_{i,t,j})\mu_{i,t,j}. \tag{35}$$

Thus the sign of $\lambda_j$ alone does not determine how the overall mean changes with the factor.

For chokepoints, PageRank and flow share follow the interior Beta specification. For betweenness, let $\rho_{i,t,2}$ denote the probability of zero:

$$\begin{aligned} \operatorname{logit}(\rho_{i,t,2}) &= c_2^C + d_2^C f_{i,t}, \\ Y_{i,t,2} \mid f_{i,t} &\sim \rho_{i,t,2}\delta_0 + (1 - \rho_{i,t,2}) \operatorname{Beta}(A_{i,t,2}, D_{i,t,2}), \end{aligned} \tag{36}$$

where $\delta_0$ denotes a point mass at zero. The overall conditional mean is $(1 - \rho_{i,t,2})\mu_{i,t,2}$. The zero-probability loading is estimated separately from the interior-mean loading.

For binary response $k$,

$$B_{i,t,k} \mid f_{i,t} \sim \text{Bernoulli}(p_{i,t,k}), \tag{37}$$

$$\operatorname{logit}(p_{i,t,k}) = a_k + \ell_k f_{i,t}. \tag{38}$$

### 5.1.3 Prior specification

The following priors specify the shared response parameters and the rank-one baseline factor regression. Additional priors and identification constraints required by Models 2 and 3 will be stated with those extensions. The terminal and chokepoint models are estimated separately, with independent parameters.

The prior distributions were chosen to provide regularisation while allowing sufficient flexibility in the relationships between the latent factor and the observed responses. Let $q$ denote the number of covariates, with $q = 7$ for terminals and $q = 4$ for chokepoints. The factor-regression coefficients were assigned independent zero-centred Normal priors, $\beta_h \sim \mathcal{N}(0, 0.5^2)$ for $h = 1, \ldots, q$, favouring moderate covariate associations on the scale of the latent factor.

The loading associated with PageRank was constrained to be positive to fix the orientation of the latent factor, while the remaining loadings were allowed to take either sign. For the three bounded responses, the loading priors were $\lambda_1 \sim \text{HalfNormal}(1)$ and $\lambda_j \sim \mathcal{N}(0, 1)$ for $j = 2, 3$. The binary-response loadings were assigned independent priors $\ell_k \sim \mathcal{N}(0, 1)$ for $k = 1, \ldots, K$, where $K = 3$ for terminals and $K = 2$ for chokepoints.

For the interior components of the terminal bounded responses, intercept priors were specified on the logit scale, with centres chosen to reflect differences in the empirical scales of the corresponding network measures. In particular, PageRank typically takes values close to zero, making a zero-centred intercept prior, corresponding to an inverse-logit value of 0.5, which is inappropriate. Its intercept prior was therefore centred at $-4$, corresponding to an inverse-logit value of approximately 0.018. The intercept priors for HHI and dependence were instead centred at zero, allowing their interior means to vary around a reference value of 0.5 when the factor is zero. These choices correspond to $\alpha_j \sim \mathcal{N}(m_j, 1.5^2)$ for $j = 1, 2, 3$, with $(m_1, m_2, m_3) = (-4, 0, 0)$.

Binary-response intercepts were assigned independent $a_k \sim \mathcal{N}(0, 2^2)$ priors, allowing a relatively broad range of probabilities where the latent factor equals zero. For terminal HHI and dependence, the intercepts governing the probability of an exact one were assigned $c_j^T \sim \mathcal{N}(0, 1.5^2)$ priors, while their separate factor loadings followed $d_j^T \sim \mathcal{N}(0, 1)$, for $j = 2, 3$.

The precision parameters of the bounded continuous components were assigned independent log-Normal priors, with $\log \phi_j \sim \mathcal{N}(\log 10, 0.7^2)$ for $j = 1, 2, 3$. This specification ensures positive precision values, with prior median 10 and an approximate

central 95% interval of $[2.54, 39.44]$, allowing different degrees of dispersion around the conditional interior means.

For chokepoints, the same prior families and scales were adopted, with parameters estimated independently from the terminal model. The interior intercept centres were set to $(m_1, m_2, m_3) = (-4, -5, -4)$ for PageRank, betweenness, and flow share, respectively, reflecting the relatively small values typically observed for these measures. In particular, the inverse-logit of $-5$ is approximately 0.0067, comparable to the empirical median of chokepoint betweenness.

The betweenness zero-component intercept was assigned $c_2^C \sim \mathcal{N}(-5, 1.5^2)$, expressing a prior preference for a low probability of exact zeros at factor value zero. Its separate factor loading followed $d_2^C \sim \mathcal{N}(0, 1)$. Given the small number of observed zeros, inference on this component may be particularly sensitive to the prior specification.

All response-parameter priors were specified independently of one another and of the factor-regression coefficients. These distributions were adopted as initial regularising priors rather than independently elicited prior information. Their prior predictive implications and the sensitivity of the posterior findings to alternative specifications are examined in Section 6.4.

## 5.2 Model specifications

### 5.2.1 Baseline rank-one specification

For each node type separately, let $\boldsymbol{x}_{i,t}$ be the standardised covariate vector and $f_{i,t}$ be a scalar latent factor. The factor regression is

$$f_{i,t} \mid \boldsymbol{x}_{i,t}, \boldsymbol{\beta} \sim \mathcal{N}(\boldsymbol{x}_{i,t}^{\mathsf{T}}\boldsymbol{\beta}, 1). \tag{39}$$

The residual factor variance is fixed at one to define its scale. The PageRank loading is constrained to be positive to orient the factor, while the other loadings are unrestricted. Conditional on the factors and parameters, the responses are assumed

independent, and factors for different node-months are independent given their covariates and regression coefficients.

Let $\boldsymbol{\gamma}$ denote the vector of loadings for all response components:

$$\boldsymbol{\gamma} = \begin{cases} (\lambda_1, \lambda_2, \lambda_3, \ell_1, \ell_2, \ell_3, d_2^T, d_3^T)^\mathsf{T}, & \text{for terminals,} \\ (\lambda_1, \lambda_2, \lambda_3, \ell_1, \ell_2, d_2^C)^\mathsf{T}, & \text{for chokepoints.} \end{cases}$$

For component $m = 1, \ldots, M$, let $\eta_{i,t,m}$ denote the linear predictor on the logit scale: the logit of the interior Beta mean, the binary-response probability, or the boundary probability, according to the component. It can be written as $\eta_{i,t,m} = \xi_m + \gamma_m f_{i,t}$, where $\xi_m$ and $\gamma_m$ denote the corresponding intercept and loading, respectively, with $M = 8$ for terminals and $M = 6$ for chokepoints.

Substituting $f_{i,t} = \boldsymbol{x}_{i,t}^\mathsf{T}\boldsymbol{\beta} + z_{i,t}$, where $z_{i,t} \sim \mathcal{N}(0, 1)$ independently across node–month observations, gives

$$\eta_{i,t,m} = \xi_m + \boldsymbol{x}_{i,t}^\mathsf{T}(\gamma_m \boldsymbol{\beta}) + \gamma_m z_{i,t}. \tag{40}$$

Thus, the covariate coefficient vector for component $m$ is $\gamma_m\boldsymbol{\beta}$. Stacking these vectors as rows gives

$$\begin{pmatrix} \gamma_1\beta_1 & \cdots & \gamma_1\beta_q \\ \vdots & \ddots & \vdots \\ \gamma_M\beta_1 & \cdots & \gamma_M\beta_q \end{pmatrix} = \boldsymbol{\gamma}\boldsymbol{\beta}^\mathsf{T},$$

where $q$ denotes the number of covariates. Every row is a scalar multiple of $\boldsymbol{\beta}^\mathsf{T}$, so the matrix has rank at most one. This structure provides a parsimonious summary of how the covariates are associated with multiple network measures through a shared latent dimension, while allowing each response component to relate to that dimension with a different strength and sign.

Inference for this model is based on the joint posterior distribution of both the parameters and latent factors, which describes their uncertainty conditional on the observed responses and covariates. By using the Bayes' theorem, this distribution is proportional to the product of the response likelihood, the conditional distribution of the latent factors given the covariates and regression coefficients, and the parameter

prior densities (Gelman et al., 2013). These components are specified below, followed by the joint posterior distribution and the full conditional posterior kernels. Detailed derivations are provided in Appendix D.

Let $\mathcal{O}$ denote the set of node–month observations included in estimation. To simplify the notations, each observation $(i,t) \in \mathcal{O}$ is indexed by $n$, so that $f_n$, $\boldsymbol{x}_n$, $Y_{nj}$, and $B_{nk}$ correspond to $f_{i,t}$, $\boldsymbol{x}_{i,t}$, $Y_{i,t,j}$, and $B_{i,t,k}$, respectively.

Let $\mathbf{f} = (f_n)_{n\in\mathcal{O}}$ collect the latent factors. The observed responses are $Y_{nj}$ for the three bounded measures, $j = 1, 2, 3$, and $B_{nk}$ for the binary measures, $k = 1, \ldots, K$, where $K = 3$ for terminals and $K = 2$ for chokepoints.

Let $\boldsymbol{\theta}$ collect the response-specific parameters: the bounded-response intercepts $\alpha_j$, loadings $\lambda_j$, and precisions $\phi_j$; the binary-response intercepts $a_k$ and loadings $\ell_k$; and the boundary-component parameters $c_j$ and $d_j$. The latter denote $c_j^T, d_j^T$ for terminal HHI and dependence, $j = 2, 3$, or $c_2^C, d_2^C$ for chokepoint betweenness.

Under the conditional-independence assumption stated above, the joint likelihood contribution for a single node–month is the product of the individual response likelihood, denoted by $g_j(Y_{nj} \mid f_n, \boldsymbol{\theta})$, and the Bernoulli probabilities of the binary responses:

$$p(\mathbf{Y}_n, \mathbf{B}_n \mid f_n, \boldsymbol{\theta}) = \prod_{j=1}^{3} g_j(Y_{nj} \mid f_n, \boldsymbol{\theta}) \prod_{k=1}^{K} p_{nk}^{B_{nk}} (1 - p_{nk})^{1-B_{nk}}.$$

Combining the response contributions over all observations in $\mathcal{O}$ gives the likelihood

$$\begin{aligned} L(\boldsymbol{\theta}, \mathbf{f}) &= \prod_{n\in\mathcal{O}} p(\mathbf{Y}_n, \mathbf{B}_n \mid f_n, \boldsymbol{\theta}) \\ &= \prod_{n\in\mathcal{O}} \left[ \prod_{j=1}^{3} g_j(Y_{nj} \mid f_n, \boldsymbol{\theta}) \prod_{k=1}^{K} p_{nk}^{B_{nk}} (1 - p_{nk})^{1-B_{nk}} \right]. \end{aligned} \tag{41}$$

The factor regression in Equation (39) gives the joint conditional density of the latent factors:

$$p(\mathbf{f} \mid \boldsymbol{\beta}, \mathbf{X}) = \prod_{n\in\mathcal{O}} \mathcal{N}\big(f_n; \boldsymbol{x}_n^{\mathsf{T}}\boldsymbol{\beta}, 1\big), \tag{42}$$

where $\mathbf{X}$ is the covariate design matrix, with row $\boldsymbol{x}_n^{\mathsf{T}}$ for observation $n$.

Combining these components and denoting the observed responses and covariates

by $\mathcal{D}$, the joint posterior density is therefore proportional to

$$p(\boldsymbol{\theta}, \boldsymbol{\beta}, \mathbf{f} \mid \mathcal{D}) \propto L(\boldsymbol{\theta}, \mathbf{f})\, p(\mathbf{f} \mid \boldsymbol{\beta}, \mathbf{X})\, p(\boldsymbol{\theta}) p(\boldsymbol{\beta}). \tag{43}$$

where $p(\boldsymbol{\theta})$ and $p(\boldsymbol{\beta})$ are the products of the independent prior densities introduced in Section 5.1.3.

The full conditional posterior kernels are obtained by retaining only the terms in Equation (43) that depend on the parameter of interest, with all remaining unknown quantities held fixed. In the expressions below, *rest* denotes these remaining quantities together with the observed data and covariates.

Define $\mathcal{I}_j = \{n \in \mathcal{O} : 0 < Y_{nj} < 1\}$, so that for ordinary Beta responses, $\mathcal{I}_j = \mathcal{O}$, while for boundary-inflated responses it contains only the interior observations.

**Binary-response loadings.** For each binary-response loading $k = 1, \ldots, K$, the conditional posterior distribution is

$$p(\ell_k \mid \text{rest}) \propto \left[ \prod_{n \in \mathcal{O}} \frac{\exp\{B_{nk}(a_k + \ell_k f_n)\}}{1 + \exp(a_k + \ell_k f_n)} \right] \exp\left( -\frac{\ell_k^2}{2} \right). \tag{44}$$

**Bounded-response interior loadings.** For the unrestricted interior-mean loadings $\lambda_j$, $j = 2, 3$, corresponding to terminal HHI and dependence or chokepoint betweenness and flow share, the full conditional kernels are

$$p(\lambda_j \mid \text{rest}) \propto \left[ \prod_{n \in \mathcal{I}_j} \frac{Y_{nj}^{\mu_{nj}\phi_j - 1}(1 - Y_{nj})^{(1-\mu_{nj})\phi_j - 1}}{\Gamma(\mu_{nj}\phi_j)\Gamma((1-\mu_{nj})\phi_j)} \right] \exp\left( -\frac{\lambda_j^2}{2} \right), \qquad j = 2, 3, \tag{45}$$

For PageRank, the half-Normal prior restricts the loading to positive values:

$$p(\lambda_1 \mid \text{rest}) \propto \left[ \prod_{n \in \mathcal{O}} \frac{Y_{n1}^{\mu_{n1}\phi_1 - 1}(1 - Y_{n1})^{(1-\mu_{n1})\phi_1 - 1}}{\Gamma(\mu_{n1}\phi_1)\Gamma((1-\mu_{n1})\phi_1)} \right] \exp\left( -\frac{\lambda_1^2}{2} \right) \mathbb{I}(\lambda_1 > 0). \tag{46}$$

**Boundary-probability loadings.** For terminal HHI and dependence, define $Z_{nj} = \mathbb{1}_{\{Y_{nj}=1\}}$, $j = 2, 3$. The full conditional kernels are

$$p(d_j^T \mid \text{rest}) \propto \left[\prod_{n\in\mathcal{O}} \frac{\exp\{Z_{nj}(c_j^T + d_j^T f_n)\}}{1+\exp(c_j^T + d_j^T f_n)}\right] \exp\left[-\frac{(d_j^T)^2}{2}\right], \qquad j = 2, 3. \tag{47}$$

For chokepoint betweenness, let $Z_{n2} = \mathbb{1}_{\{Y_{n2}=0\}}$ The full conditional kernel is

$$p(d_2^C \mid \text{rest}) \propto \left[\prod_{n\in\mathcal{O}} \frac{\exp\{Z_{n2}(c_2^C + d_2^C f_n)\}}{1+\exp(c_2^C + d_2^C f_n)}\right] \exp\left[-\frac{(d_2^C)^2}{2}\right]. \tag{48}$$

**Factor-regression coefficients.** The Gaussian factor regression and the independent $\mathcal{N}(0, 0.5^2)$ coefficient priors give the full conditional kernel

$$p(\boldsymbol{\beta} \mid \text{rest}) \propto \exp\left\{ -\frac{1}{2}\Big[\boldsymbol{\beta}^\mathsf{T}(\mathbf{X}^\mathsf{T}\mathbf{X} + 4\mathbf{I}_q)\boldsymbol{\beta} - 2\boldsymbol{\beta}^\mathsf{T}\mathbf{X}^\mathsf{T}\mathbf{f}\Big]\right\}. \tag{49}$$

Recognising the kernel of a multivariate Normal distribution gives the conjugate full conditional

$$\boldsymbol{\beta} \mid \text{rest} \sim \mathcal{N}_q\Big((\mathbf{X}^\mathsf{T}\mathbf{X} + 4\mathbf{I}_q)^{-1}\mathbf{X}^\mathsf{T}\mathbf{f}, (\mathbf{X}^\mathsf{T}\mathbf{X} + 4\mathbf{I}_q)^{-1}\Big). \tag{50}$$

Matrix notation is used here for compactness, with $\mathbf{I}_q$ denoting the $q \times q$ identity matrix.

**Latent factors.** For each node–month observation, the full conditional of the factor combines the Gaussian regression density with the likelihood contributions of all observed responses:

$$p(f_n \mid \text{rest}) \propto L_n(f_n) \exp\left[-\frac{1}{2}(f_n - \boldsymbol{x}_n^\mathsf{T}\boldsymbol{\beta})^2\right]. \tag{51}$$

For terminals, the observation-specific likelihood is

$$\begin{aligned} L_n^T(f_n) = p(Y_{n1} \mid f_n, \alpha_1, \lambda_1, \phi_1) \\ \times \prod_{j=2}^{3} p(Y_{nj} \mid f_n, \alpha_j, \lambda_j, \phi_j, c_j^T, d_j^T) \\ \times \prod_{k=1}^{3} p(B_{nk} \mid f_n, a_k, \ell_k). \end{aligned}$$

For chokepoints, it is

$$\begin{aligned} L_n^C(f_n) = p(Y_{n1} \mid f_n, \alpha_1, \lambda_1, \phi_1) \\ \times p(Y_{n2} \mid f_n, \alpha_2, \lambda_2, \phi_2, c_2^C, d_2^C) \\ \times p(Y_{n3} \mid f_n, \alpha_3, \lambda_3, \phi_3) \\ \times \prod_{k=1}^{2} p(B_{nk} \mid f_n, a_k, \ell_k). \end{aligned}$$

The inflated-response terms in these expressions denote the complete mixture contributions, including the boundary probabilities and, for interior observations, the Beta densities.

**Binary-response intercepts.** Under the prior $a_k \sim \mathcal{N}(0, 2^2)$, the posterior is

$$p(a_k \mid \text{rest}) \propto \exp\left\{ -\frac{a_k^2}{8} + a_k \sum_{n \in \mathcal{O}} B_{nk} - \sum_{n \in \mathcal{O}} \log\left[1 + \exp(a_k + \ell_k f_n)\right] \right\}. \tag{52}$$

**Bounded-response interior intercepts.** For $\alpha_j \sim \mathcal{N}(m_j, 1.5^2)$, the full conditional kernels are

$$p(\alpha_j \mid \text{rest}) \propto \left[ \prod_{n \in \mathcal{I}_j} \frac{Y_{nj}^{\mu_{nj}\phi_j - 1}(1 - Y_{nj})^{(1-\mu_{nj})\phi_j - 1}}{\Gamma(\mu_{nj}\phi_j)\Gamma((1-\mu_{nj})\phi_j)} \right] \exp\left[ -\frac{(\alpha_j - m_j)^2}{2(1.5)^2} \right], \qquad j = 1, 2, 3. \tag{53}$$

**Boundary-probability intercepts.** For terminal HHI and dependence, $c_j^T \sim \mathcal{N}(0, 1.5^2)$ gives

$$p(c_j^T \mid \text{rest}) \propto \left[\prod_{n\in\mathcal{O}} \frac{\exp\{Z_{nj}(c_j^T + d_j^T f_n)\}}{1+\exp(c_j^T + d_j^T f_n)}\right] \exp\left[-\frac{(c_j^T)^2}{2(1.5)^2}\right], \qquad j = 2, 3. \tag{54}$$

For chokepoint betweenness, $c_2^C \sim \mathcal{N}(-5, 1.5^2)$ gives

$$p(c_2^C \mid \text{rest}) \propto \exp\left\{ -\frac{(c_2^C+5)^2}{2(1.5)^2} + c_2^C \sum_{n\in\mathcal{O}} Z_{n2} - \sum_{n\in\mathcal{O}} \log\left[1+\exp(c_2^C + d_2^C f_n)\right] \right\}. \tag{55}$$

**Beta precision parameters.** Under the prior assumption $\log\phi_j \sim \mathcal{N}(\log 10, 0.7^2)$, the full conditional posterior for each bounded-response component $j = 1, 2, 3$ is

$$p(\phi_j \mid \text{rest}) \propto \left[\prod_{n\in\mathcal{I}_j} \frac{\Gamma(\phi_j) Y_{nj}^{\mu_{nj}\phi_j - 1}(1-Y_{nj})^{(1-\mu_{nj})\phi_j - 1}}{\Gamma(\mu_{nj}\phi_j)\Gamma((1-\mu_{nj})\phi_j)}\right] \frac{1}{\phi_j} \exp\left[-\frac{(\log\phi_j - \log 10)^2}{2(0.7)^2}\right], \tag{56}$$

with $\phi_j > 0$.

Except for the Normal full conditional distribution in Equation (50), the full conditionals above do not generally have standard conjugate forms. The joint posterior density is known up to a normalising constant, but the integrals required for normalisation and marginalisation are analytically intractable. Inference is therefore based on numerical sampling using the No-U-Turn Sampler (NUTS), whose implementation and associated diagnostic criteria are described in Section 5.3.

### 5.2.2 Random-intercept rank-one specification

The baseline rank-one specification assumes that the conditional mean of the latent factor is entirely determined by the observed covariates. However, nodes may exhibit persistent differences in their network characteristics that are not fully captured by these predictors. To account for such unobserved heterogeneity, the model is extended by introducing a node-specific random intercept into the latent-factor regression.

Let $i = 1, \ldots, N$ index the distinct nodes included in the estimation sample. The extended factor regression is

$$f_{i,t} \mid \boldsymbol{x}_{i,t}, \boldsymbol{\beta}, \mu_i \sim \mathcal{N}\left(\boldsymbol{x}_{i,t}^{\mathsf{T}}\boldsymbol{\beta} + \mu_i, 1\right), \tag{57}$$

where $\mu_i$ represents a persistent node-specific deviation from the conditional factor mean implied by the observed covariates. Equivalently,

$$f_{i,t} = \boldsymbol{x}_{i,t}^{\mathsf{T}}\boldsymbol{\beta} + \mu_i + z_{i,t}, \qquad z_{i,t} \sim \mathcal{N}(0,1). \tag{58}$$

The effect $\mu_i$ is common to all observations associated with node $i$, while $z_{i,t}$ represents observation-specific residual variation. The residual variance remains fixed at one, as in the baseline specification.

The node-specific intercepts are assigned a hierarchical Gaussian prior,

$$\mu_i \mid \sigma_\mu \overset{\text{iid}}{\sim} \mathcal{N}(0, \sigma_\mu^2), \qquad i = 1, \ldots, N, \tag{59}$$

with

$$\sigma_\mu \sim \text{HalfNormal}(1). \tag{60}$$

The parameter $\sigma_\mu$ controls the magnitude of persistent between-node heterogeneity: small values imply stronger shrinkage of the node-specific effects towards zero, making the model close to the baseline specification, whereas larger values allow greater differences across nodes.

The response-specific likelihoods, prior distributions of the baseline parameters, and positive PageRank loading constraint remain unchanged. The extension therefore preserves the rank-one structure of the covariate coefficient matrix $\boldsymbol{\gamma}\boldsymbol{\beta}^{\mathsf{T}}$, while introducing persistent node-specific contributions $\gamma_m \mu_i$ to each response component's linear predictor.

Using the observation index $n \in \mathcal{O}$ introduced in the baseline specification, let $i(n)$ denote the node associated with observation $n$. Define $\boldsymbol{\mu} = (\mu_1, \ldots, \mu_N)^{\mathsf{T}}$. The conditional density of the latent factors is

$$p(\mathbf{f} \mid \boldsymbol{\beta}, \boldsymbol{\mu}, \mathbf{X}) = \prod_{n \in \mathcal{O}} \mathcal{N}\left(f_n; \boldsymbol{x}_n^\top \boldsymbol{\beta} + \mu_{i(n)}, 1\right). \tag{61}$$

Combining this density with the response likelihood and the hierarchical prior gives the joint posterior

$$p(\boldsymbol{\theta}, \boldsymbol{\beta}, \mathbf{f}, \boldsymbol{\mu}, \sigma_\mu \mid \mathcal{D}) \propto L(\boldsymbol{\theta}, \mathbf{f})\, p(\mathbf{f} \mid \boldsymbol{\beta}, \boldsymbol{\mu}, \mathbf{X}) p(\boldsymbol{\theta}) p(\boldsymbol{\beta}) p(\boldsymbol{\mu} \mid \sigma_\mu) p(\sigma_\mu), \tag{62}$$

where $L(\boldsymbol{\theta}, \mathbf{f})$ is the response likelihood defined in Equation (41), $p(\boldsymbol{\mu} \mid \sigma_\mu) = \prod_{i=1}^{N} p(\mu_i \mid \sigma_\mu)$, and $p(\sigma_\mu)$ is the Half-Normal prior as in Equation (60).

The full conditional posterior kernels follow from Equation (62). Since the response likelihood and its parameters remain unchanged, the corresponding full conditional kernels retain the forms derived for the baseline model, conditional on the latent factors. Only the conditional distributions involving the factor regression and the newly introduced hierarchical parameters require modification.

**Factor-regression coefficients.** Define the $|\mathcal{O}|$-dimensional vector $\boldsymbol{\mu}_{\mathcal{O}}$ whose $n^{th}$ element is $\mu_{i(n)}$. Combining the Gaussian factor regression with the independent $\mathcal{N}(0, 0.5^2)$ coefficient priors gives the following posterior density

$$p(\boldsymbol{\beta} \mid \text{rest}) \propto \exp\left\{ -\frac{1}{2}\Big[\boldsymbol{\beta}^\top (\mathbf{X}^\top \mathbf{X} + 4\mathbf{I}_q)\boldsymbol{\beta} - 2\boldsymbol{\beta}^\top \mathbf{X}^\top (\mathbf{f} - \boldsymbol{\mu}_{\mathcal{O}})\Big] \right\}, \tag{63}$$

which is the multivariate Normal kernel defined as

$$\boldsymbol{\beta} \mid \text{rest} \sim \mathcal{N}_q\Big((\mathbf{X}^\top \mathbf{X} + 4\mathbf{I}_q)^{-1} \mathbf{X}^\top (\mathbf{f} - \boldsymbol{\mu}_{\mathcal{O}}), (\mathbf{X}^\top \mathbf{X} + 4\mathbf{I}_q)^{-1}\Big). \tag{64}$$

**Latent factors.** For each node–month observation, the full conditional kernel combines the unchanged response likelihood with the modified Gaussian factor regression:

$$p(f_n \mid \text{rest}) \propto L_n(f_n) \exp\left[ -\frac{1}{2}\left(f_n - \boldsymbol{x}_n^\top \boldsymbol{\beta} - \mu_{i(n)}\right)^2 \right], \tag{65}$$

where $L_n(f_n)$ denotes the same observation-specific likelihood defined for terminals and chokepoints in the baseline specification.

**Node-specific intercepts.** For each node $i$, let $\mathcal{O}_i = \{n \in \mathcal{O} : i(n) = i\}$ denote the observations associated with node $i$ in the estimation sample, and let $n_i = |\mathcal{O}_i|$. Combining the Gaussian prior for $\mu_i$ with the factor regression gives

$$p(\mu_i \mid \text{rest}) \propto \exp\left(-\frac{\mu_i^2}{2\sigma_\mu^2}\right) \exp\left[-\frac{1}{2}\sum_{n\in\mathcal{O}_i}\left(f_n - \boldsymbol{x}_n^\top\boldsymbol{\beta} - \mu_i\right)^2\right]. \tag{66}$$

Completing the square yields the conjugate Normal full conditional

$$\mu_i \mid \text{rest} \sim \mathcal{N}(m_{\mu_i}, V_{\mu_i}), \tag{67}$$

where

$$V_{\mu_i} = \left(n_i + \frac{1}{\sigma_\mu^2}\right)^{-1}, \qquad m_{\mu_i} = V_{\mu_i}\sum_{n\in\mathcal{O}_i}\left(f_n - \boldsymbol{x}_n^\top\boldsymbol{\beta}\right).$$

**Node-effect scale parameter.** The full conditional kernel for $\sigma_\mu$ combines its Half-Normal prior with the Gaussian hierarchical distribution of the node-specific intercepts:

$$p(\sigma_\mu \mid \text{rest}) \propto \mathbb{I}(\sigma_\mu > 0)\,\sigma_\mu^{-N} \exp\left[-\frac{1}{2}\left(\sigma_\mu^2 + \frac{\sum_{i=1}^N \mu_i^2}{\sigma_\mu^2}\right)\right]. \tag{68}$$

Unlike the conditional distribution of the node-specific intercepts, this kernel does not correspond to a standard conjugate distribution for $\sigma_\mu$.

For posterior computation, the hierarchical model is implemented through an equivalent non-centred parametrisation,

$$u_i \overset{\text{iid}}{\sim} \mathcal{N}(0,1), \qquad \mu_i = \sigma_\mu u_i. \tag{69}$$

This transformation preserves the prior $\mu_i \mid \sigma_\mu \sim \mathcal{N}(0, \sigma_\mu^2)$ and therefore leaves the posterior distribution of the model parameters and latent factors unchanged. It only

changes the coordinates used for sampling to improve computational efficiency (Papaspiliopoulos et al., 2007).

### 5.2.3 Rank-two specification

The rank-two specification extends the baseline model by introducing a second latent factor, allowing the dependence structure among the observed responses to be represented in a two-dimensional latent space. This extension is motivated by the limitations of the rank-one specification, particularly its restricted ability to reproduce certain patterns of association among network criticality indicators.

For each node $i$ and month $t$, the scalar latent factor is replaced by a two-dimensional vector

$$\mathbf{f}_{i,t} = \begin{pmatrix} f_{i,t,1} \\ f_{i,t,2} \end{pmatrix}.$$

The conditional distribution of the latent factors is specified as

$$\mathbf{f}_{i,t} \mid \mathbf{x}_{i,t}, \mathbf{B} \sim \mathcal{N}_2\left(\mathbf{B}^\top \mathbf{x}_{i,t}, \mathbf{I}_2\right), \tag{70}$$

where $\mathbf{B} \in \mathbb{R}^{q\times 2}$ is the matrix of factor-regression coefficients and $\mathbf{I}_2$ denotes the two-dimensional identity matrix. Each element of $\mathbf{B}$ is assigned the same Gaussian prior used for the regression coefficients in the baseline model, $B_{kh} \sim \mathcal{N}(0, 0.5^2), h = 1, 2$. The two latent dimensions therefore have independent unit-variance Gaussian residuals, conditional on the observed covariates.

The observation models retain the likelihood families introduced in Section 5.2.1. The main modification concerns the linear predictors, in which the scalar product between a response-specific loading and the latent factor is replaced by the inner product between two-dimensional vectors.

For each continuous response $j$, the conditional mean of the interior Beta component is given by

$$\text{logit}(\mu_{i,t,j}) = \alpha_j + \boldsymbol{\lambda}_j^\top \mathbf{f}_{i,t},$$

where $\boldsymbol{\lambda}_j \in \mathbb{R}^2$ contains the two factor loadings associated with response $j$. Similarly, for each binary response $k$,

$$\text{logit}(p_{i,t,k}) = a_k + \boldsymbol{\ell}_k^\top \mathbf{f}_{i,t},$$

where $\boldsymbol{\ell}_k \in \mathbb{R}^2$ denotes the corresponding loading vector.

The probabilities associated with the boundary components of the inflated Beta distributions are extended analogously, with a separate two-dimensional loading vector for each boundary response.

The prior distributions of the intercepts and precision parameters remain unchanged, while the loading priors are extended independently to the two latent dimensions. As in the baseline model, the first PageRank loading is constrained to be positive through a half-normal prior, $\lambda_{\text{PR},1} \sim \text{halfNormal}(1)$, whereas the second PageRank loading follows an unconstrained Gaussian prior $\lambda_{\text{PR},2} \sim \mathcal{N}(0, 1)$. All remaining loadings follow the corresponding Gaussian priors specified for the rank-one model.

The joint distribution of the rank-two model retains the conditional-independence structure of the baseline rank-one specification. Let $\mathbf{F} = \{\mathbf{f}_n\}_{n=1}^N$ denote the collection of two-dimensional latent factors, and let $\boldsymbol{\theta}$ contain the factor-regression coefficients, response-specific loadings, intercepts, and precision parameters. The joint distribution can be expressed as

$$p(\mathbf{Y}, \mathbf{F}, \boldsymbol{\theta} \mid \mathbf{X}) = p(\boldsymbol{\theta}) \prod_{n=1}^{N} \left[ \mathcal{N}_2\left(\mathbf{f}_n; \mathbf{B}^\top \mathbf{x}_n, \mathbf{I}_2\right) \prod_{j=1}^{J} p(Y_{nj} \mid \mathbf{f}_n, \boldsymbol{\theta}) \right]. \tag{71}$$

Here, $p(\boldsymbol{\theta})$ represents the joint prior distribution of the model parameters, while the response likelihoods retain the distributions defined for the corresponding terminal and chokepoint models in Section 5.2.1.

The rank-two specification is estimated separately for terminals and chokepoints,

using the same response variables, covariates, and estimation samples as the corresponding rank-one models. No node-specific random intercepts are included in this extension, allowing the effect of increasing the latent rank to be examined separately from the introduction of persistent node heterogeneity.

## 5.3 Posterior computation

### 5.3.1 NUTS sampling algorithm and implementation

The models were implemented in Python 3.12.15 using PyMC 5.20.1 (Abril-Pla et al., 2023), with PyTensor 2.27.1 as the computational backend. Posterior inference was performed using the No-U-Turn Sampler (NUTS) (Hoffman and Gelman, 2014), an adaptive variant of Hamiltonian Monte Carlo (HMC) (Neal, 2011). Unlike traditional random-walk MCMC methods, HMC exploits gradient information to construct proposals that can explore high-dimensional posterior distributions more efficiently. NUTS extends HMC by automatically determining the trajectory length, avoiding the need to manually specify the number of integration steps. The principles underlying HMC and its NUTS extension are outlined below.

Let $\boldsymbol{\omega} = (\boldsymbol{\theta}, \boldsymbol{\beta}, \mathbf{f})$ be the vector containing all unknown response parameters, regression coefficients, and latent factors. The sampler operates on an unconstrained representation of these parameters $\mathbf{q} = T(\boldsymbol{\omega})$. By the change-of-variables formula, the transformed log-posterior density is

$$\mathcal{L}(\mathbf{q}) = \log p_{\boldsymbol{\omega}}\big(T^{-1}(\mathbf{q}) \mid \mathcal{D}\big) + \log \left| \det \frac{\partial T^{-1}(\mathbf{q})}{\partial \mathbf{q}} \right|. \tag{72}$$

Following Neal (2011), an auxiliary momentum vector $\mathbf{v} \sim \mathcal{N}(\mathbf{0}, \mathbf{M})$ is introduced, where $\mathbf{M}$ is a positive-definite mass matrix. Intuitively, HMC treats the current parameter values $\mathbf{q}$ as the position of a moving particle. The negative transformed log-posterior defines a potential-energy landscape, with lower values in regions of higher transformed posterior density. The momentum allows the particle to move through this landscape, while the gradient of $\mathcal{L}(\mathbf{q})$ guides its trajectory.

The negative log-density of the Gaussian momentum, up to an additive constant, defines the kinetic energy $\frac{1}{2}\mathbf{v}^\mathsf{T}\mathbf{M}^{-1}\mathbf{v}$. Together with the potential energy $-\mathcal{L}(\mathbf{q})$, this defines the Hamiltonian:

$$H(\mathbf{q}, \mathbf{v}) = -\mathcal{L}(\mathbf{q}) + \frac{1}{2}\mathbf{v}^\mathsf{T}\mathbf{M}^{-1}\mathbf{v}. \tag{73}$$

The corresponding equations of motion are

$$\frac{d\mathbf{q}}{d\tau} = \nabla_{\mathbf{v}} H(\mathbf{q}, \mathbf{v}) = \mathbf{M}^{-1}\mathbf{v}, \qquad \frac{d\mathbf{v}}{d\tau} = -\nabla_{\mathbf{q}} H(\mathbf{q}, \mathbf{v}) = \nabla_{\mathbf{q}}\mathcal{L}(\mathbf{q}),$$

with $\tau$ denoting an artificial integration time.

These equations are approximated numerically using leapfrog integration: for a step size $\epsilon$, each integration step comprises the updates

$$\begin{aligned} \mathbf{v}_{1/2} &= \mathbf{v} + \frac{\epsilon}{2}\nabla_{\mathbf{q}}\mathcal{L}(\mathbf{q}), \\ \mathbf{q}' &= \mathbf{q} + \epsilon\mathbf{M}^{-1}\mathbf{v}_{1/2}, \\ \mathbf{v}' &= \mathbf{v}_{1/2} + \frac{\epsilon}{2}\nabla_{\mathbf{q}}\mathcal{L}(\mathbf{q}'). \end{aligned} \tag{74}$$

These updates act on the complete state vector, so regression coefficients, response parameters, and latent factors are explored jointly rather than through sequential parameter-specific full conditional updates.

NUTS constructs trajectories by recursively expanding a tree of leapfrog states in either the forward or backward direction until the no-U-turn criterion detects that the trajectory is doubling back, a numerical stopping criterion is met, or the maximum tree depth is reached. A state from the constructed trajectory is then selected according to a rule that preserves the target distribution (Hoffman and Gelman, 2014).

Algorithm 1 summarises the sampling procedure for the rank-one baseline model.

**Algorithm 1** NUTS sampling for rank-one baseline model

**Require:** Data $\mathcal{D}$; chains $C$; warm-up iterations $W$; retained draws $S$; target acceptance $\delta$.

**for** each chain $c = 1, \ldots, C$ **do**
  Initialise the unconstrained parameter vector $\mathbf{q}$.
  Initialise step size $\epsilon$ and diagonal mass matrix $\mathbf{M}$.
  **for** $m = 1, \ldots, W + S$ **do**
    Sample momentum $\mathbf{v} \sim \mathcal{N}(\mathbf{0}, \mathbf{M})$.
    Initialise the trajectory at $(\mathbf{q}, \mathbf{v})$.
    **repeat**
      Sample a direction $d \sim \text{Uniform}(\{-1, +1\})$.
      Extend the tree in direction $d$ by doubling the number of leapfrog steps.
      Update the candidate state according to the NUTS selection rule.
    **until** a stopping criterion is met
    Set $\mathbf{q}$ to the selected candidate position.
    **if** $m \leq W$ **then**
      Adapt $\epsilon$ to approach target acceptance $\delta$.
      Adapt $\mathbf{M}$ to account for differences in parameter scales.
    **else**
      Store the joint posterior draw $\boldsymbol{\omega}^{(c,m-W)} = T^{-1}(\mathbf{q})$.
      Compute factor means $\mathbf{x}_n^\top \boldsymbol{\beta}$ and residuals $f_n - \mathbf{x}_n^\top \boldsymbol{\beta}$ for all $n \in \mathcal{O}$.
    **end if**
  **end for**
**end for**
**return** $S$ posterior draws per chain.

Each node-type model was estimated using four parallel chains, with 20,000 burn-in iterations and 30,000 retained draws per chain. The use of multiple chains allows convergence to be assessed by comparing within-chain and between-chain variability (Gelman et al., 2013), while the number of iterations was increased to allow more extensive sampler adaptation during warm-up and improve the Monte Carlo precision of posterior summaries through a larger number of retained draws.

The target acceptance probability was set to 0.80, corresponding to the target acceptance rate used during step-size adaptation in the warm-up phase (Hoffman and Gelman, 2014). The random seed was set to 20261008.

### 5.3.2 Sampling diagnostics

Posterior summaries and convergence diagnostics were computed using ArviZ 0.21.0, with NumPy 1.26.4, pandas 2.3.3, and Matplotlib 3.9.2 used for data processing and visualisation.

The assessment covered regression coefficients, response parameters, latent factors, factor means, and factor residuals. Several complementary diagnostics were considered together to examine agreement across chains, sampling efficiency, within-chain stationarity, and the Monte Carlo precision of posterior estimates.

Agreement across chains was assessed using rank-normalised $\widehat{R}$, with values close to one indicating consistency between within-chain and between-chain variability. Bulk and tail effective sample sizes (ESS) were used to assess how much information the correlated draws provided about the centre and tails of the posterior distribution, respectively (Vehtari et al., 2021).

Post-burn-in divergences were also monitored, as they signal numerical difficulties in following Hamiltonian trajectories and may indicate that some posterior regions have not been adequately explored (Betancourt, 2017).

The Geweke and Heidelberger–Welch diagnostics provided additional checks on individual chains (Geweke, 1992; Heidelberger and Welch, 1983).

The Geweke diagnostic compares the means of an early and a late segment of a chain, standardising their difference using a standard error that accounts for autocorrelation. Under suitable stationarity conditions, the statistic has an asymptotic standard Normal distribution, with large absolute values suggesting inconsistency between the two segments.

The Heidelberger–Welch procedure first assesses stationarity using a test based on the Cramér–von Mises statistic. If necessary, the test is repeated after removing progressively larger initial portions of the chain, up to half of the available draws. Once stationarity is accepted, a half-width test checks whether the Monte Carlo confidence interval for the posterior mean, computed from the remaining draws, is sufficiently

narrow relative to the magnitude of the estimated mean.

Trace plots and autocorrelation functions (ACFs) for selected parameters complemented these numerical checks. Trace plots were inspected for persistent trends, differences between chains, and whether the chains explored similar posterior regions. ACFs showed how dependence between draws changed with increasing lag, with faster decay generally indicating more efficient sampling (Vehtari et al., 2021).

# Chapter 6

# Results and discussion

## 6.1 Rank-one model results

This section presents the posterior results for the terminal model and the chokepoint model, under the rank-one factor specification. Covariate associations, relationships among responses, and the interpretation of the latent factor in each case are discussed. Furthermore, posterior predictive checks are performed for each type of node to assess how well the model reproduces the observed responses distributions and associations.

### 6.1.1 Terminal posterior analysis

The rank-one terminal model was estimated using four MCMC chains, each with 20,000 warm-up iterations and 30,000 retained draws. The posterior estimates of the factor-regression coefficients, computed as the mean of the retained MCMC samples, are presented in Table 9, together with their 95% highest-density intervals (HDIs).

The most pronounced positive associations are observed for lagged exporter status and terminal capacity. Conditional on the other predictors, being classified as an exporter in the preceding month is associated with an increase of approximately 1.4 units in the expected factor value. This is substantial relative to the residual standard deviation of the factor, which is fixed at one.

Neighbour activity and voyage distance in the previous month are also positively related to the factor, but their estimated associations are considerably weaker. By contrast, the posterior intervals for terminal age and the seasonal terms include zero,

indicating considerable uncertainty about their individual contributions.

Table 9: Factor-regression coefficients for the terminal model. Numeric predictors are standardised; exporter status is a zero–one indicator.

| Predictor | Posterior mean | 95% HDI |
|---|---|---|
| Lagged neighbour activity share | 0.096 | $[0.058, 0.134]$ |
| $\log(1 + \text{lagged voyage distance})$ | 0.062 | $[0.022, 0.102]$ |
| Month sine | $-0.032$ | $[-0.068, 0.003]$ |
| Month cosine | $-0.003$ | $[-0.038, 0.033]$ |
| $\log(1 + \text{capacity})$ | 0.409 | $[0.367, 0.452]$ |
| Terminal age | $-0.019$ | $[-0.057, 0.019]$ |
| Lagged exporter indicator | 1.395 | $[1.303, 1.489]$ |

The response loadings in Table 10 provide further insight into the interpretation of the terminal factor by describing associations with the interior means of bounded responses, the probabilities of binary outcomes, and the boundary probabilities.

The clearest pattern combines increasing PageRank with decreasing counterparty concentration. In particular, higher factor values correspond to both a lower interior mean for HHI and a reduced probability that HHI equals one.

The structural indicators show contrasting relationships: the probability of being an articulation point increases with the factor, while the probability of being incident to a global bridge decreases. For the indicator of adjency to an articulation point, the posterior interval includes zero, providing no clear evidence of a directional association.

Table 10: Loadings on the latent factor. Bounded interior means and all probabilities use logit links.

| Response component | Posterior mean | 95% HDI |
|---|---|---|
| PageRank interior mean | 0.359 | $[0.348, 0.370]$ |
| HHI interior mean | $-0.535$ | $[-0.567, -0.503]$ |
| Dependence interior mean | 0.420 | $[0.395, 0.445]$ |
| Articulation-point probability | 0.770 | $[0.684, 0.858]$ |
| Incident-global-bridge probability | $-0.638$ | $[-0.713, -0.563]$ |
| Connected-to-articulation probability | 0.109 | $[-0.014, 0.235]$ |
| Probability of HHI equal to one | $-3.985$ | $[-4.699, -3.303]$ |
| Probability of dependence equal to one | $-0.545$ | $[-0.636, -0.451]$ |

The magnitude of the HHI relationship becomes clearer when expressed in the

original response units. As the factor increases from 0 to 1, the posterior median of the expected HHI decreases from approximately 0.66 to 0.38, with a median difference of $-0.289$ (95% equal-tailed credible interval: $[-0.311, -0.266]$), incorporating both the interior Beta component and the point mass at one.

Dependence exhibits a more complex pattern. Its interior mean increases with the factor, whereas the probability of an exact value of one declines. Combining these components, as defined in Equation (35), yields a mildly U-shaped conditional expectation over the displayed factor range.

Figure 6 displays the fitted response functions, allowing the individual response components and their combined effects to be examined across the estimated factor range.




Figure 6: Fitted response functions against the terminal latent factor. Lines represent posterior medians, and shaded bands indicate equal-tailed 95% credible intervals for conditional expected responses or probabilities. HHI and dependence means include the point mass at one. The horizontal axis spans the full empirical range of posterior mean latent factor scores in the estimation sample.

To relate these posterior patterns to observed network positions, two terminal-month observations whose posterior mean latent factor values are closest to the 5th

and 95th percentiles of the estimation sample are considered. Mugardos LNG Terminal, located in Galicia, Spain, has an estimated latent factor score of approximately $-1.0$ in October 2022, compared with 2.73 for Sabine Pass LNG Terminal, located in Cameron Parish, Louisiana, in June 2024. Their observed PageRank values are 0.011 and 0.039, respectively, while their counterparty HHI values are 1.00 and 0.17. Mugardos is classified as an importer and has an incident global bridge, whereas Sabine Pass is an exporter without one. These observations were selected to illustrate contrasting positions along the latent dimension while avoiding the most extreme cases, and display differences which are broadly consistent with the estimated relationships between the latent factor and the terminal responses presented above, although the two observations also differ in commercial role.

An additional case of interest is Rasgas LNG Terminal 3, Qatar, which exhibits the highest posterior mean latent factor score in the estimation sample, approximately 6.38 in June 2020. Its posterior mean factor score can be decomposed into a systematic component $\mathbf{x}_n^\top \boldsymbol{\beta} \approx 1.84$ and a residual component $z_n \approx 4.55$, indicating a substantial deviation from the position explained by the observed covariates. Importantly, this deviation is not confined to a single observation: across its 59 terminal-month observations, Rasgas exhibits a positive posterior mean residual in every month, with an average of approximately 3.13. Similar, although less pronounced, patterns are observed for other terminals, including Fos Tonkin, Yamal and Sabine Pass, whose residual components remain predominantly positive over time. Conversely, terminals such as Cameroon FLNG and Sakhalin II exhibit consistently negative posterior mean residuals. Table E.1 in the Appendix provides descriptive summaries of these patterns.

To complement the observation-level examples, Figure 7 illustrates the geographical distribution of posterior mean latent factor scores averaged over time for each terminal in the estimation sample. The highest average factor scores are observed among major LNG export terminals located in geographically distinct regions, including the Middle East, North America, Russia and West Africa. This pattern is consistent with the strong positive association between lagged exporter status and the

latent factor identified in Table 9. Conversely, terminals with the lowest average scores are also distributed across different geographical areas, with several located in East and Southeast Asia. Overall, the spatial distribution does not suggest that high or low factor values are confined to a single geographical region. Rather, the geographical representation reinforces the interpretation of the latent factor as a network-related dimension associated with terminal characteristics and commercial roles, rather than geographical location alone.




Figure 7: Geographical distribution of time-averaged posterior mean latent factor scores for LNG terminals under the baseline rank-one model. The colour scale is centred at zero.

The persistent residual patterns described above suggest that part of the variation left unexplained by the baseline covariates may reflect systematic differences across terminals rather than exclusively observation-specific fluctuations. This provides an empirical motivation for extending the rank-one model by introducing node-specific effects, allowing persistent terminal-level heterogeneity to be represented separately from residual variation. The results of this extension are presented in Section 6.2.

While the preceding examples illustrate the interpretation of the latent factor and highlight potential limitations of the baseline specification, the extent to which the model reproduces the observed response patterns is examined through the posterior predictive checks in the following subsection.

**Terminal posterior predictive assessment**

This section evaluates how well the fitted model reproduces the observed response distributions and associations by comparing statistics computed from the terminal-month observations with those obtained from replicated responses conditional on the posterior factors. These comparisons are conducted both overall and within lagged-role groups, with 95% equal-tailed intervals summarising the variability of the statistics across 200 conditional in-sample replications.

The model closely reproduces the HHI overall response means and the overall frequencies of dependence equal to one, with the observed overall binary proportions also lying within their respective replicated intervals. Nevertheless, discrepancies emerge in the dispersion, quantiles and role-specific boundary frequencies of the bounded responses. For PageRank, the model overestimates dispersion among lagged importers and underestimates it among lagged exporters. This difference is also reflected in the tails of the distributions: among importers, the observed 5th percentile (0.0089) exceeds the replicated 95% interval of $[0.0062, 0.0068]$, indicating that the model generates excessively low values, whereas among exporters, the observed 95th percentile (0.0714) exceeds the corresponding replicated interval of $[0.0587, 0.0648]$, indicating an underestimation in the upper tail. For HHI, the model underestimates both the frequency of values equal to one (14.86%, included in a 95% interval of $[13.48\%, 16.23\%]$, compared with an observed 18.00%) and the overall dispersion among exporters, where the observed mean 0.3103 exceeds its replicated interval $[0.2818, 0.2993]$. For dependence, the discrepancies differ between terminal roles. The observed frequency of dependence equal to one among importers (26.45%) slightly exceeds the replicated interval of $[21.72\%, 26.07\%]$, whereas the corresponding frequency among exporters (9.04%) falls below the replicated interval of $[11.79\%, 16.14\%]$. Moreover, the model underestimates dependence dispersion among importers and overestimates it among exporters. The observed standard deviations are 0.3574 and 0.2296, respectively, compared with replicated intervals of $[0.3256, 0.3402]$ and $[0.2709, 0.2882]$. The complete marginal posterior predictive checks are reported in Table E.2 in the Appendix.

These findings indicate that, although the baseline model captures several features of the overall response distributions, it does not fully reproduce their role-specific dispersion, tail behaviour and boundary frequencies.

Association between responses is assessed by comparing observed pairwise Spearman correlations with those obtained from 200 replicated datasets selected evenly across the four chains. Table 11 focuses on the three response pairs associated with the five largest absolute discrepancies between observed correlations and posterior predictive medians, reporting results for the full sample, importers, and exporters to allow systematic comparisons across terminal roles. The complete set of 45 pairwise correlation checks is provided in Table E.3 in the Appendix.

Table 11: Selected joint posterior predictive checks. Entries are Spearman correlations. Replicated intervals use 200 conditional in-sample replicates.

| Response pair | Lagged role | Observed | Rep. median | 95% interval |
|---|---|---|---|---|
| AP–GB | All | 0.290 | -0.124 | $[-0.162, -0.095]$ |
| AP–GB | Importer | 0.137 | -0.037 | $[-0.082, 0.003]$ |
| AP–GB | Exporter | 0.546 | -0.158 | $[-0.209, -0.102]$ |
| HHI–GB | All | 0.533 | 0.295 | $[0.248, 0.335]$ |
| HHI–GB | Importer | 0.644 | 0.189 | $[0.137, 0.226]$ |
| HHI–GB | Exporter | 0.398 | 0.336 | $[0.268, 0.397]$ |
| DEP–GB | All | -0.066 | -0.070 | $[-0.113, -0.037]$ |
| DEP–GB | Importers | -0.154 | -0.020 | $[-0.062, 0.016]$ |
| DEP–GB | Exporters | 0.195 | -0.118 | $[-0.186, -0.065]$ |

The largest discrepancy concerns the association between articulation-point and incident-global-bridge status among exporters, for which the observed correlation is positive (0.546), whereas the posterior predictive median is negative ($-0.158$). The same sign reversal is observed for the full sample, while the replicated correlation among importers is closer to zero, although still negative. This indicates that the model fails to reproduce the positive association between these two indicators across terminal roles. The association between counterparty HHI and incident-global-bridge status also exhibits substantial discrepancies, particularly among importers, for whom the observed correlation (0.644) considerably exceeds the posterior predictive median (0.189). In contrast, the corresponding discrepancy among exporters is relatively small.

For dependence and incident-global-bridge status, the model reproduces the observed negative association reasonably well in the full sample but underestimates its magnitude among importers. Among exporters, the observed correlation is positive (0.195), whereas the posterior predictive median is negative ($-0.118$), indicating a further sign reversal.

These discrepancies suggest limitations in the rank-one model's ability to reproduce the dependence structure among responses. In particular, the single latent factor has opposite loadings on articulation-point and incident-global-bridge probabilities, while the assumption of conditional independence does not allow for additional residual associations between these responses. Consequently, the model struggles to reproduce their positive observed co-occurrence.

### 6.1.2 Chokepoint posterior analysis

This section examines how the latent factor relates to network characteristics and measures of centrality, flow exposure, and structural importance for the chokepoint specification. The adequacy of the fitted model is subsequently evaluated through posterior predictive checks.

For chokepoints, the factor-regression results in Table 12 indicate relatively weak associations with the available predictors. Greater neighbour activity and longer average voyage distances in the preceding month are associated with higher factor values; both posterior intervals exclude zero, although the lower bound for voyage distance is close to zero, suggesting greater uncertainty about this association. The seasonal coefficients provide limited evidence of seasonal variation in the conditional factor mean: the interval for the negative cosine coefficient excludes zero, whereas the interval for the sine coefficient includes zero.

With respect to the responses, Table 13 shows positive loadings across the interior means and binary structural probabilities, with all reported HDIs excluding zero. These relationships are further illustrated by the fitted response curves in Figure 8.

Table 12: Factor-regression coefficients for the chokepoint model. Numeric predictors are standardised. Intervals are 94% highest-density intervals (HDIs), as exported by ArviZ.

| Predictor | Posterior mean | 94% HDI |
|---|---|---|
| Lagged neighbour activity share | 0.103 | $[0.042, 0.162]$ |
| $\log(1 + \text{lagged voyage distance})$ | 0.065 | $[0.004, 0.123]$ |
| Month sine | $-0.036$ | $[-0.096, 0.021]$ |
| Month cosine | $-0.065$ | $[-0.123, -0.007]$ |

The factor is positively associated with PageRank, betweenness, and flow share, indicating that higher scores correspond to chokepoints occupying more central positions and carrying larger proportions of the network's LNG flows. Both articulation-point status and connection to an articulation point become more likely as the factor increases, unlike the terminal specification, where the latter association is not clearly established. The negative loading on the probability that betweenness equals zero is consistent with this pattern, as higher factor values are associated with a lower probability of zero betweenness, further reinforcing the positive relationship between the factor and its overall conditional expectation.

Table 13: Loadings on the latent factor for the chokepoint model. Bounded interior means and all probabilities use logit links. Intervals are 94% HDIs, as exported by ArviZ.

| Response component | Posterior mean | 94% HDI |
|---|---|---|
| PageRank interior mean | 0.263 | $[0.244, 0.281]$ |
| Betweenness interior mean | 1.024 | $[0.971, 1.079]$ |
| Flow-share interior mean | 0.619 | $[0.580, 0.658]$ |
| Articulation-point probability | 0.719 | $[0.566, 0.876]$ |
| Connected-to-articulation probability | 1.206 | $[0.945, 1.461]$ |
| Probability of betweenness equal to zero | $-1.658$ | $[-2.536, -0.797]$ |

To illustrate how these relationships are reflected in individual observations, consider Cape of Good Hope in May 2021 and Taiwan Strait in October 2021. Their posterior mean factor values, located near the 5th and 95th percentiles of the estimation sample, are approximately $-1.63$ and 1.55, respectively. Taiwan Strait has substantially higher observed PageRank (0.035 versus 0.007), betweenness (0.040 versus




Figure 8: Fitted response functions against the chokepoint latent factor. Lines are posterior medians and bands are equal-tailed 95% credible intervals for conditional expected responses or binary probabilities. The betweenness mean includes the mass at zero. The horizontal axis spans the full empirical range of posterior mean latent factor scores in the estimation sample.

0.0006), and flow share (3.6% versus 0.3%). Nevertheless, both chokepoint-month observations are classified as articulation points and are connected to articulation points. Their different factor values therefore reflect variation in the continuous measures even though these binary structural indicators coincide, illustrating how the estimated factor summarises different aspects of chokepoint centrality, flow exposure, and structural position.

Although the chokepoint latent factor does not exhibit values as extreme as those observed for some terminals, a similar pattern of persistent residual deviations emerges. The highest and lowest posterior mean factor scores occur for Gibraltar Strait in November 2024 (2.394) and Sunda Strait in December 2022 ($-2.282$), respectively. In both cases, the systematic components explained by the covariates are close to zero ($-0.014$ and 0.044), whereas the posterior mean residual components account for nearly the entire factor scores (2.408 and $-2.326$, respectively).

As observed for the terminals, these deviations are not limited to isolated months. Gibraltar Strait exhibits positive posterior mean residuals in 96.6% of its 59 observations, with an average of 1.294, whereas Sunda Strait exhibits negative residuals in 95.9% of its 49 observations, averaging $-1.179$. Similar patterns are found across other chokepoints: among the 20 with at least 20 observations, 13 exhibit residuals of the same sign in more than 85% of their observed months. Descriptive summaries of these cases are provided in Appendix E.1.2, Table E.4.

The five highest and five lowest time-averaged scores are highlighted in Figure 9 to illustrate the spatial persistent variations in the estimated latent dimension. The highest average factor scores are associated with Gibraltar Strait, Taiwan Strait, Suez Canal, Malacca Strait, and Bab el-Mandeb Strait, all located along major maritime corridors connecting Europe, the Middle East, and Asia. Conversely, the lowest average scores are observed for Mona Passage and Windward Passage in the Caribbean, Magellan Strait at the southern tip of South America, and Torres Strait and Sunda Strait in the Indo-Pacific region. Notably, the coexistence of relatively high and low factor scores among geographically proximate chokepoints in Southeast Asia suggests

that geographical proximity does not necessarily correspond to similar positions along the estimated latent dimension.




Figure 9: Geographical distribution of time-averaged posterior mean latent factor scores for chokepoints under the baseline rank-one model. The colour scale is centred at zero.

The persistent residual patterns identified above reinforce the evidence of node-level heterogeneity already observed for terminals, suggesting that the baseline covariates may not fully account for systematic differences across chokepoints and providing further motivation for the node-specific extension discussed in Section 6.2.

### Chokepoint posterior predictive assessment

Following the same conditional, in-sample approach used for terminals, the chokepoint posterior predictive checks indicate that the model reproduces overall response means and binary proportions reasonably closely.

The observed PageRank mean (0.01764) is close to the replicated mean (0.01767), and its observed 5th, median, and 95th percentiles fall within their respective replicated intervals. The model also reproduces the low proportion of observations with zero betweenness reasonably well (0.516% observed versus 0.584% replicated). Nevertheless, the observed fifth percentile of betweenness (0.000575) lies just below the corresponding replicated interval $[0.000582, 0.000988]$; its observed median (0.007677)

is also slightly below the replicated interval $[0.007731, 0.008545]$.

The largest marginal discrepancies concern flow share. The observed standard deviation (0.02010) exceeds the 95% replicated interval $[0.01693, 0.01933]$, indicating that the model underestimates the dispersion of this response. The observed median (0.01589) falls below its replicated interval $[0.01796, 0.02040]$, whereas the observed 95th percentile (0.06426) exceeds the corresponding interval $[0.05522, 0.06304]$. These discrepancies indicate that, despite reproducing the overall mean (0.02341 observed versus 0.02354 replicated), the model does not fully capture the distribution of flow share.

Posterior predictive checks also examine whether the model reproduces the observed associations between responses. Pairwise Spearman correlations are compared with those obtained from 200 replicated datasets selected evenly across the four chains. As shown in Table 14, the model reproduces the PageRank–betweenness association reasonably closely but underestimates associations involving flow share, as well as the PageRank–articulation association. The observed PageRank–articulation correlation lies slightly above the upper endpoint of the replicated interval.

Table 14: Selected joint predictive checks for chokepoints. Entries are Spearman correlations; intervals use 200 conditional in-sample replicates.

| Response pair | Observed | Rep. median | 95% interval |
|---|---|---|---|
| PageRank–betweenness | 0.685 | 0.655 | $[0.608, 0.699]$ |
| PageRank–flow share | 0.672 | 0.531 | $[0.481, 0.569]$ |
| Betweenness–flow share | 0.800 | 0.718 | $[0.683, 0.746]$ |
| PageRank–articulation | 0.287 | 0.204 | $[0.137, 0.272]$ |

### 6.1.3 Sampling diagnostics

Sampling performance was evaluated using rank-normalised $\widehat{R}$, bulk and tail effective sample sizes (ESS), post-burn-in divergences, and the Geweke and Heidelberger–Welch diagnostics. The Geweke and Heidelberger–Welch diagnostics were applied separately

to each chain for every global parameter in both node-specific models, using the retained posterior draws. Observation-level latent factors were excluded from these additional tests to limit the computational burden associated with evaluating thousands of observation-specific quantities across four chains. Their convergence and sampling efficiency were nevertheless assessed through rank-normalised $\widehat{R}$ and bulk and tail effective sample sizes.

Neither model presented post-burn-in divergences. The maximum rank-normalised $\widehat{R}$, rounded to three decimal places, was 1.001 for terminals and 1.003 for chokepoints.

For terminals, the minimum bulk ESS was 2,931, associated with the lagged exporter coefficient, and the minimum tail ESS was 6,726 for the HHI boundary loading. For chokepoints, the corresponding values were 1,693 and 3,698, both associated with the betweenness precision parameter, indicating satisfactory convergence and sampling efficiency.

Trace plots and within-chain autocorrelation functions are presented in Appendix F.2 and provide additional visual evidence of sampling behaviour. In particular, the betweenness loading and precision parameters exhibit relatively persistent autocorrelation in the chokepoint model, consistent with their comparatively lower effective sample sizes.

The Geweke diagnostic was used to assess differences between the early and late portions of each chain. For the terminal model, 92 out of 104 parameter-chain tests (88.5%) satisfied $|Z| \leq 1.96$. The remaining 12 tests exceeded the threshold, with a maximum absolute $Z$-score of 3.07.

For the chokepoint model, 71 out of 76 tests (93.4%) satisfied the same criterion. The five exceedances were concentrated primarily among the precision parameters of the continuous responses, with one additional exceedance involving the betweenness boundary-component loading. The maximum absolute $Z$-score was 2.46.

Although these results indicate some differences between the early and late portions of individual chains, the tests were conducted separately for multiple parameters

and chains without adjustment for multiple comparisons. Consequently, isolated exceedances should not be interpreted as evidence of non-convergence, particularly in the presence of satisfactory $\widehat{R}$, ESS, and divergence diagnostics.

The Heidelberger–Welch procedure was used to assess stationarity and the Monte Carlo precision of posterior mean estimates. All 104 terminal and 76 chokepoint parameter-chain tests satisfied the stationarity criterion. For terminals, 93 tests passed without excluding any additional draws, while 11 required the exclusion of between 1,000 and 3,000 initial draws. For chokepoints, 74 tests passed without additional exclusions, while the remaining two required the exclusion of 3,000 and 4,000 draws, respectively.

The half-width test assessed whether the half-width of the 95% Monte Carlo confidence interval for the posterior mean was below 10% of its absolute estimated value. All 76 chokepoint parameter-chain tests satisfied this criterion. For terminals, 96 out of 104 tests (92.3%) passed, while the remaining eight failures involved only two parameters: the seasonal regression coefficient associated with month_cos and the continuous intercept for counterparty HHI, each across all four chains. Notably, both parameters had posterior means close to zero, making the relative half-width criterion particularly sensitive to small absolute differences. These failures therefore do not necessarily indicate substantial Monte Carlo uncertainty in absolute terms.

Table 15 summarises the convergence and mixing diagnostics, together with the results of the Geweke and Heidelberger–Welch tests for both baseline models, while additional results are reported in Appendix F.2.

Overall, the combined diagnostics provide satisfactory sampling behaviour for both baseline models. The absence of divergences, $\widehat{R}$ values close to one, and sufficiently large effective sample sizes provide complementary evidence of satisfactory posterior exploration and sampling efficiency.

Table 15: Summary of sampling diagnostics for the baseline terminal and chokepoint models. Geweke and Heidelberger–Welch results refer to parameter-chain tests on distinct global parameters.

| Diagnostic | Terminals | Chokepoints |
|---|---|---|
| *MCMC convergence and mixing* | | |
| Post-warm-up divergences | 0 | 0 |
| Maximum $\widehat{R}$ | 1.001 | 1.003 |
| Minimum bulk ESS | 2,931 | 1,693 |
| Minimum tail ESS | 6,726 | 3,698 |
| *Geweke diagnostic* | | |
| Total tests | 104 | 76 |
| Passed | 92 (88.5%) | 71 (93.4%) |
| Failed | 12 (11.5%) | 5 (6.6%) |
| Maximum $\lvert Z \rvert$ | 3.07 | 2.46 |
| *Heidelberger–Welch diagnostic* | | |
| Stationarity passed | 104 (100%) | 76 (100%) |
| Half-width passed | 96 (92.3%) | 76 (100%) |
| Half-width failed | 8 (7.7%) | 0 (0%) |

## 6.2 Node-specific intercept model results

This section presents the posterior results of the rank-one model extended with node-specific random intercepts. The analysis focuses on the magnitude of between-node heterogeneity, changes in the factor-regression coefficients, and the estimated node-specific effects. The response loadings are also examined to assess whether the interpretation of the latent factor changes under the extended specification. Finally, posterior predictive checks and sampling diagnostics are reported.

Both models were estimated using four MCMC chains, each with 20,000 warm-up iterations and 30,000 retained draws. The results for terminals and chokepoints are presented separately, as in the baseline analysis.

### 6.2.1 Terminal posterior analysis

The posterior distribution of the random-effect scale parameter provides evidence of substantial persistent heterogeneity across terminals. The posterior mean of $\sigma_\mu$ is 1.373, with a 94% highest-density interval (HDI) of $[1.193, 1.565]$. Since the residual

variance of the latent factor remains fixed at one, this estimate indicates that persistent between-terminal differences represent an important component of the variation in latent factor scores, beyond that explained by the observed covariates.

The introduction of node-specific intercepts also changes the estimated associations between the covariates and the latent factor. Table 16 reports the posterior estimates alongside those obtained under the baseline specification.

Table 16: Factor-regression coefficients for the terminal random-intercept model, compared with the baseline rank-one specification. Numeric predictors are standardised; exporter status is a zero–one indicator.

| Predictor | Baseline mean | Node mean | Node 94% HDI |
|---|---|---|---|
| Lagged neighbour activity | 0.096 | 0.016 | $[-0.039, 0.072]$ |
| $\log(1 + \text{lagged voyage distance})$ | 0.062 | 0.053 | $[-0.022, 0.129]$ |
| Month sine | $-0.032$ | $-0.038$ | $[-0.080, 0.004]$ |
| Month cosine | $-0.003$ | 0.004 | $[-0.038, 0.045]$ |
| $\log(1 + \text{capacity})$ | 0.409 | 0.725 | $[0.493, 0.957]$ |
| Terminal age | $-0.019$ | $-0.326$ | $[-0.558, -0.089]$ |
| Lagged exporter indicator | 1.395 | 1.309 | $[0.823, 1.783]$ |

Compared with the baseline specification, the coefficient associated with lagged neighbour activity decreases from 0.096 to 0.016, and its posterior interval now includes zero. The coefficient for lagged voyage distance remains positive but is less precisely estimated, with its interval also including zero. These changes suggest that some of the associations identified under the baseline model may have reflected persistent differences between terminals that are now represented by the random intercepts. In contrast, terminal capacity and lagged exporter status remain positively associated with the latent factor even after accounting for unobserved terminal-specific heterogeneity. However, their posterior intervals are considerably wider than under the baseline specification, indicating greater uncertainty in separating covariate associations from persistent node-specific differences. A further change concerns terminal age: whereas its posterior interval included zero under the baseline specification, the random-intercept model yields a negative posterior mean of $-0.326$, with a 94% HDI of $[-0.558, -0.089]$, indicating a negative conditional association between terminal age

and the latent factor.

The response loadings reported in Table 17 preserve the main directional relationships identified under the baseline specification, although the magnitudes of several estimates are smaller.

Table 17: Posterior estimates of response loadings for the terminal random-intercept model. Bounded interior means and probabilities use logit links. Intervals are 94% HDIs.

| Response component | Posterior mean | 94% HDI |
|---|---|---|
| PageRank interior mean | 0.197 | $[0.189, 0.206]$ |
| HHI interior mean | $-0.283$ | $[-0.301, -0.266]$ |
| Dependence interior mean | 0.232 | $[0.217, 0.248]$ |
| Articulation-point probability | 0.394 | $[0.347, 0.441]$ |
| Incident-global-bridge probability | $-0.311$ | $[-0.351, -0.271]$ |
| Connected-to-articulation probability | 0.096 | $[0.021, 0.172]$ |
| Probability of HHI equal to one | $-1.872$ | $[-2.141, -1.618]$ |
| Probability of dependence equal to one | $-0.318$ | $[-0.378, -0.262]$ |

Higher factor values remain associated with higher PageRank, lower counterparty concentration, and a greater probability of being an articulation point. The probability of having an incident global bridge continues to decrease as the factor increases. As in the baseline model, dependence exhibits opposing relationships between its interior and boundary components: the interior mean increases with the factor, whereas the probability of dependence equal to one decreases. One difference concerns connection to an articulation point: its loading remains positive and its 94% HDI now excludes zero, whereas the corresponding interval included zero under the baseline specification.

To illustrate persistent terminal-level heterogeneity, Figure 10 illustrates the posterior distributions of the node-specific intercepts for the five terminals with the highest and the five with the lowest posterior mean effects, restricting the selection to terminals with at least 20 observations in the estimation sample. Findings are generally consistent with the persistent residual patterns identified under the baseline model: the largest positive posterior mean effect is associated with Rasgas LNG Terminal 3, approximately 7.615, with a 95% equal-tailed credible interval of $[6.942, 8.320]$. Other terminals exhibiting substantial positive effects include Fos Tonkin LNG Terminal

(3.69), Yamal LNG Terminal (3.18), Sabine Pass LNG Terminal (3.11), and Pluto LNG Terminal (3.10), all of which also exhibited predominantly positive latent-factor residuals under the baseline specification. Their positive random intercepts therefore suggest that the systematic deviations observed under the baseline specification can, at least partly, be represented by persistent node-level differences. At the opposite end, Sakhalin II LNG Terminal and Cameroon FLNG Terminal exhibit the most negative effects among the selected terminals, consistent with the predominantly negative residuals identified in the baseline analysis. Oman Qalhat LNG Terminal, Punta Europa LNG Terminal, and Elba Island LNG Terminal also display negative node-specific intercepts, indicating that their latent factor values tend to be lower than those predicted by the observed covariates alone.




Figure 10: Posterior distributions of selected terminal-specific random intercepts under the rank-one model. The five largest and five smallest posterior mean effects are shown among terminals with at least 20 observations in the estimation sample. Points represent posterior means, and horizontal bars indicate 95% equal-tailed credible intervals. The dashed vertical line denotes zero.

### Posterior predictive assessment

Posterior predictive checks were conducted to examine whether the introduction of node-specific intercepts improves the reproduction of marginal response distributions and associations between responses. The same conditional, in-sample approach adopted for the baseline model was used: observed response statistics were compared with those obtained from 200 replicated datasets generated conditional on posterior latent factor values. Table 18 shows that the random-intercept model closely reproduces the overall means of the observed terminal responses.

Table 18: Marginal posterior predictive checks for the terminal random-intercept model. Observed and replicated means are reported on the original response scale and rounded to five decimal places to highlight differences. For binary responses, means correspond to proportions.

| Response | Observed mean | Replicated mean |
|---|---|---|
| PageRank | 0.01852 | 0.01854 |
| HHI | 0.58872 | 0.58933 |
| Dependence | 0.49478 | 0.49618 |
| Articulation point (AP) | 0.10571 | 0.10637 |
| Incident global bridge (GB) | 0.58500 | 0.58509 |

However, as in the baseline model, agreement in overall means does not necessarily imply adequate reproduction of dispersion, tail behaviour, or associations between responses. In particular, the joint posterior predictive checks reveal that several discrepancies identified under the baseline specification persist. Table 19 compares selected response pairs exhibiting these discrepancies under the baseline and random-intercept models, allowing an assessment of whether introducing node-specific intercepts improves the model's ability to reproduce the observed associations.

Table 19: Selected joint posterior predictive checks for terminals. Entries are Spearman correlations.

| Response pair | Model | Observed | Rep. median | 95% interval |
|---|---|---|---|---|
| AP–GB | Baseline | 0.290 | $-0.124$ | $[-0.162, -0.095]$ |
| AP–GB | Node intercept | 0.290 | $-0.108$ | $[-0.143, -0.070]$ |
| HHI–GB | Baseline | 0.533 | 0.295 | $[0.248, 0.335]$ |
| HHI–GB | Node intercept | 0.533 | 0.242 | $[0.204, 0.289]$ |

The most pronounced discrepancy continues to concern the association between articulation-point and incident-global-bridge status, which the model still fails to reproduce. The discrepancy is particularly pronounced among exporters. Their observed AP–GB correlation is 0.546, compared with a replicated median of $-0.150$ under the random-intercept model. The association between HHI and incident-global-bridge status also remains underestimated. For the full terminal sample, the observed correlation is 0.533, whereas the replicated median decreases from 0.295 under the baseline specification to 0.242 under the random-intercept model.

Thus, introducing node-specific intercepts does not uniformly improve the reproduction of joint response associations. This finding is consistent with the structure of the extension. The random intercepts account for persistent heterogeneity in the common latent factor, but the model retains a single latent dimension and conditional independence of the responses given that factor. Consequently, the additional node-specific variation does not necessarily provide the flexibility required to reproduce associations that are poorly represented by the rank-one structure.

Overall, the random-intercept specification provides a more explicit representation of persistent between-terminal heterogeneity, while preserving the main directional relationships between the latent factor and the observed responses. Nevertheless, the posterior predictive results indicate that important limitations of the baseline rank-one specification remain, particularly in reproducing the joint occurrence of articulation-point and incident-global-bridge status.

The complete marginal and joint posterior predictive results are reported in the Appendix E.2.

### 6.2.2 Chokepoint posterior analysis

The random-intercept extension also identifies substantial persistent heterogeneity across chokepoints. The posterior mean of the random-effect scale parameter is 2.377, with a 94% HDI of $[1.806, 2.973]$. As for terminals, the estimated random-effect scale suggests that persistent differences between chokepoints contribute substantially to

variation in the latent factor beyond that associated with the observed covariates. The posterior estimates of the factor-regression coefficients are reported in Table 20 and suggest that the introduction of node-specific effects changes the relative contributions of the observed covariates.

Table 20: Factor-regression coefficients for the chokepoint random-intercept model, compared with the baseline rank-one specification. Numeric predictors are standardised. Intervals for the random-intercept model are 94% HDIs.

| Predictor | Baseline mean | Node mean | Node 94% HDI |
|---|---|---|---|
| Lagged neighbour activity | 0.103 | 0.316 | $[0.229, 0.401]$ |
| $\log(1 + \text{lagged voyage distance})$ | 0.065 | 0.088 | $[-0.067, 0.244]$ |
| Month sine | $-0.036$ | $-0.107$ | $[-0.175, -0.040]$ |
| Month cosine | $-0.065$ | $-0.085$ | $[-0.152, -0.018]$ |

The coefficient associated with lagged neighbour activity increases from 0.103 under the baseline model to 0.316 under the random-intercept specification, with a posterior interval excluding zero. The coefficient for lagged voyage distance also remains positive, but its posterior interval includes zero, indicating greater uncertainty about its conditional association with the latent factor. Both seasonal coefficients are negative and their reported HDIs exclude zero, whereas, under the baseline specification, only the cosine coefficient had an interval excluding zero. Given the sinusoidal encoding of calendar months, the combination of these negative coefficients implies a cyclical pattern in the expected latent factor, with lower values around February and higher values around August.

Turning to the response loadings, Table 21 shows that the estimated relationships between the latent factor and the response loadings retain the directions observed under the baseline model. Higher latent factor values correspond to higher interior means for PageRank, betweenness, and flow share, as well as higher probabilities of being an articulation point and being connected to an articulation point. The negative loading on the zero-betweenness component indicates that higher factor values are associated with a lower probability that betweenness equals zero. Thus, the estimated factor continues to represent a dimension associated with chokepoint centrality, flow

Table 21: Posterior estimates of response loadings for the chokepoint random-intercept model. Bounded interior means and probabilities use logit links. Intervals are 94% HDIs.

| Response component | Posterior mean | 94% HDI |
|---|---|---|
| PageRank interior mean | 0.107 | $[0.097, 0.117]$ |
| Betweenness interior mean | 0.303 | $[0.280, 0.327]$ |
| Flow-share interior mean | 0.382 | $[0.350, 0.414]$ |
| Articulation-point probability | 0.362 | $[0.289, 0.435]$ |
| Connected-to-articulation probability | 0.354 | $[0.253, 0.457]$ |
| Probability of betweenness equal to zero | $-1.048$ | $[-1.576, -0.532]$ |

exposure, and structural importance.

The node-specific intercepts provide further evidence of persistent heterogeneity across chokepoints. Figure 11 presents their posterior distributions for all 24 chokepoints, ordered by posterior mean. The estimates reveal considerable heterogeneity across maritime passages, with both positive and negative effects relative to the covariate-dependent component of the latent factor. The Strait of Hormuz exhibits the largest positive node-specific effect, with a posterior mean of approximately 4.79. Other substantial positive effects are associated with Malacca Strait (3.51), Gibraltar Strait (3.50), Taiwan Strait (3.13), Dover Strait (2.85), Suez Canal (2.73), and Bab el-Mandeb Strait (2.54). Indeed, several of these passages occupy strategically important positions along major international shipping corridors. Conversely, Torres Strait exhibits the most negative effect (approximately $-4.07$), followed by Windward Passage ($-3.75$), Magellan Strait ($-3.73$), Mona Passage ($-3.17$), and Sunda Strait ($-2.25$). These negative intercepts indicate systematically lower latent factor values relative to the covariate-based predictions, a pattern which is broadly consistent with the lower factor scores observed for several of these passages in the baseline specification.

Although the posterior summaries indicate substantial heterogeneity across chokepoints, the convergence diagnostics discussed in Section 6.2.3 raise concerns about the reliability of some estimates. The corresponding interpretations should therefore be treated with caution.




Figure 11: Posterior distributions of the node-specific random intercepts for all 24 chokepoints under the rank-one model. Points represent posterior means, and horizontal bars indicate 95% equal-tailed credible intervals. The dashed vertical line denotes zero. Estimates are preliminary due to unresolved sampling diagnostics.

**Posterior predictive assessment**

To assess whether the random-intercept specification adequately captures the observed characteristics of the chokepoint network, posterior predictive checks were conducted using 200 conditional in-sample replicates. As for the preceeding models, the assessment considers both the marginal distributions of individual responses and the associations between them.

At the marginal level, the model closely reproduces the overall response means. The observed PageRank mean is 0.01764, compared with a replicated mean of 0.01765. For betweenness, the corresponding values are 0.01321 and 0.01323, while for flow share they are 0.02341 and 0.02342.

Nevertheless, discrepancies remain in the associations between responses. Table 22 reports four illustrative response pairs involving different combinations of centrality, flow exposure, and structural importance measures, for which the observed Spearman correlations exceed the upper bounds of the 95% posterior predictive intervals, highlighting the model's tendency to underestimate certain positive associations.

Table 22: Selected joint posterior predictive checks for chokepoints under the random-intercept model. Entries are Spearman correlations. Replicated intervals are based on 200 conditional in-sample replicates.

| Response pair | Observed | Rep. median | 95% interval |
|---|---|---|---|
| PageRank–betweenness | 0.685 | 0.473 | $[0.417, 0.516]$ |
| PageRank–flow share | 0.672 | 0.609 | $[0.562, 0.644]$ |
| Betweenness–flow share | 0.800 | 0.700 | $[0.659, 0.739]$ |
| PageRank–articulation | 0.287 | 0.201 | $[0.140, 0.271]$ |

The largest discrepancy among the selected associations concerns PageRank and betweenness, for which the replicated median correlation is 0.473, compared with an observed value of 0.685. This represents a deterioration relative to the baseline specification, which yielded a replicated median of approximately 0.655. The model also underestimates the positive associations between PageRank and flow share, betweenness and flow share, and PageRank and articulation-point status, indicating that introducing node-specific intercepts does not fully resolve the limitations of the baseline

rank-one model in reproducing the observed dependence structure among chokepoint responses. The complete marginal and joint posterior predictive results are reported in Appendix E.2.2.

### 6.2.3 Sampling diagnostics

Sampling performance was assessed using the same diagnostic procedures as for the baseline specification. Table 23 summarises the main results.

Table 23: Sampling diagnostics for the terminal and chokepoint random-intercept models. Geweke and Heidelberger–Welch results refer to parameter-chain tests for global parameters.

| Diagnostic | Terminals | Chokepoints |
|---|---|---|
| Post-burn-in divergences | 0 | 183 |
| Maximum $\widehat{R}$ | 1.003* | 1.040 |
| | | betweenness intercept |
| Minimum bulk ESS | 795 | 112 |
| | PageRank intercept | flow-share intercept |
| Minimum tail ESS | 1,734 | 202 |
| | PageRank intercept | betweenness intercept |
| Passed Geweke | 115 (95.8%) | 85 (92.4%) |
| Failed Geweke | 5 (4.2%) | 7 (7.6%) |
| Passed stationarity | 120 (100%) | 92 (100%) |
| Passed half-width | 113 (94.2%) | 92 (100%) |

* The maximum reported $\widehat{R}$ for terminals is shared by the PageRank interior-mean intercept, the HHI interior-mean intercept, and the HHI boundary-component intercept.

The terminal model exhibits satisfactory overall sampling behaviour: no post-burn-in divergences were recorded, and the maximum rank-normalised $\widehat{R}$ was 1.003. The minimum bulk and tail ESS were 795 and 1,734, respectively, and although these values are lower than those obtained under the baseline specification, they remain compatible with adequate posterior exploration for the reported summaries. The Geweke diagnostic was satisfied in 115 of the 120 parameter-chain tests, all tests passed the Heidelberger–Welch stationarity criterion, and the half-width test was passed by 113 of the 120 tests.

Conversely, the chokepoint model exhibited less satisfactory sampling diagnostics. Although 85 of the 92 parameter-chain tests satisfied the Geweke criterion, and all

tests passed the Heidelberger–Welch stationarity and half-width criteria, a total of 183 post-warm-up divergences were recorded, with 168 concentrated in one chain and 15 in another. The maximum rank-normalised $\widehat{R}$ was 1.040, while the minimum bulk and tail ESS were 112 and 202, respectively. These results suggest incomplete exploration of the posterior distribution, potentially associated with difficult posterior geometry and insufficient mixing. In particular, the presence of divergences, together with the low ESS values, indicates that the available posterior draws may not adequately represent the target distribution. Consequently, the chokepoint posterior summaries should be interpreted cautiously, as the reported credible intervals and posterior predictive results may be affected by incomplete posterior exploration.

Additional trace plots, autocorrelation functions, and detailed diagnostic results are provided in Appendix F.3.

## 6.3 Rank-two model results

The rank-two specifications were estimated to investigate whether introducing a second latent dimension could capture patterns of network criticality that were not adequately represented by the baseline rank-one and random-intercept models. The posterior results, however, provide limited evidence of a clearly identifiable additional dimension, particularly in light of the substantial sampling difficulties discussed below.

For terminals, the factor-regression coefficients exhibit considerable uncertainty, with most 94% highest-density intervals including zero. In contrast to the baseline rank-one specification, the previously strong positive associations with terminal capacity and lagged exporter status are not clearly recovered. The response loadings are similarly uncertain, with many intervals spanning zero, making it difficult to identify distinct interpretations for the two factors. The posterior predictive checks also indicate that several discrepancies identified under the simpler specifications persist. In particular, the positive association between articulation-point and incident-global-bridge indicators remains underestimated, with an observed Spearman correlation of 0.290 compared with a replicated median of 0.166. The association between PageRank

and dependence is likewise underestimated (0.314 versus 0.196).

For chokepoints, the factor-regression coefficients are generally small and uncertain, with no clear evidence of systematic associations between the available covariates and either latent dimension. The response loadings suggest a stronger relationship between the first factor and PageRank, betweenness, and monthly flow share, whereas the second-factor loadings are considerably less stable. Descriptive correlations between posterior mean factor coordinates and observed responses are consistent with this distinction, although they should not be interpreted as independent evidence of model adequacy. Compared with the baseline rank-one model, the joint posterior predictive checks show closer agreement for several response pairs. For example, the observed betweenness–flow-share correlation of 0.800 is reproduced with a median of 0.785, while the PageRank–flow-share correlation of 0.672 has a replicated median of 0.646. Nevertheless, some discrepancies remain, including the association between PageRank and articulation-point status.

The sampling diagnostics raise substantial concerns for both specifications. The terminal model exhibits 8,096 post-warm-up divergences, a maximum $\widehat{R}$ of 1.642, and a minimum bulk ESS of 6.8. The corresponding chokepoint model exhibits 2,630 divergences, a maximum $\widehat{R}$ of 1.372, and a minimum bulk ESS of 9.1. In both cases, the poorest mixing is associated with latent-factor coordinates and response loadings, indicating difficulties in adequately exploring the posterior distribution. These findings are also reflected in the Geweke and Heidelberger–Welch diagnostics, particularly for terminals.

Overall, although the rank-two specification appears to improve the reproduction of some joint response relationships, especially for chokepoints, the severe convergence and mixing difficulties prevent a reliable assessment of whether the additional latent dimension provides a meaningful improvement over the baseline and random-intercept specifications.

Table 24: Terminal rank-two model: factor-regression coefficients.

| Parameter | Mean | 94% HDI | $\widehat{R}$ |
|---|---|---|---|
| Regression [Lagged neighbour activity, F1] | -0.063 | [-0.184, 0.159] | 1.577 |
| Regression [Lagged neighbour activity, F2] | -0.108 | [-0.197, -0.033] | 1.231 |
| Regression [Log lagged voyage distance, F1] | -0.068 | [-0.171, 0.109] | 1.522 |
| Regression [Log lagged voyage distance, F2] | -0.069 | [-0.149, 0.006] | 1.199 |
| Regression [Month sine, F1] | -0.026 | [-0.072, 0.020] | 1.067 |
| Regression [Month sine, F2] | 0.023 | [-0.049, 0.076] | 1.360 |
| Regression [Month cosine, F1] | -0.038 | [-0.093, 0.032] | 1.281 |
| Regression [Month cosine, F2] | -0.012 | [-0.070, 0.041] | 1.268 |
| Regression [Log terminal capacity, F1] | 0.028 | [-0.238, 0.415] | 1.554 |
| Regression [Log terminal capacity, F2] | -0.315 | [-0.468, 0.105] | 1.388 |
| Regression [Terminal age, F1] | -0.106 | [-0.193, 0.059] | 1.388 |
| Regression [Terminal age, F2] | -0.024 | [-0.154, 0.078] | 1.457 |
| Regression [Lagged exporter, F1] | 0.645 | [0.008, 1.412] | 1.326 |
| Regression [Lagged exporter, F2] | -0.823 | [-1.497, 1.018] | 1.609 |

Table 25: Terminal rank-two model: continuous response loadings.

| Parameter | Mean | 94% HDI | $\widehat{R}$ |
|---|---|---|---|
| Continuous loading[PageRank, F1] | 0.174 | [0.010, 0.374] | 1.320 |
| Continuous loading[PageRank, F2] | -0.220 | [-0.385, 0.273] | 1.642 |
| Continuous loading[HHI, F1] | -0.118 | [-0.551, 0.202] | 1.491 |

| Parameter | Mean | 94% HDI | $\widehat{R}$ |
|---|---|---|---|
| Continuous loading[HHI, F2] | 0.404 | [-0.236, 0.600] | 1.524 |
| Continuous loading[Dependence, F1] | 0.193 | [-0.007, 0.436] | 1.340 |
| Continuous loading[Dependence, F2] | -0.261 | [-0.463, 0.305] | 1.619 |

Table 26: Terminal rank-two model: binary response loadings.

| Parameter | Mean | 94% HDI | $\widehat{R}$ |
|---|---|---|---|
| Binary loading[Articulation point, F1] | 1.516 | [-0.385, 2.346] | 1.342 |
| Binary loading[Articulation point, F2] | 0.026 | [-1.249, 1.975] | 1.576 |
| Binary loading[Incident global bridge, F1] | 3.174 | [-5.439, 7.035] | 1.606 |
| Binary loading[Incident global bridge, F2] | 3.555 | [1.023, 6.684] | 1.249 |
| Binary loading[Connected to articulation point, F1] | 0.005 | [-0.176, 0.180] | 1.131 |
| Binary loading[Connected to articulation point, F2] | -0.074 | [-0.222, 0.086] | 1.097 |

Table 27: Terminal rank-two model: boundary-component loadings.

| Parameter | Mean | 94% HDI | $\widehat{R}$ |
|---|---|---|---|
| One-inflation loading[HHI, F1] | 0.291 | [-6.419, 4.535] | 1.597 |
| One-inflation loading[HHI, F2] | 4.931 | [-0.807, 7.582] | 1.326 |
| One-inflation loading[Dependence, F1] | -0.023 | [-0.495, 0.317] | 1.537 |
| One-inflation loading[Dependence, F2] | 0.368 | [-0.118, 0.588] | 1.347 |

Table 28: Terminal rank-two model: response intercepts and precisions.

| Parameter | Mean | 94% HDI | $\widehat{R}$ |
|---|---|---|---|
| Continuous intercept[PageRank] | -4.265 | [-4.282, -4.248] | 1.001 |
| Continuous intercept[HHI] | 0.009 | [-0.034, 0.052] | 1.001 |
| Continuous intercept[Dependence] | -0.833 | [-0.870, -0.797] | 1.001 |
| Binary intercept[Articulation point] | -4.138 | [-4.579, -3.715] | 1.001 |
| Binary intercept[Incident global bridge] | 2.010 | [1.577, 2.458] | 1.002 |
| Binary intercept[Connected to articulation point] | 2.830 | [2.680, 2.977] | 1.000 |
| One-inflation intercept[HHI] | -1.658 | [-2.056, -1.265] | 1.002 |
| One-inflation intercept[Dependence] | -1.258 | [-1.346, -1.171] | 1.000 |
| Beta precision[PageRank] | 1587.156 | [1332.244, 1859.695] | 1.002 |
| Beta precision[HHI] | 18.174 | [16.747, 19.676] | 1.000 |
| Beta precision[Dependence] | 6.818 | [6.459, 7.181] | 1.002 |

Table 29: Terminal rank-two model: overall MCMC diagnostics.

| Diagnostic | Value |
|---|---|
| Divergences | 8096 |
| Maximum $\widehat{R}$ | 1.642 |
| Minimum bulk ESS | 6.8 |
| Minimum tail ESS | 12.9 |
| Geweke passed | 84/188 |
| Maximum $\lvert Z\rvert$ (Geweke) | 17.63 |
| Heidelberger–Welch stationarity passed | 156/188 |
| Heidelberger–Welch halfwidth passed | 72/188 |
| Heidelberger–Welch halfwidth not evaluated | 32 |

| Diagnostic | Value |
|---|---|
| Basic sampling checks passed | False |

Table 30: Terminal rank-two model: twelve parameters with the lowest bulk ESS (including observation-level parameters).

| Parameter | $\widehat{R}$ | Bulk ESS | Tail ESS |
|---|---|---|---|
| PageRank_second_loading | 1.642 | 6.8 | 16.4 |
| Continuous loading[PageRank, F2] | 1.642 | 6.8 | 16.4 |
| factor[1676, F2] | 1.615 | 6.9 | 14.9 |
| factor[293, F2] | 1.619 | 6.9 | 15.4 |
| factor[125, F2] | 1.618 | 6.9 | 14.9 |
| factor[674, F2] | 1.616 | 6.9 | 14.7 |
| factor[953, F2] | 1.617 | 6.9 | 15.2 |
| factor[209, F2] | 1.620 | 6.9 | 15.4 |
| factor[568, F2] | 1.616 | 6.9 | 15.2 |
| other_loadings[Dependence, F2] | 1.619 | 6.9 | 15.7 |
| Continuous loading[Dependence, F2] | 1.619 | 6.9 | 15.7 |
| factor[342, F2] | 1.614 | 6.9 | 15.3 |

Table 31: Terminal rank-two model: divergences by chain.

| chain | draws | divergences | divergence_rate |
|---|---|---|---|
| 0 | 30000 | 1290 | 0.043 |
| 1 | 30000 | 3349 | 0.1116333333333333 |
| 2 | 30000 | 98 | 0.0032666666666666 |
| 3 | 30000 | 3359 | 0.1119666666666666 |

Table 32: Terminal rank-two model: univariate posterior predictive checks (95% replicated intervals).

| Group / response | Stat. | Observed | Replicated 95% | Outside |
|---|---|---|---|---|
| All / PageRank | mean | 0.0185 | [0.0184, 0.0187] | No |
| All / PageRank | sd | 0.0133 | [0.0124, 0.0130] | Yes |

| Group / response | Stat. | Observed | Replicated 95% | Outside |
|---|---|---|---|---|
| All / PageRank | q05 | 0.0089 | [0.0073, 0.0079] | Yes |
| All / PageRank | q50 | 0.0147 | [0.0150, 0.0154] | Yes |
| All / PageRank | q95 | 0.0391 | [0.0384, 0.0405] | No |
| All / HHI | mean | 0.5887 | [0.5838, 0.5958] | No |
| All / HHI | sd | 0.3254 | [0.3188, 0.3267] | No |
| All / HHI | one_share | 0.3409 | [0.3301, 0.3510] | No |
| All / HHI | q05 | 0.1268 | [0.1238, 0.1430] | No |
| All / HHI | q50 | 0.5045 | [0.4896, 0.5113] | No |
| All / HHI | q95 | 1.0000 | [1.0000, 1.0000] | No |
| All / Dependence | mean | 0.4948 | [0.4821, 0.5098] | No |
| All / Dependence | sd | 0.3219 | [0.3120, 0.3247] | No |
| All / Dependence | one_share | 0.2076 | [0.1917, 0.2251] | No |
| All / Dependence | q05 | 0.0716 | [0.0723, 0.0892] | Yes |
| All / Dependence | q50 | 0.4283 | [0.3983, 0.4342] | No |
| All / Dependence | q95 | 1.0000 | [1.0000, 1.0000] | No |
| Importer / PageRank | mean | 0.0145 | [0.0141, 0.0144] | Yes |
| Importer / PageRank | sd | 0.0045 | [0.0051, 0.0054] | Yes |
| Importer / PageRank | q05 | 0.0089 | [0.0069, 0.0074] | Yes |
| Importer / PageRank | q50 | 0.0136 | [0.0133, 0.0138] | No |
| Importer / PageRank | q95 | 0.0233 | [0.0234, 0.0246] | Yes |
| Importer / HHI | mean | 0.6745 | [0.6597, 0.6755] | No |
| Importer / HHI | sd | 0.2966 | [0.2949, 0.3027] | No |
| Importer / HHI | one_share | 0.4189 | [0.4000, 0.4271] | No |
| Importer / HHI | q05 | 0.2446 | [0.2267, 0.2499] | No |
| Importer / HHI | q50 | 0.5701 | [0.5721, 0.6016] | Yes |
| Importer / HHI | q95 | 1.0000 | [1.0000, 1.0000] | No |
| Importer / Dependence | mean | 0.4812 | [0.4530, 0.4880] | No |
| Importer / Dependence | sd | 0.3574 | [0.3234, 0.3379] | Yes |
| Importer / Dependence | one_share | 0.2645 | [0.2129, 0.2538] | Yes |
| Importer / Dependence | q05 | 0.0614 | [0.0594, 0.0787] | No |
| Importer / Dependence | q50 | 0.3472 | [0.3504, 0.3874] | Yes |
| Importer / Dependence | q95 | 1.0000 | [1.0000, 1.0000] | No |
| Exporter / PageRank | mean | 0.0269 | [0.0270, 0.0276] | Yes |
| Exporter / PageRank | sd | 0.0198 | [0.0174, 0.0187] | Yes |
| Exporter / PageRank | q05 | 0.0088 | [0.0090, 0.0100] | Yes |
| Exporter / PageRank | q50 | 0.0210 | [0.0221, 0.0232] | Yes |
| Exporter / PageRank | q95 | 0.0714 | [0.0624, 0.0683] | Yes |
| Exporter / HHI | mean | 0.4119 | [0.4180, 0.4380] | Yes |
| Exporter / HHI | sd | 0.3103 | [0.3026, 0.3181] | No |
| Exporter / HHI | one_share | 0.1800 | [0.1746, 0.2013] | No |
| Exporter / HHI | q05 | 0.0877 | [0.0532, 0.0745] | Yes |

| Group / response | Stat. | Observed | Replicated 95% | Outside |
|---|---|---|---|---|
| Exporter / HHI | q50 | 0.3333 | [0.3252, 0.3504] | No |
| Exporter / HHI | q95 | 1.0000 | [1.0000, 1.0000] | No |
| Exporter / Dependence | mean | 0.5228 | [0.5258, 0.5666] | Yes |
| Exporter / Dependence | sd | 0.2296 | [0.2756, 0.2933] | Yes |
| Exporter / Dependence | one_share | 0.0904 | [0.1321, 0.1871] | Yes |
| Exporter / Dependence | q05 | 0.1852 | [0.1083, 0.1446] | Yes |
| Exporter / Dependence | q50 | 0.4972 | [0.4840, 0.5400] | No |
| Exporter / Dependence | q95 | 1.0000 | [1.0000, 1.0000] | No |

Table 33: Terminal rank-two model: joint posterior predictive Spearman correlations (200 replicates).

| Group / pair | Observed | Rep. median | Rep. 95% | Outside |
|---|---|---|---|---|
| All / PageRank–HHI | -0.639 | -0.661 | [-0.681, -0.640] | Yes |
| All / PageRank–Dependence | 0.314 | 0.196 | [0.153, 0.236] | Yes |
| All / PageRank–Articulation point | 0.372 | 0.307 | [0.270, 0.338] | Yes |
| All / PageRank–Incident global bridge | -0.238 | -0.201 | [-0.228, -0.171] | Yes |
| All / PageRank–Connected to articulation point | -0.015 | 0.022 | [-0.023, 0.063] | No |
| All / HHI–Dependence | -0.171 | -0.138 | [-0.184, -0.100] | No |
| All / HHI–Articulation point | -0.290 | -0.159 | [-0.195, -0.117] | Yes |
| All / HHI–Incident global bridge | 0.533 | 0.502 | [0.475, 0.526] | Yes |
| All / HHI–Connected to articulation point | -0.038 | -0.023 | [-0.062, 0.022] | No |
| All / Dependence–Articulation point | 0.157 | 0.121 | [0.087, 0.154] | Yes |
| All / Dependence–Incident global bridge | -0.066 | 0.009 | [-0.031, 0.051] | Yes |
| All / Dependence–Connected to articulation point | -0.082 | 0.005 | [-0.034, 0.033] | Yes |
| All / Articulation point–Incident global bridge | 0.290 | 0.166 | [0.132, 0.192] | Yes |

| Group / pair | Observed | Rep. median | Rep. 95% | Outside |
|---|---|---|---|---|
| All / Articulation point–Connected to articulation point | -0.063 | 0.007 | [-0.033, 0.034] | Yes |
| All / Incident global bridge–Connected to articulation point | 0.060 | -0.017 | [-0.056, 0.026] | Yes |
| Importer / PageRank–HHI | -0.416 | -0.503 | [-0.531, -0.471] | Yes |
| Importer / PageRank–Dependence | 0.209 | 0.103 | [0.060, 0.145] | Yes |
| Importer / PageRank–Articulation point | 0.232 | 0.116 | [0.076, 0.154] | Yes |
| Importer / PageRank–Incident global bridge | -0.258 | -0.193 | [-0.229, -0.155] | Yes |
| Importer / PageRank–Connected to articulation point | -0.043 | 0.013 | [-0.033, 0.050] | Yes |
| Importer / HHI–Dependence | -0.163 | -0.046 | [-0.092, 0.007] | Yes |
| Importer / HHI–Articulation point | -0.137 | 0.053 | [0.012, 0.096] | Yes |
| Importer / HHI–Incident global bridge | 0.644 | 0.577 | [0.546, 0.603] | Yes |
| Importer / HHI–Connected to articulation point | 0.014 | -0.017 | [-0.058, 0.038] | No |
| Importer / Dependence–Articulation point | 0.033 | 0.044 | [0.008, 0.083] | No |
| Importer / Dependence–Incident global bridge | -0.154 | 0.024 | [-0.021, 0.073] | Yes |
| Importer / Dependence–Connected to articulation point | -0.045 | 0.000 | [-0.051, 0.038] | No |
| Importer / Articulation point–Incident global bridge | 0.137 | 0.150 | [0.120, 0.175] | No |
| Importer / Articulation point–Connected to articulation point | 0.019 | 0.002 | [-0.050, 0.039] | No |

| Group / pair | Observed | Rep. median | Rep. 95% | Outside |
|---|---|---|---|---|
| Importer / Incident global bridge–Connected to articulation point | 0.131 | -0.013 | [-0.060, 0.030] | Yes |
| Exporter / PageRank–HHI | -0.818 | -0.763 | [-0.783, -0.740] | Yes |
| Exporter / PageRank–Dependence | 0.516 | 0.300 | [0.238, 0.360] | Yes |
| Exporter / PageRank–Articulation point | 0.276 | 0.389 | [0.337, 0.449] | Yes |
| Exporter / PageRank–Incident global bridge | -0.198 | -0.213 | [-0.249, -0.173] | No |
| Exporter / PageRank–Connected to articulation point | 0.166 | 0.033 | [-0.029, 0.091] | Yes |
| Exporter / HHI–Dependence | -0.102 | -0.232 | [-0.295, -0.182] | Yes |
| Exporter / HHI–Articulation point | -0.159 | -0.214 | [-0.280, -0.145] | No |
| Exporter / HHI–Incident global bridge | 0.398 | 0.417 | [0.368, 0.458] | No |
| Exporter / HHI–Connected to articulation point | -0.230 | -0.031 | [-0.098, 0.038] | Yes |
| Exporter / Dependence–Articulation point | 0.307 | 0.170 | [0.111, 0.223] | Yes |
| Exporter / Dependence–Incident global bridge | 0.195 | -0.015 | [-0.076, 0.050] | Yes |
| Exporter / Dependence–Connected to articulation point | -0.106 | 0.012 | [-0.044, 0.061] | Yes |
| Exporter / Articulation point–Incident global bridge | 0.546 | 0.239 | [0.179, 0.296] | Yes |
| Exporter / Articulation point–Connected to articulation point | -0.049 | 0.011 | [-0.054, 0.065] | No |
| Exporter / Incident global bridge–Connected to articulation point | -0.047 | -0.021 | [-0.084, 0.049] | No |

Table 34: Terminal rank-two model: descriptive associations with posterior mean latent coordinates (orientation-dependent).

| Group | Factor | Response | Spearman $\rho$ |
|---|---|---|---|
| All | F1 | PageRank | 0.568 |
| All | F1 | HHI | -0.085 |
| All | F1 | Dependence | 0.280 |
| All | F1 | Articulation point | 0.508 |
| All | F1 | Incident global bridge | 0.572 |
| All | F1 | Connected to articulation point | 0.012 |
| All | F2 | PageRank | -0.752 |
| All | F2 | HHI | 0.917 |
| All | F2 | Dependence | -0.202 |
| All | F2 | Articulation point | -0.190 |
| All | F2 | Incident global bridge | 0.688 |
| All | F2 | Connected to articulation point | -0.007 |
| Importer | F1 | PageRank | 0.352 |
| Importer | F1 | HHI | 0.375 |
| Importer | F1 | Dependence | 0.115 |
| Importer | F1 | Articulation point | 0.276 |
| Importer | F1 | Incident global bridge | 0.760 |
| Importer | F1 | Connected to articulation point | 0.094 |
| Importer | F2 | PageRank | -0.623 |
| Importer | F2 | HHI | 0.886 |
| Importer | F2 | Dependence | -0.178 |
| Importer | F2 | Articulation point | -0.076 |
| Importer | F2 | Incident global bridge | 0.797 |
| Importer | F2 | Connected to articulation point | 0.061 |
| Exporter | F1 | PageRank | 0.678 |
| Exporter | F1 | HHI | -0.349 |
| Exporter | F1 | Dependence | 0.653 |
| Exporter | F1 | Articulation point | 0.675 |
| Exporter | F1 | Incident global bridge | 0.533 |
| Exporter | F1 | Connected to articulation point | 0.035 |
| Exporter | F2 | PageRank | -0.868 |
| Exporter | F2 | HHI | 0.913 |
| Exporter | F2 | Dependence | -0.227 |
| Exporter | F2 | Articulation point | 0.031 |
| Exporter | F2 | Incident global bridge | 0.590 |
| Exporter | F2 | Connected to articulation point | -0.216 |


PageRank
Counterparty HHI
Dependence
Expected response
Latent factor 1 (other factor = 0)
Is Articulation Point
Has Incident Global Bridge
Adjacent to articulation point
Probability
Latent factor 1 (other factor = 0)
PageRank
Counterparty HHI
Dependence
Expected response
Latent factor 2 (other factor = 0)
Is Articulation Point
Has Incident Global Bridge
Adjacent to articulation point
Probability
Latent factor 2 (other factor = 0)


Lagged exporter status, Factor 2 — Regression coefficient
Chain 1
Chain 2
Chain 3
Chain 4
Autocorrelation
Lag


PageRank, Factor 2 — Continuous-response loading
Chain 1
Chain 2
Chain 3
Chain 4
Autocorrelation
Lag
Lagged neighbour activity, Factor 1 — Regression coefficient
PageRank, Factor 1 — Continuous-response loading


Lagged neighbour activity, Factor 1 — Regression coefficient
Parameter value
Lagged neighbour activity, Factor 1 — Regression coefficient
Iteration
Lagged neighbour activity, Factor 2 — Regression coefficient
Parameter value
Lagged neighbour activity, Factor 2 — Regression coefficient
Iteration
Lagged voyage distance, Factor 1 — Regression coefficient
Parameter value
Lagged voyage distance, Factor 1 — Regression coefficient
Iteration
Lagged voyage distance, Factor 2 — Regression coefficient
Parameter value
Lagged voyage distance, Factor 2 — Regression coefficient
Iteration
Chain 1
Chain 2
Chain 3
Chain 4
Seasonality (sine), Factor 1 — Regression coefficient
Parameter value
Seasonality (sine), Factor 1 — Regression coefficient
Iteration
Seasonality (sine), Factor 2 — Regression coefficient
Parameter value
Seasonality (sine), Factor 2 — Regression coefficient
Iteration
Seasonality (cosine), Factor 1 — Regression coefficient
Parameter value
Seasonality (cosine), Factor 1 — Regression coefficient
Iteration
Seasonality (cosine), Factor 2 — Regression coefficient
Parameter value
Seasonality (cosine), Factor 2 — Regression coefficient
Iteration
Chain 1
Chain 2
Chain 3
Chain 4


Terminal capacity, Factor 1 — Regression coefficient
Parameter value
Terminal capacity, Factor 1 — Regression coefficient
Iteration
Terminal capacity, Factor 2 — Regression coefficient
Parameter value
Terminal capacity, Factor 2 — Regression coefficient
Iteration
Terminal age, Factor 1 — Regression coefficient
Parameter value
Terminal age, Factor 1 — Regression coefficient
Iteration
Terminal age, Factor 2 — Regression coefficient
Parameter value
Terminal age, Factor 2 — Regression coefficient
Iteration
Chain 1
Chain 2
Chain 3
Chain 4


Lagged exporter status, Factor 1 — Regression coefficient
Parameter value
Lagged exporter status, Factor 1 — Regression coefficient
Iteration
Lagged exporter status, Factor 2 — Regression coefficient
Parameter value
Lagged exporter status, Factor 2 — Regression coefficient
Iteration
PageRank, Factor 1 — Continuous-response loading
Parameter value
PageRank, Factor 1 — Continuous-response loading
Iteration
PageRank, Factor 2 — Continuous-response loading
Parameter value
PageRank, Factor 2 — Continuous-response loading
Iteration
Chain 1
Chain 2
Chain 3
Chain 4


Counterparty HHI, Factor 1 — Continuous-response loading
Parameter value
Counterparty HHI, Factor 1 — Continuous-response loading
Iteration
Counterparty HHI, Factor 2 — Continuous-response loading
Parameter value
Counterparty HHI, Factor 2 — Continuous-response loading
Iteration
Dependence, Factor 1 — Continuous-response loading
Parameter value
Dependence, Factor 1 — Continuous-response loading
Iteration
Dependence, Factor 2 — Continuous-response loading
Parameter value
Dependence, Factor 2 — Continuous-response loading
Iteration
Chain 1
Chain 2
Chain 3
Chain 4
Is Articulation Point, Factor 1 — Binary-response loading
Parameter value
Is Articulation Point, Factor 1 — Binary-response loading
Iteration
Is Articulation Point, Factor 2 — Binary-response loading
Parameter value
Is Articulation Point, Factor 2 — Binary-response loading
Iteration
Has Incident Global Bridge, Factor 1 — Binary-response loading
Parameter value
Has Incident Global Bridge, Factor 1 — Binary-response loading
Iteration
Has Incident Global Bridge, Factor 2 — Binary-response loading
Parameter value
Has Incident Global Bridge, Factor 2 — Binary-response loading
Iteration
Chain 1
Chain 2
Chain 3
Chain 4


Adjacent to articulation point, Factor 1 — Binary-response loading
Parameter value
Iteration
Adjacent to articulation point, Factor 2 — Binary-response loading
Parameter value
Iteration
PageRank — Continuous-response intercept
Parameter value
Iteration
Counterparty HHI — Continuous-response intercept
Parameter value
Iteration
Chain 1
Chain 2
Chain 3
Chain 4
Dependence — Continuous-response intercept
Parameter value
Iteration
Is Articulation Point — Binary-response intercept
Parameter value
Iteration
Has Incident Global Bridge — Binary-response intercept
Parameter value
Iteration
Adjacent to articulation point — Binary-response intercept
Parameter value
Iteration
Chain 1
Chain 2
Chain 3
Chain 4


PageRank — Beta precision
Parameter value
PageRank — Beta precision
Iteration
Counterparty HHI — Beta precision
Parameter value
Counterparty HHI — Beta precision
Iteration
Dependence — Beta precision
Parameter value
Dependence — Beta precision
Iteration
Counterparty HHI — One-mass intercept
Parameter value
Counterparty HHI — One-mass intercept
Iteration
Chain 1
Chain 2
Chain 3
Chain 4
Dependence — One-mass intercept
Parameter value
Dependence — One-mass intercept
Iteration
Counterparty HHI, Factor 1 — One-mass loading
Parameter value
Counterparty HHI, Factor 1 — One-mass loading
Iteration
Counterparty HHI, Factor 2 — One-mass loading
Parameter value
Counterparty HHI, Factor 2 — One-mass loading
Iteration
Dependence, Factor 1 — One-mass loading
Parameter value
Dependence, Factor 1 — One-mass loading
Iteration
Chain 1
Chain 2
Chain 3
Chain 4


Chain 1
Chain 2
Chain 3
Chain 4


Post-warm-up divergences by chain
Divergent transitions
Chain

non-divergent
divergent
beta [log1p_capacity_mtpa, factor_2]
loading_one [counterparty_hhi_terminal, factor_2]
loading_cont [role_specific_pagerank, factor_1]
loading_binary [has_incident_global_bridge, factor_2]
loading_cont [counterparty_hhi_terminal, factor_1]
loading_one [counterparty_hhi_terminal, factor_1]
loading_cont [role_specific_pagerank, factor_2]
loading_binary [is_articulation_point, factor_2]




Dashed lines: unadjusted two-sided 5% threshold. NA: diagnostic unavailable.




Dashed lines: unadjusted two-sided 5% threshold. NA: diagnostic unavailable.




Half-width: 95% interval, relative target 0.10; sensitive to means near zero. See CSV for discarded draws.




Half-width: 95% interval, relative target 0.10; sensitive to means near zero. See CSV for discarded draws.

Table 35: Chokepoint rank-two model: factor-regression coefficients.

| Parameter | Mean | 94% HDI | $\widehat{R}$ |
|---|---|---|---|
| Regression [Lagged neighbour activity, F1] | 0.057 | [-0.026, 0.135] | 1.039 |
| Regression [Lagged neighbour activity, F2] | 0.005 | [-0.114, 0.121] | 1.229 |
| Regression [Log lagged voyage distance, F1] | 0.047 | [-0.207, 0.270] | 1.350 |
| Regression [Log lagged voyage distance, F2] | 0.137 | [-0.045, 0.273] | 1.066 |
| Regression [Month sine, F1] | -0.024 | [-0.099, 0.060] | 1.082 |
| Regression [Month sine, F2] | 0.024 | [-0.057, 0.100] | 1.125 |
| Regression [Month cosine, F1] | -0.039 | [-0.144, 0.079] | 1.265 |
| Regression [Month cosine, F2] | -0.049 | [-0.144, 0.062] | 1.116 |

Table 36: Chokepoint rank-two model: continuous response loadings.

| Parameter | Mean | 94% HDI | $\widehat{R}$ |
|---|---|---|---|
| Continuous loading[PageRank, F1] | 0.174 | [0.021, 0.277] | 1.067 |
| Continuous loading[PageRank, F2] | -0.011 | [-0.270, 0.258] | 1.359 |
| Continuous loading[Betweenness, F1] | 0.667 | [0.089, 1.073] | 1.159 |
| Continuous loading[Betweenness, F2] | 0.095 | [-0.977, 1.033] | 1.356 |
| Continuous loading[Monthly flow share, F1] | 0.546 | [-0.196, 0.947] | 1.103 |
| Continuous loading[Monthly flow share, F2] | -0.246 | [-0.947, 0.723] | 1.372 |

Table 37: Chokepoint rank-two model: binary response loadings.

| Parameter | Mean | 94% HDI | $\widehat{R}$ |
|---|---|---|---|
| Binary loading[Articulation point, F1] | 0.476 | [-0.082, 0.866] | 1.058 |
| Binary loading[Articulation point, F2] | -0.133 | [-0.836, 0.706] | 1.353 |
| Binary loading[Connected to articulation point, F1] | 1.057 | [-1.123, 2.506] | 1.344 |
| Binary loading[Connected to articulation point, F2] | 0.901 | [-1.416, 2.473] | 1.219 |

Table 38: Chokepoint rank-two model: boundary-component loadings.

| Parameter | Mean | 94% HDI | $\widehat{R}$ |
|---|---|---|---|
| Zero-inflation loading[Betweenness, F1] | -0.872 | [-2.547, 1.363] | 1.170 |
| Zero-inflation loading[Betweenness, F2] | 0.875 | [-1.309, 2.501] | 1.268 |

Table 39: Chokepoint rank-two model: response intercepts and precisions.

| Parameter | Mean | 94% HDI | $\widehat{R}$ |
|---|---|---|---|
| Continuous intercept[PageRank] | -4.053 | [-4.074, -4.032] | 1.000 |
| Continuous intercept[Betweenness] | -4.788 | [-4.853, -4.727] | 1.001 |
| Continuous intercept[Monthly flow share] | -4.096 | [-4.148, -4.045] | 1.000 |
| Binary intercept[Articulation point] | 1.149 | [1.007, 1.296] | 1.000 |
| Binary intercept[Connected to articulation point] | 3.503 | [2.880, 4.167] | 1.000 |
| Zero-inflation intercept[Betweenness] | -7.031 | [-8.480, -5.591] | 1.000 |

| Parameter | Mean | 94% HDI | $\widehat{R}$ |
|---|---|---|---|
| Beta precision[PageRank] | 792.046 | [726.953, 856.703] | 1.000 |
| Beta precision[Betweenness] | 1019.967 | [764.417, 1288.509] | 1.000 |
| Beta precision[Monthly flow share] | 2509.068 | [1449.260, 3709.450] | 1.004 |

Table 40: Chokepoint rank-two model: overall MCMC diagnostics.

| Diagnostic | Value |
|---|---|
| Divergences | 2630 |
| Maximum $\widehat{R}$ | 1.372 |
| Minimum bulk ESS | 9.1 |
| Minimum tail ESS | 41.5 |
| Geweke passed | 113/140 |
| Maximum $\lvert Z\rvert$ (Geweke) | 4.52 |
| Heidelberger–Welch stationarity passed | 137/140 |
| Heidelberger–Welch halfwidth passed | 43/140 |
| Heidelberger–Welch halfwidth not evaluated | 3 |
| Basic sampling checks passed | False |

Table 41: Chokepoint rank-two model: twelve parameters with the lowest bulk ESS (including observation-level parameters).

| Parameter | $\widehat{R}$ | Bulk ESS | Tail ESS |
|---|---|---|---|
| factor[1129, F1] | 1.364 | 9.1 | 49.9 |
| factor[1131, F1] | 1.364 | 9.2 | 49.9 |
| Continuous loading[Flow share, F2] | 1.372 | 9.2 | 51.1 |
| factor[979, F1] | 1.360 | 9.2 | 48.0 |
| factor[521, F2] | 1.360 | 9.2 | 48.3 |
| factor[1135, F1] | 1.359 | 9.2 | 52.1 |
| factor[860, F1] | 1.361 | 9.2 | 49.3 |
| factor[998, F1] | 1.361 | 9.2 | 51.9 |
| factor[1000, F1] | 1.360 | 9.2 | 45.7 |
| factor[1004, F1] | 1.355 | 9.2 | 45.4 |
| factor[862, F1] | 1.359 | 9.3 | 49.6 |
| factor[962, F1] | 1.357 | 9.3 | 48.0 |

Table 42: Chokepoint rank-two model: divergences by chain.

| chain | draws | divergences | divergence_rate |
|---|---|---|---|
| 0 | 30000 | 0 | 0.0 |
| 1 | 30000 | 2157 | 0.0719 |
| 2 | 30000 | 0 | 0.0 |
| 3 | 30000 | 473 | 0.0157666666666666 |

Table 43: Chokepoint rank-two model: univariate posterior predictive checks (95% replicated intervals).

| Group / response | Stat. | Observed | Replicated 95% | Outside |
|---|---|---|---|---|
| All / PageRank | mean | 0.0176 | [0.0172, 0.0180] | No |
| All / PageRank | sd | 0.0069 | [0.0061, 0.0068] | Yes |
| All / PageRank | q05 | 0.0084 | [0.0078, 0.0088] | No |
| All / PageRank | q50 | 0.0166 | [0.0165, 0.0174] | No |
| All / PageRank | q95 | 0.0302 | [0.0281, 0.0302] | No |
| All / Betweenness | mean | 0.0132 | [0.0130, 0.0135] | No |
| All / Betweenness | sd | 0.0135 | [0.0130, 0.0140] | No |
| All / Betweenness | zero_share | 0.0052 | [0.0017, 0.0121] | No |
| All / Betweenness | q05 | 0.0006 | [0.0006, 0.0010] | Yes |
| All / Betweenness | q50 | 0.0077 | [0.0076, 0.0085] | No |
| All / Betweenness | q95 | 0.0413 | [0.0397, 0.0434] | No |
| All / Monthly flow share | mean | 0.0234 | [0.0231, 0.0236] | No |
| All / Monthly flow share | sd | 0.0201 | [0.0196, 0.0205] | No |
| All / Monthly flow share | q05 | 0.0030 | [0.0029, 0.0035] | No |
| All / Monthly flow share | q50 | 0.0159 | [0.0154, 0.0165] | No |
| All / Monthly flow share | q95 | 0.0643 | [0.0620, 0.0668] | No |

Table 44: Chokepoint rank-two model: joint posterior predictive Spearman correlations (200 replicates).

| Group / pair | Observed | Rep. median | Rep. 95% | Outside |
|---|---|---|---|---|
| All / PageRank–Betweenness | 0.685 | 0.646 | [0.606, 0.681] | Yes |
| All / PageRank–Monthly flow share | 0.672 | 0.646 | [0.599, 0.682] | No |
| All / PageRank–Articulation point | 0.287 | 0.210 | [0.144, 0.267] | Yes |

| Group / pair | Observed | Rep. median | Rep. 95% | Outside |
|---|---|---|---|---|
| All / PageRank–Connected to articulation point | 0.216 | 0.204 | [0.136, 0.260] | No |
| All / Betweenness–Monthly flow share | 0.800 | 0.785 | [0.766, 0.802] | No |
| All / Betweenness–Articulation point | 0.252 | 0.259 | [0.195, 0.329] | No |
| All / Betweenness–Connected to articulation point | 0.320 | 0.320 | [0.259, 0.368] | No |
| All / Monthly flow share–Articulation point | 0.291 | 0.292 | [0.213, 0.359] | No |
| All / Monthly flow share–Connected to articulation point | 0.182 | 0.173 | [0.116, 0.238] | No |
| All / Articulation point–Connected to articulation point | 0.166 | 0.079 | [0.017, 0.145] | Yes |

Table 45: Chokepoint rank-two model: descriptive associations with posterior mean latent coordinates (orientation-dependent).

| Group | Factor | Response | Spearman $\rho$ |
|---|---|---|---|
| All | F1 | PageRank | 0.742 |
| All | F1 | Betweenness | 0.961 |
| All | F1 | Monthly flow share | 0.915 |
| All | F1 | Articulation point | 0.298 |
| All | F1 | Connected to articulation point | 0.346 |
| All | F2 | PageRank | -0.050 |
| All | F2 | Betweenness | 0.152 |
| All | F2 | Monthly flow share | -0.412 |
| All | F2 | Articulation point | -0.065 |
| All | F2 | Connected to articulation point | 0.366 |


PageRank
Counterparty HHI
Dependence
Expected response
Latent factor 1 (other factor = 0)
Is Articulation Point
Has Incident Global Bridge
Adjacent to articulation point
Probability
PageRank
Counterparty HHI
Dependence
Expected response
Latent factor 2 (other factor = 0)
Is Articulation Point
Has Incident Global Bridge
Adjacent to articulation point
Probability


Lagged exporter status, Factor 2 — Regression coefficient
Chain 1
Chain 2
Chain 3
Chain 4
Autocorrelation
Lag


PageRank, Factor 2 — Continuous-response loading
Chain 1
Chain 2
Chain 3
Chain 4
Autocorrelation
Lag
Lagged neighbour activity, Factor 1 — Regression coefficient
PageRank, Factor 1 — Continuous-response loading


Lagged neighbour activity, Factor 1 — Regression coefficient
Parameter value
Lagged neighbour activity, Factor 1 — Regression coefficient
Iteration
Lagged neighbour activity, Factor 2 — Regression coefficient
Parameter value
Lagged neighbour activity, Factor 2 — Regression coefficient
Iteration
Lagged voyage distance, Factor 1 — Regression coefficient
Parameter value
Lagged voyage distance, Factor 1 — Regression coefficient
Iteration
Lagged voyage distance, Factor 2 — Regression coefficient
Parameter value
Lagged voyage distance, Factor 2 — Regression coefficient
Iteration
Chain 1
Chain 2
Chain 3
Chain 4
Seasonality (sine), Factor 1 — Regression coefficient
Parameter value
Seasonality (sine), Factor 1 — Regression coefficient
Iteration
Seasonality (sine), Factor 2 — Regression coefficient
Parameter value
Seasonality (sine), Factor 2 — Regression coefficient
Iteration
Seasonality (cosine), Factor 1 — Regression coefficient
Parameter value
Seasonality (cosine), Factor 1 — Regression coefficient
Iteration
Seasonality (cosine), Factor 2 — Regression coefficient
Parameter value
Seasonality (cosine), Factor 2 — Regression coefficient
Iteration
Chain 1
Chain 2
Chain 3
Chain 4


Terminal capacity, Factor 1 — Regression coefficient
Parameter value
Terminal capacity, Factor 1 — Regression coefficient
Iteration
Terminal capacity, Factor 2 — Regression coefficient
Parameter value
Terminal capacity, Factor 2 — Regression coefficient
Iteration
Terminal age, Factor 1 — Regression coefficient
Parameter value
Terminal age, Factor 1 — Regression coefficient
Iteration
Terminal age, Factor 2 — Regression coefficient
Parameter value
Terminal age, Factor 2 — Regression coefficient
Iteration
Chain 1
Chain 2
Chain 3
Chain 4
Lagged exporter status, Factor 1 — Regression coefficient
Parameter value
Lagged exporter status, Factor 1 — Regression coefficient
Iteration
Lagged exporter status, Factor 2 — Regression coefficient
Parameter value
Lagged exporter status, Factor 2 — Regression coefficient
Iteration
PageRank, Factor 1 — Continuous-response loading
Parameter value
PageRank, Factor 1 — Continuous-response loading
Iteration
PageRank, Factor 2 — Continuous-response loading
Parameter value
PageRank, Factor 2 — Continuous-response loading
Iteration
Chain 1
Chain 2
Chain 3
Chain 4


Counterparty HHI, Factor 1 — Continuous-response loading
Parameter value
Counterparty HHI, Factor 1 — Continuous-response loading
Iteration
Counterparty HHI, Factor 2 — Continuous-response loading
Parameter value
Counterparty HHI, Factor 2 — Continuous-response loading
Iteration
Dependence, Factor 1 — Continuous-response loading
Parameter value
Dependence, Factor 1 — Continuous-response loading
Iteration
Dependence, Factor 2 — Continuous-response loading
Parameter value
Dependence, Factor 2 — Continuous-response loading
Iteration
Chain 1
Chain 2
Chain 3
Chain 4
Is Articulation Point, Factor 1 — Binary-response loading
Parameter value
Is Articulation Point, Factor 1 — Binary-response loading
Iteration
Is Articulation Point, Factor 2 — Binary-response loading
Parameter value
Is Articulation Point, Factor 2 — Binary-response loading
Iteration
Has Incident Global Bridge, Factor 1 — Binary-response loading
Parameter value
Has Incident Global Bridge, Factor 1 — Binary-response loading
Iteration
Has Incident Global Bridge, Factor 2 — Binary-response loading
Parameter value
Has Incident Global Bridge, Factor 2 — Binary-response loading
Iteration
Chain 1
Chain 2
Chain 3
Chain 4


Adjacent to articulation point, Factor 1 — Binary-response loading
Parameter value
Adjacent to articulation point, Factor 1 — Binary-response loading
Iteration
Adjacent to articulation point, Factor 2 — Binary-response loading
Parameter value
Adjacent to articulation point, Factor 2 — Binary-response loading
Iteration
PageRank — Continuous-response intercept
Parameter value
PageRank — Continuous-response intercept
Iteration
Counterparty HHI — Continuous-response intercept
Parameter value
Counterparty HHI — Continuous-response intercept
Iteration
Chain 1
Chain 2
Chain 3
Chain 4
Dependence — Continuous-response intercept
Parameter value
Dependence — Continuous-response intercept
Iteration
Is Articulation Point — Binary-response intercept
Parameter value
Is Articulation Point — Binary-response intercept
Iteration
Has Incident Global Bridge — Binary-response intercept
Parameter value
Has Incident Global Bridge — Binary-response intercept
Iteration
Adjacent to articulation point — Binary-response intercept
Parameter value
Adjacent to articulation point — Binary-response intercept
Iteration
Chain 1
Chain 2
Chain 3
Chain 4


PageRank — Beta precision
Parameter value
PageRank — Beta precision
Iteration
Counterparty HHI — Beta precision
Parameter value
Counterparty HHI — Beta precision
Iteration
Dependence — Beta precision
Parameter value
Dependence — Beta precision
Iteration
Counterparty HHI — One-mass intercept
Parameter value
Counterparty HHI — One-mass intercept
Iteration
Chain 1
Chain 2
Chain 3
Chain 4
Dependence — One-mass intercept
Parameter value
Dependence — One-mass intercept
Iteration
Counterparty HHI, Factor 1 — One-mass loading
Parameter value
Counterparty HHI, Factor 1 — One-mass loading
Iteration
Counterparty HHI, Factor 2 — One-mass loading
Parameter value
Counterparty HHI, Factor 2 — One-mass loading
Iteration
Dependence, Factor 1 — One-mass loading
Parameter value
Dependence, Factor 1 — One-mass loading
Iteration
Chain 1
Chain 2
Chain 3
Chain 4







Dashed lines: unadjusted two-sided 5% threshold. NA: diagnostic unavailable.




Dashed lines: unadjusted two-sided 5% threshold. NA: diagnostic unavailable.




Half-width: 95% interval, relative target 0.10; sensitive to means near zero. See CSV for discarded draws.




Half-width: 95% interval, relative target 0.10; sensitive to means near zero. See CSV for discarded draws.




PageRank
Betweenness centrality
Monthly flow share
Is Articulation Point
Adjacent to articulation point
Expected response
Probability
Latent factor 2 (other factor = 0)


Lagged voyage distance, Factor 1 — Regression coefficient
Autocorrelation
Lag
Chain 1
Chain 2
Chain 3
Chain 4


Monthly flow share, Factor 2 — Continuous-response loading
Autocorrelation
Lag
Chain 1
Chain 2
Chain 3
Chain 4


Is Articulation Point, Factor 2 — Binary-response loading
Autocorrelation
Lag
Chain 1
Chain 2
Chain 3
Chain 4


PageRank, Factor 1 — Continuous-response loading
Autocorrelation
Lag
Chain 1
Chain 2
Chain 3
Chain 4


PageRank, Factor 2 — Continuous-response loading
Autocorrelation
Lag
Chain 1
Chain 2
Chain 3
Chain 4


Lagged neighbour activity, Factor 1 — Regression coefficient
Parameter value
Lagged neighbour activity, Factor 1 — Regression coefficient
Iteration
Lagged neighbour activity, Factor 2 — Regression coefficient
Parameter value
Lagged neighbour activity, Factor 2 — Regression coefficient
Iteration
Lagged voyage distance, Factor 1 — Regression coefficient
Parameter value
Lagged voyage distance, Factor 1 — Regression coefficient
Iteration
Lagged voyage distance, Factor 2 — Regression coefficient
Parameter value
Lagged voyage distance, Factor 2 — Regression coefficient
Iteration
Chain 1
Chain 2
Chain 3
Chain 4


Seasonality (sine), Factor 1 — Regression coefficient
Parameter value
Seasonality (sine), Factor 1 — Regression coefficient
Iteration
Seasonality (sine), Factor 2 — Regression coefficient
Parameter value
Seasonality (sine), Factor 2 — Regression coefficient
Iteration
Seasonality (cosine), Factor 1 — Regression coefficient
Parameter value
Seasonality (cosine), Factor 1 — Regression coefficient
Iteration
Seasonality (cosine), Factor 2 — Regression coefficient
Parameter value
Seasonality (cosine), Factor 2 — Regression coefficient
Iteration
Chain 1
Chain 2
Chain 3
Chain 4

PageRank, Factor 1 — Continuous-response loading
Parameter value
PageRank, Factor 1 — Continuous-response loading
Iteration
PageRank, Factor 2 — Continuous-response loading
Parameter value
PageRank, Factor 2 — Continuous-response loading
Iteration
Betweenness centrality, Factor 1 — Continuous-response loading
Parameter value
Betweenness centrality, Factor 1 — Continuous-response loading
Iteration
Betweenness centrality, Factor 2 — Continuous-response loading
Parameter value
Betweenness centrality, Factor 2 — Continuous-response loading
Iteration
Chain 1
Chain 2
Chain 3
Chain 4


Monthly flow share, Factor 1 — Continuous-response loading
Parameter value
Monthly flow share, Factor 1 — Continuous-response loading
Iteration
Monthly flow share, Factor 2 — Continuous-response loading
Parameter value
Monthly flow share, Factor 2 — Continuous-response loading
Iteration
Is Articulation Point, Factor 1 — Binary-response loading
Parameter value
Is Articulation Point, Factor 1 — Binary-response loading
Iteration
Is Articulation Point, Factor 2 — Binary-response loading
Parameter value
Is Articulation Point, Factor 2 — Binary-response loading
Iteration
Chain 1
Chain 2
Chain 3
Chain 4


Adjacent to articulation point, Factor 1 — Binary-response loading
Parameter value
Adjacent to articulation point, Factor 1 — Binary-response loading
Iteration
Adjacent to articulation point, Factor 2 — Binary-response loading
Parameter value
Adjacent to articulation point, Factor 2 — Binary-response loading
Iteration
PageRank — Continuous-response intercept
Parameter value
PageRank — Continuous-response intercept
Iteration
Betweenness centrality — Continuous-response intercept
Parameter value
Betweenness centrality — Continuous-response intercept
Iteration
Chain 1
Chain 2
Chain 3
Chain 4


Monthly flow share — Continuous-response intercept
Parameter value
Monthly flow share — Continuous-response intercept
Iteration
Is Articulation Point — Binary-response intercept
Parameter value
Is Articulation Point — Binary-response intercept
Iteration
Adjacent to articulation point — Binary-response intercept
Parameter value
Adjacent to articulation point — Binary-response intercept
Iteration
PageRank — Beta precision
Parameter value
PageRank — Beta precision
Iteration
Chain 1
Chain 2
Chain 3
Chain 4
Betweenness centrality — Beta precision
Parameter value
Betweenness centrality — Beta precision
Iteration
Monthly flow share — Beta precision
Parameter value
Monthly flow share — Beta precision
Iteration
Betweenness centrality — Zero-mass intercept
Parameter value
Betweenness centrality — Zero-mass intercept
Iteration
Betweenness centrality, Factor 1 — Zero-mass loading
Parameter value
Betweenness centrality, Factor 1 — Zero-mass loading
Iteration
Chain 1
Chain 2
Chain 3
Chain 4







Dashed lines: unadjusted two-sided 5% threshold. NA: diagnostic unavailable.




Dashed lines: unadjusted two-sided 5% threshold. NA: diagnostic unavailable.




Half-width: 95% interval, relative target 0.10; sensitive to means near zero. See CSV for discarded draws.




Half-width: 95% interval, relative target 0.10; sensitive to means near zero. See CSV for discarded draws.

## 6.4 Prior sensitivity results

To assess the implications of the prior assumptions and the sensitivity of posterior findings, two alternative prior specifications were investigated using the same estimation samples. The analysis considers both the prior predictive distributions and the sensitivity of posterior inference to these modifications.

### 6.4.1 Alternative prior specifications

Three prior specifications were considered: the original specification (ORG), described above; a concentrated specification (CON), with more concentrated priors for selected parameters; and a response-specific precision specification (RS), allowing prior distributions to vary across response variables. The alternative specifications were informed by the observed response distributions and prior predictive simulations, rather than independently elicited prior information.

Under the original priors, prior predictive simulations generated relatively broad distributions for several network measures, particularly those with small observed values, such as PageRank, betweenness and flow share. In response to these findings, the second specification introduced more concentrated prior distributions for selected intercepts and factor loadings. Specifically, the standard deviations of the logit-intercept priors for the small-valued continuous responses were reduced from 1.5 to 0.7, while the PageRank loading scale and the loading standard deviations for betweenness and flow share were reduced from 1 to 0.5. For the boundary components, intercept prior standard deviations were reduced from 1.5 to 1, and boundary-loading standard deviations from 1 to 0.5. The prior centres were retained, as were the original priors for the terminal HHI and dependence interior components, regression coefficients, and binary-response parameters. The third specification retained the modifications introduced in the concentrated one and additionally introduced response-specific priors for the Beta precision parameters.

The corresponding prior medians were set to $(\tilde{\phi}_1, \tilde{\phi}_2, \tilde{\phi}_3) = (300, 10, 10)$ for terminal

PageRank, HHI, and dependence, respectively, and to $(\tilde{\phi}_1, \tilde{\phi}_2, \tilde{\phi}_3) = (500, 100, 100)$ for chokepoint PageRank, betweenness, and flow share, respectively. For each precision prior, the log-scale standard deviation was fixed at 0.7.

### 6.4.2 Prior predictive assessment

Prior predictive simulations were conducted under the rank-one model to examine the response distributions implied by the three prior specifications before conditioning on the observed responses. The model structure was held fixed across specifications, allowing the analysis to isolate the influence of the prior distributions. Table 46 compares the observed standard deviations with the mean standard deviations across prior predictive replications under each specification.

Table 46: Observed and mean prior predictive standard deviations under the three prior specifications.

| Node type | Response | Observed | ORG | CON | RS |
|---|---|---|---|---|---|
| Terminal | PageRank | 0.0133 | 0.1065 | 0.0584 | 0.0289 |
| | Counterparty HHI | 0.3254 | 0.2880 | 0.2935 | 0.2935 |
| | Role-specific dependence | 0.3219 | 0.2854 | 0.2867 | 0.2867 |
| Chokepoint | PageRank | 0.0069 | 0.0959 | 0.0542 | 0.0228 |
| | Betweenness | 0.0135 | 0.0599 | 0.0329 | 0.0154 |
| | Flow share | 0.0201 | 0.0870 | 0.0546 | 0.0278 |

For both node types, the original specification (ORG) generated substantially greater PageRank dispersion than those observed. The concentrated specification (CON) reduced this discrepancy, while the response-specific precision specification (RS) produced further improvements. For chokepoints, similar reductions were observed for betweenness and flow share, with RS yielding prior predictive standard deviations closer to their empirical counterparts.

For terminals, the prior predictive standard deviations of counterparty HHI and role-specific dependence remained relatively stable across specifications and were slightly below their observed values. This is consistent with the unchanged response-specific precision priors for HHI and dependence between CON and RS.

However, the prior predictive distributions continued to assign excessive probability to the boundary value of one. Under RS, the mean predicted proportions were approximately 0.492 for HHI and 0.504 for dependence, compared with observed proportions of 0.341 and 0.208, respectively. These discrepancies remained largely unchanged across the three specifications, indicating that the alternative priors did not adequately address the excessive prior predictive boundary mass. The corresponding results are reported in Appendix F.1.

### 6.4.3 Posterior sensitivity and predictive robustness

To assess the sensitivity of posterior inference to the alternative prior assumptions, the rank-one models were re-estimated under the response-specific precision specification (RS) and compared with the original specification (ORG).

The alternative terminal model produced no divergent transitions, a maximum reported $\widehat{R}$ of 1.01, a minimum bulk effective sample size (ESS) of 782, and a minimum tail ESS of 2,037. The corresponding values under ORG were 1.01, 817, and 1,958, respectively.

For chokepoints, neither specification produced divergent transitions, and the maximum reported $\widehat{R}$ was 1.01 in both cases. However, the minimum bulk ESS decreased from 472 under ORG to 359 under RS, while the minimum tail ESS decreased from 905 to 780. The lowest bulk ESS under both specifications was associated with the betweenness precision parameter. Under RS, this value fell below the commonly recommended minimum of 400 effective samples, indicating reduced Monte Carlo precision for this parameter. The complete diagnostic comparison, including the parameters associated with the reported extrema, is provided in Table F.2 of Appendix F.1.

Table 47 compares the posterior means and 95% highest-density intervals (HDIs) of the regression coefficients under the original and response-specific prior specifications, showing that the regression coefficients remained relatively stable, with similar posterior means and credible intervals.

Table 47: Posterior means and 95% highest-density intervals (HDIs) for regression coefficients under the original (ORG) and response-specific precision (RS) prior specifications.

| Covariate | ORG | RS |
|---|---|---|
| *Terminals* | | |
| Lagged neighbour activity | 0.096 [0.057, 0.137] | 0.093 [0.055, 0.132] |
| Log lagged voyage distance | 0.062 [0.020, 0.105] | 0.058 [0.015, 0.099] |
| Month sine | -0.032 [-0.070, 0.004] | -0.036 [-0.071, 0.003] |
| Month cosine | -0.004 [-0.042, 0.032] | -0.005 [-0.042, 0.032] |
| Log capacity | 0.409 [0.363, 0.453] | 0.408 [0.363, 0.453] |
| Terminal age | -0.019 [-0.057, 0.021] | -0.022 [-0.061, 0.019] |
| Lagged exporter indicator | 1.392 [1.296, 1.488] | 1.407 [1.311, 1.504] |
| *Chokepoints* | | |
| Lagged neighbour activity | 0.103 [0.042, 0.170] | 0.103 [0.040, 0.168] |
| Log lagged voyage distance | 0.065 [0.005, 0.128] | 0.066 [0.007, 0.131] |
| Month sine | -0.036 [-0.098, 0.023] | -0.035 [-0.098, 0.028] |
| Month cosine | -0.064 [-0.128, -0.004] | -0.065 [-0.125, -0.004] |

For terminals, the positive associations with lagged neighbour activity, voyage distance, capacity and lagged exporter status were preserved, while the intervals for terminal age and the seasonal terms continued to include zero. For chokepoints, the positive associations with lagged neighbour activity and voyage distance were also preserved, and the cosine coefficient remained negative, with its interval just excluding zero.

Greater sensitivity emerged for selected response-specific parameters. In particular, the terminal loading on the probability of HHI equal to one changed from $-3.982$ to $-3.116$, while the chokepoint loading on the probability of zero betweenness changed from $-1.659$ to $-0.980$. Both retained their negative directions, although their magnitudes changed.

In-sample posterior predictive checks were also compared to examine whether the alternative specification improved the posterior estimation of selected observed statistics. Table 48 reports the corresponding replicated statistics under both specifications.

The findings show that the alternative specification produced little change in the

Table 48: Selected posterior predictive statistics under the original (ORG) and response-specific precision (RS) prior specifications. Replicated values are posterior predictive means, except for correlations, for which medians are reported. Checks were based on 200 replicated datasets.

| Model | Statistic | Observed | ORG | RS |
|---|---|---|---|---|
| Terminal | Articulation–bridge correlation | 0.290 | -0.122 | -0.121 |
| Terminal | Exporter HHI = 1 (%) | 18.00 | 14.89 | 14.92 |
| Terminal | Exporter dependence = 1 (%) | 9.04 | 14.23 | 14.31 |
| Chokepoint | Flow-share standard deviation | 0.0201 | 0.0182 | 0.0182 |
| Chokepoint | Flow-share median | 0.0159 | 0.0190 | 0.0190 |

reported posterior predictive statistics, leaving the main discrepancies largely unresolved. For terminals, the replicated association between articulation-point and incident-global-bridge status remained negative, despite its positive observed correlation, and the boundary frequencies of HHI and dependence among exporters remained inaccurately reproduced; for chokepoints, both specifications underestimated the observed dispersion of flow share and overestimated its median.

Given the limited changes in posterior inference and predictive performance, together with the minimum bulk ESS falling below the recommended threshold in the alternative chokepoint fit, the original specification (ORG) was retained for the main analysis. The response-specific precision specification (RS) therefore serves as a sensitivity analysis, indicating that the main posterior findings are reasonably robust under the alternative prior specifications considered, while the identified predictive limitations remain largely unresolved.

## 6.5 Comparative discussion and limitations

### 6.5.1 Model limitations and implications

The bounded rank-one models provide an initial Bayesian framework for examining common patterns across multiple measures of node importance in the LNG maritime network. Both models accommodate the bounded nature of the responses and their observed boundary values, while the posterior estimates satisfy the MCMC diagnostic

screening criteria considered.

The terminal latent factor is associated with network centrality and counterparty diversification, whereas the chokepoint factor reflects centrality, flow exposure and structural position. These interpretations suggest that a shared latent dimension can summarise several aspects of network position, although its meaning differs between the two node types.

Nevertheless, satisfactory sampling diagnostics do not guarantee an adequate representation of the observed response distributions. The posterior predictive checks identified limitations in the ability of the terminal model to reproduce role-specific distributions and associations between structural indicators. For chokepoints, the main discrepancies concern the distribution of flow share and its relationships with other centrality measures.

The prior sensitivity analysis further showed that, although the main regression coefficients were relatively stable, some response-specific parameters were more sensitive to prior choices. Moreover, the alternative specification did not substantially improve the posterior predictive results, highlighting that the limitations identified remain relevant under the considered prior modifications.

These findings motivate several possible extensions. For terminals, allowing response-specific intercepts or factor loadings to vary by terminal role could help capture differences between importers and exporters that are not fully represented by the current model. This distinction may be particularly relevant for counterparty concentration: for importing terminals, a high HHI indicates dependence on a limited number of LNG suppliers, potentially increasing vulnerability to supply disruptions; for exporting terminals, by contrast, a low HHI indicates a more diversified set of destinations, potentially reflecting broader commercial reach within international LNG trade. The current specification imposes a common relationship between HHI and the latent factor across terminal roles, potentially overlooking such differences.

Another possible extension would be to relax the assumption of conditional independence between responses given the latent factor. This could allow the model to

capture residual associations that are not adequately represented by the shared factor alone.

More flexible likelihood specifications could also improve the representation of bounded responses, especially where the observed dispersion or distributional shape is not adequately reproduced, as in the case of chokepoint flow share.

Finally, node-specific random effects could account for persistent heterogeneity across terminals or chokepoints and introduce dependence among repeated observations of the same node. A further extension would be to develop a joint model for terminals and chokepoints, which are currently analysed separately despite belonging to the same maritime network. Such a framework could allow for dependencies between the two node types while retaining response specifications appropriate to their different characteristics.

These extensions are not mutually exclusive, and the posterior predictive discrepancies do not establish which modelling assumptions are primarily responsible for the observed shortcomings. Their adequacy would therefore need to be evaluated through further model comparisons and predictive assessments.

An additional limitation concerns the estimation sample. The models are fitted only to node-month observations corresponding to consecutive active months. Consequently, the latent factors describe network position conditional on sustained observed activity, without modelling the process determining whether a node is active. Incorporating an activity component would extend the analysis to both network participation and node importance conditional on participation.

Taken together, these limitations suggest that the estimated latent factors should be interpreted as preliminary summaries of network position rather than comprehensive measures of node criticality. The present models therefore provide a baseline for subsequent methodological developments.

# Chapter 7

# Conclusions and future work

# Appendix A

# Chokepoint geometry registry

Table A.1 documents the geometries used to assess route crossings for all 28 initial PortWatch chokepoints. Published spatial geometries were used directly where available. Geometries based on published routeing coordinates were reconstructed as polygons. For the remaining passages, operational corridors or crossing gates were constructed using the geographic and navigational references indicated below. These constructed geometries serve to classify route crossings and do not represent official maritime boundaries. Four of the 28 passages were not traversed under the final route-assignment specification and were excluded from the monthly network.

Table A.1: Geometry registry for the 28 initial chokepoints

| ID | Chokepoint | Geometry source | Geometry used for route classification |
|---|---|---|---|
| CP001 | Suez Canal | World Bank | Published polygon. |
| CP002 | Panama Canal | Marine Regions reference | Operational canal centreline with an 8 km buffer. |
| CP003 | Bosporus Strait* | Marine Regions reference | Operational strait centreline with a 5 km buffer. |
| CP004 | Bab el-Mandeb Strait | World Bank | Published polygon. |
| CP005 | Malacca Strait | Marine Regions, IHO | Detailed source MultiPolygon. |
| CP006 | Strait of Hormuz | Marine Regions, SeaVoX | Detailed source MultiPolygon. |
| CP007 | Cape of Good Hope | World Bank | Published polygon. |
| CP008 | Gibraltar Strait | Marine Regions, IHO | Detailed source MultiPolygon. |
| CP009 | Dover Strait | Marine Regions, SeaVoX | Detailed source MultiPolygon. |

| ID | Chokepoint | Geometry source | Geometry used for route classification |
|---|---|---|---|
| CP010 | Oresund Strait | Marine Regions reference | Crossing gate, 45 × 8 km, oriented east–west. |
| CP011 | Taiwan Strait | Marine Regions reference | Crossing gate, 220 × 12 km, oriented northwest–southeast. |
| CP012 | Korea Strait | Marine Regions reference | Crossing gate, 130 × 10 km, oriented northwest–southeast. |
| CP013 | Tsugaru Strait | Marine Regions reference | Crossing gate, 55 × 8 km, oriented north–south and centred on the Marine Regions location. |
| CP014 | Luzon Strait | Marine Regions reference | Crossing gate, 220 × 12 km, oriented north–south. |
| CP015 | Lombok Strait | IMO routeing coordinates | Polygon reconstructed from three traffic-separation-scheme sections. |
| CP016 | Ombai Strait | IMO sea-lane reference | Crossing gate, 115 × 12 km, oriented north–south. |
| CP017 | Bohai Strait | NGA GeoNames reference | Crossing gate, 120 × 15 km, oriented north–south. |
| CP018 | Torres Strait | IMO routeing coordinates | Polygon reconstructed from the Prince of Wales Channel two-way route. |
| CP019 | Sunda Strait | IMO routeing coordinates | Polygon reconstructed from two traffic-separation-scheme sections. |
| CP020 | Makassar Strait | Marine Regions, IHO | Detailed source MultiPolygon. |
| CP021 | Magellan Strait | Marine Regions reference | Crossing gate, 55 × 8 km, oriented north–south. |
| CP022 | Yucatan Channel | NOAA/NCEI reference | Crossing gate, 230 × 15 km, oriented east–west. |
| CP023 | Windward Passage | Marine Regions reference | Crossing gate, 105 × 12 km, oriented northwest–southeast. |
| CP024 | Mona Passage | Marine Regions reference | Crossing gate, 145 × 12 km, oriented east–west. |
| CP025 | Balabac Strait* | Marine Regions reference | Crossing gate, 100 × 10 km, oriented north–south. |
| CP026 | Bering Strait* | IMO routeing coordinates | Polygon reconstructed from the two-way route through the strait. |
| CP027 | Mindoro Strait | Marine Regions reference | Crossing gate, 95 × 10 km, oriented northeast–southwest. |

| | | | |
|---|---|---|---|
| CP028 | Kerch Strait* | Marine Regions reference | Crossing gate, 26 × 5 km, oriented east–west. |

*Notes:* IDs correspond to the study registry; the original PortWatch identifiers and reference coordinates are retained in the accompanying digital file. Gate dimensions are span × thickness. Crossing gates are centred on the PortWatch reference point. The asterisk identifies passages excluded from the final monthly network because no reconstructed route crossed them under the selected 25 km tolerance. The complete geometry registry and the GeoJSON file containing the final geometries are available in the project's GitHub repository.

# Appendix B

# Sensitivity analysis of chokepoint assignments

## B.1 Purpose and method

The identification of the chokepoints traversed by each reconstructed maritime route may be affected by small spatial discrepancies between the `SeaRoute` paths and the operational chokepoint geometries. Such discrepancies can arise from the spatial resolution and simplification of the routing network, the geographic representation adopted for each passage, and small differences between a reconstructed route centreline and the corresponding transit geometry.

A sensitivity analysis was therefore conducted to quantify how the route–chokepoint assignments changed under alternative spatial specifications. Four classifications were compared: exact geometric intersection and intersections obtained after applying uniform buffers of 5, 10, and 25 km around each of the 28 candidate chokepoint geometries. The 25-km specification corresponds to the classification retained for the final network.

For each chokepoint, distances and buffers were calculated in a local azimuthal-equidistant projection centred on the geometry, allowing the tolerance distance to be measured in metres rather than in geographic degrees. Each of the 1,037 reconstructed origin–destination routes was then compared with the original and buffered chokepoint geometries. For every route–chokepoint pair, the analysis recorded the minimum geometric distance, exact-intersection status, and whether the route fell within each alternative tolerance. The same tolerance was applied to every candidate chokepoint, thereby avoiding passage-specific adjustments based on the observed routes.

## B.2 Sensitivity results

Table B.1 reports, for each candidate chokepoint, the number of reconstructed routes identified through exact intersection and under the three alternative tolerance buffers. The final column gives the number of assignments added by the 25-km specification relative to exact intersection.

Table B.1: Sensitivity of route–chokepoint assignments to alternative spatial tolerances

| **Chokepoint** | **Exact** | **5 km** | **10 km** | **25 km** | **Additional at 25 km** |
|---|---|---|---|---|---|
| Suez Canal | 151 | 151 | 151 | 151 | 0 |
| Panama Canal | 148 | 148 | 148 | 148 | 0 |
| Bosporus Strait | 0 | 0 | 0 | 0 | 0 |
| Bab el-Mandeb Strait | 147 | 147 | 147 | 147 | 0 |
| Malacca Strait | 133 | 133 | 133 | 133 | 0 |
| Strait of Hormuz | 122 | 122 | 122 | 122 | 0 |
| Cape of Good Hope | 65 | 65 | 65 | 65 | 0 |
| Gibraltar Strait | 216 | 222 | 222 | 222 | 6 |
| Dover Strait | 91 | 91 | 91 | 91 | 0 |
| Oresund Strait | 47 | 47 | 47 | 47 | 0 |
| Taiwan Strait | 152 | 152 | 152 | 152 | 0 |
| Korea Strait | 3 | 3 | 3 | 68 | 65 |
| Tsugaru Strait | 50 | 50 | 52 | 52 | 2 |
| Luzon Strait | 43 | 56 | 56 | 56 | 13 |
| Lombok Strait | 69 | 69 | 69 | 74 | 5 |
| Ombai Strait | 57 | 57 | 57 | 58 | 1 |
| Bohai Strait | 35 | 35 | 35 | 35 | 0 |
| Torres Strait | 5 | 5 | 5 | 5 | 0 |
| Sunda Strait | 51 | 51 | 51 | 51 | 0 |
| Makassar Strait | 58 | 58 | 58 | 58 | 0 |
| Magellan Strait | 2 | 2 | 2 | 2 | 0 |
| Yucatan Channel | 137 | 137 | 137 | 137 | 0 |
| Windward Passage | 10 | 10 | 10 | 10 | 0 |
| Mona Passage | 17 | 17 | 17 | 17 | 0 |
| Balabac Strait | 0 | 0 | 0 | 0 | 0 |
| Bering Strait | 0 | 0 | 0 | 0 | 0 |
| Mindoro Strait | 0 | 0 | 8 | 63 | 63 |
| Kerch Strait | 0 | 0 | 0 | 0 | 0 |
| **Total assignments** | **1,809** | **1,828** | **1,838** | **1,964** | **155** |
| **Observed chokepoints** | **23** | **23** | **24** | **24** | — |

Exact intersection identified 1,809 route–chokepoint assignments involving 23 of the 28 candidate passages. Under the final 25-km specification, the number of assignments increased to 1,964 and the number of identified chokepoints increased to 24, resulting in 155 additional assignments, or an 8.6% increase relative to exact intersection.

The assignments were stable across all four specifications for 17 of the 24 chokepoints retained in the final network. Most of the additional assignments under the final specification were concentrated in Korea Strait and Mindoro Strait, indicating that the results for these two passages were particularly sensitive to the selected spatial tolerance.

Four candidate chokepoints—Bosporus Strait, Balabac Strait, Bering Strait, and Kerch Strait—were not traversed under any of the evaluated specifications: their minimum distances from the closest reconstructed routes were approximately 71.3, 162.0, 2,617.3, and 829.7 km, respectively. They were therefore excluded from the final monthly network. The exclusion indicates that none of the reconstructed LNG routes in the study period passed within 25 km of their operational geometries, but it does not imply that the passages have no relevance to maritime traffic outside the analysed LNG network.

The 25-km tolerance was retained because it accommodates spatial misalignments between the simplified `SeaRoute` paths and the operational transit geometries while applying a uniform and transparent classification rule to all candidate passages. The exact-intersection assignments were preserved separately for reproducibility, and the sensitivity procedure did not alter the reconstructed route geometries.

# Appendix C

# Model-ready panels variables

This appendix documents the variables contained in the three model-ready panels described in Section 3.5. Variables are grouped according to their analytical role rather than their physical order in the CSV files. Flow variables are measured monthly, binary indicators take the value one when the stated condition is satisfied, and variables carrying the suffix `lag1` refer to month $t-1$. Log-transformed variables were calculated using $\log(1+x)$.

The three panels cover January 2020–December 2024. Consequently, all one-month lagged variables are unavailable in January 2020. Measures based on historical neighbours are additionally unavailable until a node has acquired at least one neighbour through a connection observed before the current month. Commercial criticality measures that require realised trade are defined only for active terminal–months.

## C.1 Node activity panel

Table C.1: Variables in the node activity panel

| Variable | Definition | Role |
|---|---|---|
| `node_id` | Unique node identifier. | Identifier |
| `period_month` | Calendar month of observation. | Identifier |
| `node_name` | Node name. | Identifier |
| `node_type` | Node category: terminal or chokepoint. | Covariate |
| `country` | Country in which a terminal is located; not applicable to chokepoints. | Covariate |
| `active` | Equals one when the node participates in the monthly network and zero otherwise. | Outcome |
| `incoming_flow` | Total LNG volume entering the node in month (t). | Descriptive |
| `outgoing_flow` | Total LNG volume leaving the node in month (t). | Descriptive |

Continued on next page

Table C.1 continued

| Variable | Definition | Role |
|---|---|---|
| incident_flow | Sum of incoming and outgoing LNG volume incident to the node. | Descriptive |
| node_throughput | Node throughput, adjusted for double counting at chokepoints. | Descriptive |
| in_degree | Number of distinct incoming neighbours in the monthly directed network. | Descriptive |
| out_degree | Number of distinct outgoing neighbours in the monthly directed network. | Descriptive |
| total_degree | Sum of monthly in-degree and out-degree. | Descriptive |
| in_voyages | Number of incoming voyage movements associated with the node. | Descriptive |
| out_voyages | Number of outgoing voyage movements associated with the node. | Descriptive |
| incident_voyage_traversals | Sum of incoming and outgoing voyage traversals incident to the node. | Descriptive |
| total_voyages | Number of voyages associated with the node, adjusted for double counting at chokepoints. | Descriptive |
| in_route_exposure | Number of reconstructed routes entering the node. | Descriptive |
| out_route_exposure | Number of reconstructed routes leaving the node. | Descriptive |
| total_route_exposure | Total reconstructed-route exposure of the node. | Descriptive |
| betweenness | Normalised weighted betweenness centrality in the monthly segment-level network. | Descriptive |
| pagerank | Weighted PageRank in the monthly directed segment-level network. | Descriptive |
| share_monthly_network_flow | Share of total monthly node throughput associated with the node. | Descriptive |
| months_since_last_active | Number of months elapsed since the node was last active; unavailable before its first observed active month. | Supplementary |
| infrastructure_type | Terminal infrastructure role, classified as import or export; not applicable to chokepoints. | Covariate |
| capacity_mtpa | Annual terminal processing capacity in million tonnes per annum; not applicable to chokepoints. | Covariate |
| log1p_capacity_mtpa | Log-transformed annual terminal processing capacity. | Covariate |
| start_year | Year in which the terminal began operating; not applicable to chokepoints. | Covariate |
| activity_lag1 | Activity status in the preceding month. | Covariate |

Continued on next page

Table C.1 continued

| Variable | Definition | Role |
|---|---|---|
| throughput_lag1 | Node throughput in the preceding month. | Covariate |
| log1p_throughput_lag1 | Log-transformed node throughput in the preceding month. | Covariate |
| voyage_count_lag1 | Number of voyages associated with the node in the preceding month. | Covariate |
| log1p_voyage_count_lag1 | Log-transformed voyage count in the preceding month. | Covariate |
| historical_neighbor_count_lag1 | Number of neighbours connected to the node by at least one edge observed up to (t-1). | Covariate |
| active_neighbor_count_lag1 | Number of historical neighbours active in (t-1). | Covariate |
| neighbor_activity_lag1 | Share of historical neighbours active in (t-1). | Covariate |
| lag_available | Equals one when preceding-month values can be calculated and zero in January 2020. | Availability |
| year | Calendar year. | Time index |
| month | Calendar month, coded from 1 to 12. | Time index |
| month_sin | Sine transformation of the calendar month. | Covariate |
| month_cos | Cosine transformation of the calendar month. | Covariate |

## C.2 Terminal criticality panel

Table C.2: Variables in the terminal criticality panel

| Variable | Definition | Role |
|---|---|---|
| terminal_id | Unique terminal identifier. | Identifier |
| node_name | Terminal name. | Identifier |
| period_month | Calendar month of observation. | Identifier |
| year | Calendar year. | Time index |
| month | Calendar month, coded from 1 to 12. | Time index |
| node_type | Node category, equal to terminal throughout this panel. | Descriptor |
| country | Country in which the terminal is located. | Covariate |
| infrastructure_type | Terminal infrastructure role, classified as import or export. | Covariate |
| capacity_mtpa | Annual processing capacity in million tonnes per annum. | Covariate |
| log1p_capacity_mtpa | Log-transformed annual processing capacity. | Covariate |

Continued on next page

Table C.2 continued

| Variable | Definition | Role |
|---|---|---|
| unit_count | Number of infrastructure units associated with the terminal. | Covariate |
| start_year | Year in which the terminal began operating. | Covariate |
| terminal_age | Difference between the observation year and the terminal start year; unavailable before the start year. | Covariate |
| terminal_role | Observed monthly role: exporter, importer, or inactive. | Descriptor |
| active | Equals one when the terminal participates in LNG trade in month (t). | Conditioning indicator |
| export_share | Share of terminal throughput represented by outgoing flow; defined for active terminal–months. | Descriptor |
| outgoing_flow | Total LNG volume exported by the terminal. | Descriptive |
| incoming_flow | Total LNG volume imported by the terminal. | Descriptive |
| throughput | Sum of outgoing and incoming LNG volume. | Descriptive |
| outgoing_voyages | Number of voyages departing from the terminal. | Descriptive |
| incoming_voyages | Number of voyages arriving at the terminal. | Descriptive |
| voyage_count | Total number of voyages associated with the terminal. | Descriptive |
| counterparty_count | Number of distinct terminal counterparties. | Descriptive |
| counterparty_country_count | Number of distinct counterparty countries. | Descriptive |
| counterparty_hhi_terminal | Role-specific HHI of flows across terminal counterparties. | Outcome |
| counterparty_hhi_country | Role-specific HHI of flows across counterparty countries. | Outcome |
| effective_counterparties_terminal | Effective number of terminal counterparties, calculated as the inverse of the terminal-level HHI. | Supplementary |
| effective_counterparties_country | Effective number of counterparty countries, calculated as the inverse of the country-level HHI. | Supplementary |
| max_dependence_generated | Maximum recipient dependence generated by an exporting terminal. | Supplementary |
| weighted_dependence_generated | Exporter-generated dependence weighted by the distribution of its outgoing flow. | Supplementary |
| country_terminal_dependence | Importing terminal's share of total LNG imports received by its country. | Supplementary |
| role_specific_dependence | Dependence measure selected according to the terminal's observed monthly role. | Outcome |

Continued on next page

Table C.2 continued

| Variable | Definition | Role |
|---|---|---|
| import_pagerank | Weighted PageRank calculated in the observed direction of LNG flows. | Supplementary |
| export_pagerank | Weighted PageRank calculated after reversing the direction of LNG flows. | Supplementary |
| role_specific_pagerank | Directional PageRank selected according to the terminal's observed monthly role. | Outcome |
| degree_undirected | Degree in the undirected projection of the active monthly network. | Structural |
| is_articulation_point | Equals one when removal of the terminal increases the number of connected components. | Outcome |
| global_bridge_count | Number of global bridges incident to the terminal. | Structural |
| global_bridge_share | Share of incident undirected edges classified as global bridges. | Structural |
| local_bridge_only_count | Number of incident local bridges after excluding global bridges. | Structural |
| local_bridge_only_share | Share of incident undirected edges classified as local-only bridges. | Structural |
| has_incident_global_bridge | Equals one when the terminal is incident to at least one global bridge. | Outcome |
| connected_to_articulation_point | Equals one when the terminal is directly connected to an articulation point. | Outcome |
| terminal_role_lag1 | Observed terminal role in the preceding month: exporter, importer, or inactive. | Covariate |
| activity_lag1 | Terminal activity status in the preceding month. | Covariate |
| throughput_lag1 | Terminal throughput in the preceding month. | Covariate |
| log1p_throughput_lag1 | Log-transformed terminal throughput in the preceding month. | Covariate |
| voyage_count_lag1 | Terminal voyage count in the preceding month. | Covariate |
| log1p_voyage_count_lag1 | Log-transformed terminal voyage count in the preceding month. | Covariate |
| mean_voyage_distance_lag1 | Role-relevant mean voyage distance in the preceding month; undefined when no applicable voyage was observed. | Covariate |
| log1p_mean_voyage_distance_lag1 | Log-transformed preceding-month role-relevant mean voyage distance; undefined when no applicable voyage was observed. | Covariate |
| historical_neighbor_count_lag1 | Number of historical neighbours observed up to (t-1). | Covariate |
| active_neighbor_count_lag1 | Number of historical neighbours active in (t-1). | Covariate |

Continued on next page

Table C.2 continued

| Variable | Definition | Role |
|---|---|---|
| lag_available | Equals one when the preceding calendar month is available and zero for January 2020. | Availability |
| neighbor_activity_lag1 | Share of historical neighbours active in (t-1). | Covariate |
| month_sin | Sine transformation of the calendar month. | Covariate |
| month_cos | Cosine transformation of the calendar month. | Covariate |

## C.3 Chokepoint criticality panel

Table C.3: Variables in the chokepoint criticality panel

| Variable | Definition | Role |
|---|---|---|
| node_id | Unique chokepoint identifier. | Identifier |
| period_month | Calendar month of observation. | Identifier |
| node_name | Chokepoint name. | Identifier |
| node_type | Node category, equal to chokepoint throughout this panel. | Descriptor |
| active | Equals one when the chokepoint is traversed in month (t). | Conditioning indicator |
| incoming_flow | Total LNG volume entering the chokepoint. | Descriptive |
| outgoing_flow | Total LNG volume leaving the chokepoint. | Descriptive |
| incident_flow | Sum of incoming and outgoing LNG volume incident to the chokepoint. | Descriptive |
| node_throughput | Chokepoint throughput, equal to one half of incident flow. | Descriptive |
| in_degree | Number of distinct incoming neighbours. | Descriptive |
| out_degree | Number of distinct outgoing neighbours. | Descriptive |
| total_degree | Sum of directed in-degree and out-degree. | Descriptive |
| in_voyages | Number of incoming voyage traversals. | Descriptive |
| out_voyages | Number of outgoing voyage traversals. | Descriptive |
| incident_voyage_traversals | Sum of incoming and outgoing voyage traversals. | Descriptive |
| total_voyages | Chokepoint voyage count adjusted for double counting across incoming and outgoing segments. | Descriptive |
| in_route_exposure | Number of reconstructed routes entering the chokepoint. | Descriptive |
| out_route_exposure | Number of reconstructed routes leaving the chokepoint. | Descriptive |

Continued on next page

Table C.3 continued

| Variable | Definition | Role |
|---|---|---|
| total_route_exposure | Total reconstructed-route exposure of the chokepoint. | Descriptive |
| betweenness | Normalised weighted betweenness centrality. | Outcome |
| pagerank | Weighted PageRank in the monthly directed segment-level network. | Outcome |
| share_monthly_network_flow | Share of total monthly node throughput associated with the chokepoint. | Outcome |
| months_since_last_active | Number of months since the chokepoint was last active; unavailable before its first active month. | Supplementary |
| degree_undirected | Degree in the undirected projection of the active monthly network. | Structural |
| is_articulation_point | Equals one when removal of the chokepoint increases the number of connected components. | Outcome |
| global_bridge_count | Number of global bridges incident to the chokepoint. | Structural |
| global_bridge_share | Share of incident undirected edges classified as global bridges. | Structural |
| local_bridge_only_count | Number of incident local bridges after excluding global bridges. | Structural |
| local_bridge_only_share | Share of incident undirected edges classified as local-only bridges. | Structural |
| has_incident_global_bridge | Equals one when the chokepoint is incident to at least one global bridge. | Outcome |
| connected_to_articulation_point | Equals one when the chokepoint is directly connected to an articulation point. | Outcome |
| activity_lag1 | Chokepoint activity status in the preceding month. | Covariate |
| throughput_lag1 | Chokepoint throughput in the preceding month. | Covariate |
| log1p_throughput_lag1 | Log-transformed chokepoint throughput in the preceding month. | Covariate |
| voyage_count_lag1 | Chokepoint voyage count in the preceding month. | Covariate |
| log1p_voyage_count_lag1 | Log-transformed chokepoint voyage count in the preceding month. | Covariate |
| mean_voyage_distance_lag1 | Mean voyage distance for voyages traversing the chokepoint in the preceding month; undefined when no applicable voyage was observed. | Covariate |
| log1p_mean_voyage_distance_lag1 | Log-transformed preceding-month mean voyage distance. | Covariate |

Continued on next page

Table C.3 continued

| Variable | Definition | Role |
|---|---|---|
| historical_neighbor_count_lag1 | Number of historical neighbours observed up to (t-1). | Covariate |
| active_neighbor_count_lag1 | Number of historical neighbours active in (t-1). | Covariate |
| neighbor_activity_lag1 | Share of historical neighbours active in (t-1). | Covariate |
| lag_available | Equals one when preceding-month information can be calculated. | Availability |
| year | Calendar year. | Time index |
| month | Calendar month, coded from 1 to 12. | Time index |
| month_sin | Sine transformation of the calendar month. | Covariate |
| month_cos | Cosine transformation of the calendar month. | Covariate |

# Appendix D

# Full conditional posterior distributions

This appendix derives the full conditional posterior distributions for the rank-one baseline model. The derivations apply separately to terminals and chokepoints, using the response likelihoods and original prior specification introduced in Sections 5.1.2 and 5.1.3. Each full conditional is obtained from the joint posterior in Equation (43) by retaining only the terms that depend on the parameter of interest. Multiplicative terms that are constant with respect to that quantity are absorbed into the normalising constant.

The notation follows the main text: $n \in \mathcal{O}$ indexes the node–month observations, $N = |\mathcal{O}|$, and $\mathcal{I}_j = \{n \in \mathcal{O} : 0 < Y_{nj} < 1\}$ contains the interior observations for bounded response $j$. Throughout, *rest* denotes all remaining unknown quantities together with the observed data and covariates.

## D.1 Baseline rank-one specification

### D.1.1 Factor-regression coefficients

Let $\mathbf{X}$ be the $N \times q$ covariate design matrix, with row $\mathbf{x}_n^\top$, and let $\mathbf{f}$ collect the corresponding latent factors.

Conditional on $\mathbf{f}$, the response likelihood does not depend on $\boldsymbol{\beta}$. Combining the Gaussian factor-regression density with the independent $\mathcal{N}(0, 0.5^2)$ coefficient priors therefore gives

$$p(\boldsymbol{\beta} \mid \text{rest}) \propto p(\mathbf{f} \mid \boldsymbol{\beta}, \mathbf{X}) p(\boldsymbol{\beta}).$$

Writing the relevant densities explicitly,

$$p(\boldsymbol{\beta} \mid \text{rest}) \propto \left[ \prod_{n \in \mathcal{O}} \exp\left\{ -\frac{1}{2} (f_n - \mathbf{x}_n^\top \boldsymbol{\beta})^2 \right\} \right] \left[ \prod_{r=1}^{q} \exp\left\{ -\frac{\beta_r^2}{2(0.5)^2} \right\} \right].$$

Combining the exponential terms and using matrix notation yields

$$p(\boldsymbol{\beta} \mid \text{rest}) \propto \exp\left\{ -\frac{1}{2}\Big[(\mathbf{f}-\mathbf{X}\boldsymbol{\beta})^\mathsf{T}(\mathbf{f}-\mathbf{X}\boldsymbol{\beta}) + 4\boldsymbol{\beta}^\mathsf{T}\boldsymbol{\beta}\Big]\right\}.$$

The first quadratic term expands as

$$(\mathbf{f}-\mathbf{X}\boldsymbol{\beta})^\mathsf{T}(\mathbf{f}-\mathbf{X}\boldsymbol{\beta}) = \mathbf{f}^\mathsf{T}\mathbf{f} - 2\boldsymbol{\beta}^\mathsf{T}\mathbf{X}^\mathsf{T}\mathbf{f} + \boldsymbol{\beta}^\mathsf{T}\mathbf{X}^\mathsf{T}\mathbf{X}\boldsymbol{\beta}.$$

Removing $\mathbf{f}^\mathsf{T}\mathbf{f}$, which does not depend on $\boldsymbol{\beta}$, gives

$$p(\boldsymbol{\beta} \mid \text{rest}) \propto \exp\left\{ -\frac{1}{2}\Big[\boldsymbol{\beta}^\mathsf{T}(\mathbf{X}^\mathsf{T}\mathbf{X} + 4\mathbf{I}_q)\boldsymbol{\beta} - 2\boldsymbol{\beta}^\mathsf{T}\mathbf{X}^\mathsf{T}\mathbf{f}\Big]\right\},$$

where $\mathbf{I}_q$ is the $q \times q$ identity matrix.

Define

$$\mathbf{V}_\beta = (\mathbf{X}^\mathsf{T}\mathbf{X} + 4\mathbf{I}_q)^{-1}, \qquad \mathbf{m}_\beta = \mathbf{V}_\beta\mathbf{X}^\mathsf{T}\mathbf{f}.$$

Since $\mathbf{V}_\beta^{-1}\mathbf{m}_\beta = \mathbf{X}^\mathsf{T}\mathbf{f}$, completing the square gives

$$\boldsymbol{\beta}^\mathsf{T}\mathbf{V}_\beta^{-1}\boldsymbol{\beta} - 2\boldsymbol{\beta}^\mathsf{T}\mathbf{X}^\mathsf{T}\mathbf{f} = (\boldsymbol{\beta}-\mathbf{m}_\beta)^\mathsf{T}\mathbf{V}_\beta^{-1}(\boldsymbol{\beta}-\mathbf{m}_\beta) - \mathbf{m}_\beta^\mathsf{T}\mathbf{V}_\beta^{-1}\mathbf{m}_\beta.$$

The final term is constant with respect to $\boldsymbol{\beta}$. Thus,

$$p(\boldsymbol{\beta} \mid \text{rest}) \propto \exp\left\{-\frac{1}{2}(\boldsymbol{\beta}-\mathbf{m}_\beta)^\mathsf{T}\mathbf{V}_\beta^{-1}(\boldsymbol{\beta}-\mathbf{m}_\beta)\right\},$$

which yields the following conjugate full conditional

$$\boldsymbol{\beta} \mid \text{rest} \sim \mathcal{N}_q\big((\mathbf{X}^\mathsf{T}\mathbf{X} + 4\mathbf{I}_q)^{-1}\mathbf{X}^\mathsf{T}\mathbf{f},\ (\mathbf{X}^\mathsf{T}\mathbf{X} + 4\mathbf{I}_q)^{-1}\big).$$

### D.1.2 Response loadings

#### Binary-response loadings

For binary response $k$,

$$B_{nk} \mid f_n, a_k, \ell_k \sim \text{Bernoulli}(p_{nk}), \qquad \text{logit}(p_{nk}) = a_k + \ell_k f_n.$$

Holding the factors and other response parameters fixed, only the likelihood of binary response $k$ and the prior of $\ell_k$ depend on this loading. Therefore,

$$p(\ell_k \mid \text{rest}) \propto \left[\prod_{n\in\mathcal{O}} p(B_{nk} \mid f_n, a_k, \ell_k)\right] p(\ell_k).$$

The Bernoulli contribution is

$$p(B_{nk} \mid f_n, a_k, \ell_k) = p_{nk}^{B_{nk}}(1 - p_{nk})^{1-B_{nk}}.$$

Using

$$p_{nk} = \frac{\exp(a_k + \ell_k f_n)}{1 + \exp(a_k + \ell_k f_n)}, \qquad 1 - p_{nk} = \frac{1}{1 + \exp(a_k + \ell_k f_n)},$$

this becomes

$$p(B_{nk} \mid f_n, a_k, \ell_k) = \frac{\exp\{B_{nk}(a_k + \ell_k f_n)\}}{1 + \exp(a_k + \ell_k f_n)}.$$

The prior $\ell_k \sim \mathcal{N}(0, 1)$ has density

$$p(\ell_k) = \frac{1}{\sqrt{2\pi}} \exp\left(-\frac{\ell_k^2}{2}\right).$$

Combining the likelihood and prior and absorbing the Normal normalising constant of the normal prior distribution gives

$$p(\ell_k \mid \text{rest}) \propto \left[\prod_{n \in \mathcal{O}} \frac{\exp\{B_{nk}(a_k + \ell_k f_n)\}}{1 + \exp(a_k + \ell_k f_n)}\right] \exp\left(-\frac{\ell_k^2}{2}\right).$$

This expression applies to $k = 1, 2, 3$ for terminals and $k = 1, 2$ for chokepoints.

**Ordinary Beta-response loadings**

For an ordinary Beta response,

$$Y_{nj} \mid f_n, \alpha_j, \lambda_j, \phi_j \sim \text{Beta}\big(\mu_{nj}\phi_j, (1 - \mu_{nj})\phi_j\big), \qquad \text{logit}(\mu_{nj}) = \alpha_j + \lambda_j f_n.$$

Its density is

$$p(Y_{nj} \mid f_n, \alpha_j, \lambda_j, \phi_j) = \frac{\Gamma(\phi_j)}{\Gamma(\mu_{nj}\phi_j)\Gamma((1 - \mu_{nj})\phi_j)} Y_{nj}^{\mu_{nj}\phi_j - 1}(1 - Y_{nj})^{(1-\mu_{nj})\phi_j - 1}.$$

The loading enters this density through $\mu_{nj}$. Retaining the relevant likelihood and prior gives

$$p(\lambda_j \mid \text{rest}) \propto \left[\prod_{n \in \mathcal{O}} p(Y_{nj} \mid f_n, \alpha_j, \lambda_j, \phi_j)\right] p(\lambda_j).$$

**Unrestricted loading for chokepoint flow share**

For chokepoint flow share, $j = 3$ and $\lambda_3 \sim \mathcal{N}(0, 1)$, with density

$$p(\lambda_3) = \frac{1}{\sqrt{2\pi}} \exp\left(-\frac{\lambda_3^2}{2}\right).$$

Substituting the Beta likelihood and Normal prior gives

$$p(\lambda_3 \mid \text{rest}) \propto \left[\prod_{n\in\mathcal{O}} \frac{\Gamma(\phi_3)Y_{n3}^{\mu_{n3}\phi_3-1}(1-Y_{n3})^{(1-\mu_{n3})\phi_3-1}}{\Gamma(\mu_{n3}\phi_3)\Gamma((1-\mu_{n3})\phi_3)}\right]\exp\left(-\frac{\lambda_3^2}{2}\right).$$

Since $\phi_3$ is held fixed, $\Gamma(\phi_3)$ does not depend on $\lambda_3$ and can be absorbed into the normalising constant. The resulting kernel is

$$p(\lambda_3 \mid \text{rest}) \propto \left[\prod_{n\in\mathcal{O}} \frac{Y_{n3}^{\mu_{n3}\phi_3-1}(1-Y_{n3})^{(1-\mu_{n3})\phi_3-1}}{\Gamma(\mu_{n3}\phi_3)\Gamma((1-\mu_{n3})\phi_3)}\right]\exp\left(-\frac{\lambda_3^2}{2}\right).$$

**Positive PageRank loading**

For PageRank in both node types, $j = 1$ and $\lambda_1 \sim \text{HalfNormal}(1)$. Its prior density is

$$p(\lambda_1) = \sqrt{\frac{2}{\pi}}\exp\left(-\frac{\lambda_1^2}{2}\right)\mathbb{1}_{\{\lambda_1>0\}}.$$

Combining this prior with the PageRank likelihood gives

$$p(\lambda_1 \mid \text{rest}) \propto \left[\prod_{n\in\mathcal{O}} \frac{\Gamma(\phi_1)Y_{n1}^{\mu_{n1}\phi_1-1}(1-Y_{n1})^{(1-\mu_{n1})\phi_1-1}}{\Gamma(\mu_{n1}\phi_1)\Gamma((1-\mu_{n1})\phi_1)}\right]\exp\left(-\frac{\lambda_1^2}{2}\right)\mathbb{1}_{\{\lambda_1>0\}}.$$

Removing the factors $\Gamma(\phi_1)$, which are constant with respect to $\lambda_1$, yields

$$p(\lambda_1 \mid \text{rest}) \propto \left[\prod_{n\in\mathcal{O}} \frac{Y_{n1}^{\mu_{n1}\phi_1-1}(1-Y_{n1})^{(1-\mu_{n1})\phi_1-1}}{\Gamma(\mu_{n1}\phi_1)\Gamma((1-\mu_{n1})\phi_1)}\right]\exp\left(-\frac{\lambda_1^2}{2}\right)\mathbb{1}_{\{\lambda_1>0\}}.$$

**Interior-mean loadings for boundary-inflated Beta responses**

**Terminal HHI and dependence** For terminal HHI and dependence, $j = 2, 3$, define $Z_{nj} = \mathbb{1}_{\{Y_{nj}=1\}}$. The boundary probability satisfies

$$\text{logit}(\pi_{nj}) = c_j^T + d_j^T f_n.$$

Separating the boundary and interior observations, the likelihood for response $j$ is

$$\begin{aligned}&\prod_{n\in\mathcal{O}} p(Y_{nj} \mid f_n, \alpha_j, \lambda_j, \phi_j, c_j^T, d_j^T)\\ &\quad= \left[\prod_{\substack{n\in\mathcal{O}:\\ Y_{nj}=1}} \pi_{nj}\right]\left[\prod_{n\in\mathcal{I}_j}(1-\pi_{nj})\frac{\Gamma(\phi_j)Y_{nj}^{\mu_{nj}\phi_j-1}(1-Y_{nj})^{(1-\mu_{nj})\phi_j-1}}{\Gamma(\mu_{nj}\phi_j)\Gamma((1-\mu_{nj})\phi_j)}\right].\end{aligned}$$

The probabilities $\pi_{nj}$ depend on $c_j^T$, $d_j^T$, and $f_n$, but not on $\lambda_j$. These contributions are therefore constant in the full conditional of $\lambda_j$. Consequently,

$$p(\lambda_j \mid \text{rest}) \propto \left[\prod_{n\in\mathcal{I}_j} \frac{\Gamma(\phi_j) Y_{nj}^{\mu_{nj}\phi_j-1}(1-Y_{nj})^{(1-\mu_{nj})\phi_j-1}}{\Gamma(\mu_{nj}\phi_j)\Gamma((1-\mu_{nj})\phi_j)}\right] p(\lambda_j).$$

Substituting the $\mathcal{N}(0,1)$ prior and absorbing the factors $\Gamma(\phi_j)$ gives

$$p(\lambda_j \mid \text{rest}) \propto \left[\prod_{n\in\mathcal{I}_j} \frac{Y_{nj}^{\mu_{nj}\phi_j-1}(1-Y_{nj})^{(1-\mu_{nj})\phi_j-1}}{\Gamma(\mu_{nj}\phi_j)\Gamma((1-\mu_{nj})\phi_j)}\right] \exp\left(-\frac{\lambda_j^2}{2}\right), \qquad j = 2, 3.$$

Thus, exact ones do not contribute to the full conditional of the interior-mean loading when the factors and boundary parameters are held fixed.

**Chokepoint betweenness** For chokepoint betweenness, define $Z_{n2} = \mathbb{1}_{\{Y_{n2}=0\}}$, with

$$\text{logit}(\rho_{n2}) = c_2^C + d_2^C f_n.$$

The response likelihood is

$$\begin{aligned}
&\prod_{n\in\mathcal{O}} p(Y_{n2} \mid f_n, \alpha_2, \lambda_2, \phi_2, c_2^C, d_2^C) \\
&\quad = \left[\prod_{\substack{n\in\mathcal{O}:\\ Y_{n2}=0}} \rho_{n2}\right] \left[\prod_{n\in\mathcal{I}_2} (1-\rho_{n2}) \frac{\Gamma(\phi_2) Y_{n2}^{\mu_{n2}\phi_2-1}(1-Y_{n2})^{(1-\mu_{n2})\phi_2-1}}{\Gamma(\mu_{n2}\phi_2)\Gamma((1-\mu_{n2})\phi_2)}\right].
\end{aligned}$$

Since $\rho_{n2}$ does not depend on $\lambda_2$, only the interior Beta densities and the loading prior remain:

$$p(\lambda_2 \mid \text{rest}) \propto \left[\prod_{n\in\mathcal{I}_2} \frac{\Gamma(\phi_2) Y_{n2}^{\mu_{n2}\phi_2-1}(1-Y_{n2})^{(1-\mu_{n2})\phi_2-1}}{\Gamma(\mu_{n2}\phi_2)\Gamma((1-\mu_{n2})\phi_2)}\right] p(\lambda_2).$$

Using $\lambda_2 \sim \mathcal{N}(0,1)$ and removing terms constant with respect to $\lambda_2$ yields

$$p(\lambda_2 \mid \text{rest}) \propto \left[\prod_{n\in\mathcal{I}_2} \frac{Y_{n2}^{\mu_{n2}\phi_2-1}(1-Y_{n2})^{(1-\mu_{n2})\phi_2-1}}{\Gamma(\mu_{n2}\phi_2)\Gamma((1-\mu_{n2})\phi_2)}\right] \exp\left(-\frac{\lambda_2^2}{2}\right).$$

As in the one-inflated case, only interior observations contribute to this loading conditional.

**Loadings for the probability mass at one**

For terminal HHI and dependence, the indicators $Z_{nj} = \mathbb{1}_{\{Y_{nj}=1\}}$ satisfy

$$Z_{nj} \mid f_n, c_j^T, d_j^T \sim \text{Bernoulli}(\pi_{nj}), \qquad \text{logit}(\pi_{nj}) = c_j^T + d_j^T f_n, \qquad j = 2, 3.$$

When deriving the full conditional of $d_j^T$, the interior Beta densities are constant because their parameters do not depend on $d_j^T$. The remaining likelihood consists of the boundary probabilities for all observations:

$$p(d_j^T \mid \text{rest}) \propto \left[ \prod_{n \in \mathcal{O}} \pi_{nj}^{Z_{nj}} (1 - \pi_{nj})^{1 - Z_{nj}} \right] p(d_j^T).$$

From the logistic specification,

$$\pi_{nj} = \frac{\exp(c_j^T + d_j^T f_n)}{1 + \exp(c_j^T + d_j^T f_n)}, \qquad 1 - \pi_{nj} = \frac{1}{1 + \exp(c_j^T + d_j^T f_n)}.$$

Hence,

$$\pi_{nj}^{Z_{nj}} (1 - \pi_{nj})^{1 - Z_{nj}} = \frac{\exp\{Z_{nj}(c_j^T + d_j^T f_n)\}}{1 + \exp(c_j^T + d_j^T f_n)}.$$

The prior $d_j^T \sim \mathcal{N}(0, 1)$ has density

$$p(d_j^T) = \frac{1}{\sqrt{2\pi}} \exp\left( -\frac{(d_j^T)^2}{2} \right).$$

Combining these terms gives

$$p(d_j^T \mid \text{rest}) \propto \left[ \prod_{n \in \mathcal{O}} \frac{\exp\{Z_{nj}(c_j^T + d_j^T f_n)\}}{1 + \exp(c_j^T + d_j^T f_n)} \right] \exp\left( -\frac{(d_j^T)^2}{2} \right), \qquad j = 2, 3.$$

Both boundary and interior observations contribute: the former have $Z_{nj} = 1$, whereas the latter have $Z_{nj} = 0$.

**Loading for the probability mass at zero**

For chokepoint betweenness, $Z_{n2} = \mathbb{1}_{\{Y_{n2}=0\}}$, satisfies

$$Z_{n2} \mid f_n, c_2^C, d_2^C \sim \text{Bernoulli}(\rho_{n2}), \qquad \text{logit}(\rho_{n2}) = c_2^C + d_2^C f_n.$$

The interior Beta densities do not depend on $d_2^C$. Retaining the boundary-indicator

likelihood and the loading prior therefore gives

$$p(d_2^C \mid \text{rest}) \propto \left[\prod_{n\in\mathcal{O}} \rho_{n2}^{Z_{n2}}(1-\rho_{n2})^{1-Z_{n2}}\right] p(d_2^C).$$

Using

$$\rho_{n2} = \frac{\exp(c_2^C + d_2^C f_n)}{1+\exp(c_2^C + d_2^C f_n)}, \qquad 1-\rho_{n2} = \frac{1}{1+\exp(c_2^C + d_2^C f_n)},$$

the likelihood contribution becomes

$$\rho_{n2}^{Z_{n2}}(1-\rho_{n2})^{1-Z_{n2}} = \frac{\exp\{Z_{n2}(c_2^C + d_2^C f_n)\}}{1+\exp(c_2^C + d_2^C f_n)}.$$

The prior $d_2^C \sim \mathcal{N}(0,1)$ has density

$$p(d_2^C) = \frac{1}{\sqrt{2\pi}} \exp\left(-\frac{(d_2^C)^2}{2}\right).$$

The resulting full conditional kernel is

$$p(d_2^C \mid \text{rest}) \propto \left[\prod_{n\in\mathcal{O}} \frac{\exp\{Z_{n2}(c_2^C + d_2^C f_n)\}}{1+\exp(c_2^C + d_2^C f_n)}\right] \exp\left(-\frac{(d_2^C)^2}{2}\right).$$

Here exact zeros have $Z_{n2} = 1$, whereas interior observations have $Z_{n2} = 0$, so both contribute to inference on the zero-probability loading.

### D.1.3 Response intercepts

#### Bounded-response interior intercepts

For each bounded response, the interior intercept enters the Beta mean through

$$\text{logit}(\mu_{nj}) = \alpha_j + \lambda_j f_n.$$

Its prior is $\alpha_j \sim \mathcal{N}(m_j, 1.5^2)$, where $(m_1, m_2, m_3) = (-4, 0, 0)$ for terminals and $(-4, -5, -4)$ for chokepoints.

For ordinary Beta responses, all observations contribute to the likelihood of $\alpha_j$. For boundary-inflated responses, the boundary probabilities do not depend on $\alpha_j$, so only the interior Beta densities remain. Both cases can therefore be written as

$$p(\alpha_j \mid \text{rest}) \propto \left[\prod_{n\in\mathcal{I}_j} p(Y_{nj} \mid f_n, \alpha_j, \lambda_j, \phi_j, 0 < Y_{nj} < 1)\right] p(\alpha_j).$$

For ordinary Beta responses, $\mathcal{I}_j = \mathcal{O}$.

The interior density is

$$p(Y_{nj} \mid f_n, \alpha_j, \lambda_j, \phi_j, 0 < Y_{nj} < 1) = \frac{\Gamma(\phi_j) Y_{nj}^{\mu_{nj}\phi_j - 1}(1 - Y_{nj})^{(1-\mu_{nj})\phi_j - 1}}{\Gamma(\mu_{nj}\phi_j)\Gamma((1-\mu_{nj})\phi_j)}.$$

The prior density is

$$p(\alpha_j) = \frac{1}{1.5\sqrt{2\pi}} \exp\left\{ -\frac{(\alpha_j - m_j)^2}{2(1.5)^2} \right\}.$$

Substituting these densities gives

$$p(\alpha_j \mid \text{rest}) \propto \left[ \prod_{n \in \mathcal{I}_j} \frac{\Gamma(\phi_j) Y_{nj}^{\mu_{nj}\phi_j - 1}(1 - Y_{nj})^{(1-\mu_{nj})\phi_j - 1}}{\Gamma(\mu_{nj}\phi_j)\Gamma((1-\mu_{nj})\phi_j)} \right] \exp\left\{ -\frac{(\alpha_j - m_j)^2}{2(1.5)^2} \right\}.$$

Since $\phi_j$ is held fixed, the factors $\Gamma(\phi_j)$ do not depend on $\alpha_j$. Absorbing them into the normalising constant yields

$$p(\alpha_j \mid \text{rest}) \propto \left[ \prod_{n \in \mathcal{I}_j} \frac{Y_{nj}^{\mu_{nj}\phi_j - 1}(1 - Y_{nj})^{(1-\mu_{nj})\phi_j - 1}}{\Gamma(\mu_{nj}\phi_j)\Gamma((1-\mu_{nj})\phi_j)} \right] \exp\left\{ -\frac{(\alpha_j - m_j)^2}{2(1.5)^2} \right\}, \qquad j = 1, 2, 3.$$

The likelihood depends on $\alpha_j$ through $\mu_{nj}$.

**Binary-response intercepts**

For binary response $k$,

$$B_{nk} \mid f_n, a_k, \ell_k \sim \text{Bernoulli}(p_{nk}), \qquad \text{logit}(p_{nk}) = a_k + \ell_k f_n.$$

Only the likelihood of response $k$ and the prior of $a_k$ depend on this intercept. Therefore,

$$p(a_k \mid \text{rest}) \propto \left[ \prod_{n \in \mathcal{O}} p(B_{nk} \mid f_n, a_k, \ell_k) \right] p(a_k).$$

Using the logistic specification,

$$\begin{aligned} p(B_{nk} \mid f_n, a_k, \ell_k) &= p_{nk}^{B_{nk}}(1 - p_{nk})^{1 - B_{nk}} \\ &= \frac{\exp\{B_{nk}(a_k + \ell_k f_n)\}}{1 + \exp(a_k + \ell_k f_n)}. \end{aligned}$$

The prior $a_k \sim \mathcal{N}(0, 2^2)$ has density

$$p(a_k) = \frac{1}{2\sqrt{2\pi}} \exp\left( -\frac{a_k^2}{8} \right).$$

Combining likelihood and prior gives

$$p(a_k \mid \text{rest}) \propto \left[\prod_{n\in\mathcal{O}} \frac{\exp\{B_{nk}(a_k+\ell_k f_n)\}}{1+\exp(a_k+\ell_k f_n)}\right] \exp\left(-\frac{a_k^2}{8}\right).$$

Equivalently, collecting the exponential terms and removing those independent of $a_k$ gives

$$p(a_k \mid \text{rest}) \propto \exp\left\{ -\frac{a_k^2}{8} + a_k \sum_{n\in\mathcal{O}} B_{nk} - \sum_{n\in\mathcal{O}} \log[1+\exp(a_k+\ell_k f_n)] \right\}.$$

This expression applies to $k = 1, 2, 3$ for terminals and $k = 1, 2$ for chokepoints.

**Intercepts of the probability of mass at one**

For terminal HHI and dependence, define $Z_{nj} = \mathbb{1}_{\{Y_{nj}=1\}}$, with

$$Z_{nj} \mid f_n, c_j^T, d_j^T \sim \text{Bernoulli}(\pi_{nj}), \qquad \text{logit}(\pi_{nj}) = c_j^T + d_j^T f_n, \qquad j = 2, 3.$$

The interior Beta densities do not depend on $c_j^T$. Retaining the boundary-indicator likelihood and the intercept prior gives

$$p(c_j^T \mid \text{rest}) \propto \left[\prod_{n\in\mathcal{O}} \pi_{nj}^{Z_{nj}} (1-\pi_{nj})^{1-Z_{nj}}\right] p(c_j^T).$$

The logistic link implies

$$\pi_{nj}^{Z_{nj}} (1-\pi_{nj})^{1-Z_{nj}} = \frac{\exp\{Z_{nj}(c_j^T + d_j^T f_n)\}}{1+\exp(c_j^T + d_j^T f_n)}.$$

The prior $c_j^T \sim \mathcal{N}(0, 1.5^2)$ has density

$$p(c_j^T) = \frac{1}{1.5\sqrt{2\pi}} \exp\left\{ -\frac{(c_j^T)^2}{2(1.5)^2} \right\}.$$

Substituting gives

$$p(c_j^T \mid \text{rest}) \propto \left[\prod_{n\in\mathcal{O}} \frac{\exp\{Z_{nj}(c_j^T + d_j^T f_n)\}}{1+\exp(c_j^T + d_j^T f_n)}\right] \exp\left\{ -\frac{(c_j^T)^2}{2(1.5)^2} \right\}, \qquad j = 2, 3.$$

Equivalently, removing terms independent of $c_j^T$ yields

$$p(c_j^T \mid \text{rest}) \propto \exp\left\{-\frac{(c_j^T)^2}{2(1.5)^2}+c_j^T\sum_{n\in\mathcal{O}} Z_{nj}-\sum_{n\in\mathcal{O}}\log\big[1+\exp(c_j^T+d_j^T f_n)\big]\right\}, \qquad j=2,3.$$

**Intercept of the probability of mass at zero**

For chokepoint betweenness, define $Z_{n2} = \mathbb{1}_{\{Y_{n2}=0\}}$, with

$$Z_{n2} \mid f_n, c_2^C, d_2^C \sim \text{Bernoulli}(\rho_{n2}), \qquad \text{logit}(\rho_{n2}) = c_2^C + d_2^C f_n.$$

The interior Beta densities do not depend on $c_2^C$. Its full conditional therefore satisfies

$$p(c_2^C \mid \text{rest}) \propto \left[\prod_{n\in\mathcal{O}} \rho_{n2}^{Z_{n2}}(1-\rho_{n2})^{1-Z_{n2}}\right] p(c_2^C).$$

Using the logistic link,

$$\rho_{n2}^{Z_{n2}}(1-\rho_{n2})^{1-Z_{n2}} = \frac{\exp\{Z_{n2}(c_2^C+d_2^C f_n)\}}{1+\exp(c_2^C+d_2^C f_n)}.$$

The prior $c_2^C \sim \mathcal{N}(-5, 1.5^2)$ has density

$$p(c_2^C) = \frac{1}{1.5\sqrt{2\pi}}\exp\left\{-\frac{(c_2^C+5)^2}{2(1.5)^2}\right\}.$$

Combining these expressions gives

$$p(c_2^C \mid \text{rest}) \propto \left[\prod_{n\in\mathcal{O}} \frac{\exp\{Z_{n2}(c_2^C+d_2^C f_n)\}}{1+\exp(c_2^C+d_2^C f_n)}\right]\exp\left\{-\frac{(c_2^C+5)^2}{2(1.5)^2}\right\}.$$

Equivalently, removing terms independent of $c_2^C$ yields

$$p(c_2^C \mid \text{rest}) \propto \exp\left\{-\frac{(c_2^C+5)^2}{2(1.5)^2}+c_2^C\sum_{n\in\mathcal{O}} Z_{n2}-\sum_{n\in\mathcal{O}}\log\big[1+\exp(c_2^C+d_2^C f_n)\big]\right\}.$$

### D.1.4 Beta precision parameters

For each bounded response, the precision $\phi_j$ enters both Beta shape parameters:

$$A_{nj} = \mu_{nj}\phi_j, \qquad D_{nj} = (1-\mu_{nj})\phi_j.$$

For ordinary Beta responses, all observations contribute to the likelihood of the precision parameter. For boundary-inflated responses, the boundary probabilities do

not depend on $\phi_j$, so only the interior observations contribute. Retaining the relevant likelihood and prior gives

$$p(\phi_j \mid \text{rest}) \propto \left[\prod_{n \in \mathcal{I}_j} p(Y_{nj} \mid f_n, \alpha_j, \lambda_j, \phi_j, 0 < Y_{nj} < 1)\right] p(\phi_j).$$

The interior Beta density is

$$p(Y_{nj} \mid f_n, \alpha_j, \lambda_j, \phi_j, 0 < Y_{nj} < 1) = \frac{\Gamma(\phi_j) Y_{nj}^{\mu_{nj}\phi_j - 1}(1 - Y_{nj})^{(1-\mu_{nj})\phi_j - 1}}{\Gamma(\mu_{nj}\phi_j)\Gamma((1-\mu_{nj})\phi_j)}.$$

The prior is specified on the logarithmic scale:

$$\log \phi_j \sim \mathcal{N}(\log 10, 0.7^2).$$

Expressed as a density for $\phi_j$, this becomes

$$p(\phi_j) = \frac{1}{0.7\,\phi_j\sqrt{2\pi}} \exp\left\{-\frac{(\log \phi_j - \log 10)^2}{2(0.7)^2}\right\}, \qquad \phi_j > 0.$$

The factor $1/\phi_j$ follows from the change-of-variables derivative $|d \log \phi_j / d\phi_j| = 1/\phi_j$.

Substituting the likelihood and prior densities, and absorbing constants independent of $\phi_j$, gives the full conditional kernel

$$p(\phi_j \mid \text{rest}) \propto \left[\prod_{n \in \mathcal{I}_j} \frac{\Gamma(\phi_j) Y_{nj}^{\mu_{nj}\phi_j - 1}(1 - Y_{nj})^{(1-\mu_{nj})\phi_j - 1}}{\Gamma(\mu_{nj}\phi_j)\Gamma((1-\mu_{nj})\phi_j)}\right] \frac{1}{\phi_j} \exp\left\{-\frac{(\log \phi_j - \log 10)^2}{2(0.7)^2}\right\},$$

for $j = 1, 2, 3$ and $\phi_j > 0$.

### D.1.5 Latent factors

To derive the full conditional of a particular latent factor $f_n$, the regression coefficients, response parameters, and all other latent factors are held fixed. The parameter priors and the likelihood contributions from other observations are therefore constant with respect to $f_n$. Retaining the factors of the joint posterior that depend on $f_n$ gives

$$p(f_n \mid \text{rest}) \propto p(\mathbf{Y}_n, \mathbf{B}_n \mid f_n, \boldsymbol{\theta}) p(f_n \mid \boldsymbol{\beta}, \mathbf{x}_n).$$

The Gaussian factor-regression density is

$$p(f_n \mid \boldsymbol{\beta}, \mathbf{x}_n) = \frac{1}{\sqrt{2\pi}} \exp\left\{-\frac{1}{2}\left(f_n - \mathbf{x}_n^{\mathsf{T}}\boldsymbol{\beta}\right)^2\right\}.$$

Conditional independence of the responses gives

$$p(\mathbf{Y}_n, \mathbf{B}_n \mid f_n, \boldsymbol{\theta}) = \prod_{j=1}^{3} g_j(Y_{nj} \mid f_n, \boldsymbol{\theta}) \prod_{k=1}^{K} p(B_{nk} \mid f_n, a_k, \ell_k),$$

where $g_j$ is the bounded-response contribution introduced in the main text.

Denote this observation-specific likelihood by $L_n(f_n)$. For terminals, it is

$$\begin{aligned} L_n^T(f_n) = p(Y_{n1} \mid f_n, \alpha_1, \lambda_1, \phi_1) \\ \times \prod_{j=2}^{3} p(Y_{nj} \mid f_n, \alpha_j, \lambda_j, \phi_j, c_j^T, d_j^T) \\ \times \prod_{k=1}^{3} p(B_{nk} \mid f_n, a_k, \ell_k). \end{aligned}$$

The bounded-response terms consist of the ordinary Beta density for PageRank and the one-inflated Beta contributions for HHI and dependence.

For chokepoints,

$$\begin{aligned} L_n^C(f_n) = p(Y_{n1} \mid f_n, \alpha_1, \lambda_1, \phi_1) \\ \times p(Y_{n2} \mid f_n, \alpha_2, \lambda_2, \phi_2, c_2^C, d_2^C) \\ \times p(Y_{n3} \mid f_n, \alpha_3, \lambda_3, \phi_3) \\ \times \prod_{k=1}^{2} p(B_{nk} \mid f_n, a_k, \ell_k). \end{aligned}$$

Here PageRank and flow share contribute ordinary Beta densities, while betweenness contributes the zero-inflated Beta likelihood.

Combining the observation-specific likelihood with the Gaussian factor-regression density and removing its constant prefactor yields

$$p(f_n \mid \text{rest}) \propto L_n(f_n) \exp\left\{ -\frac{1}{2} \left( f_n - \mathbf{x}_n^{\mathsf{T}} \boldsymbol{\beta} \right)^2 \right\},$$

where $L_n = L_n^T$ for terminals and $L_n = L_n^C$ for chokepoints.

The factor enters the interior means, binary probabilities, and boundary probabilities. Its full conditional therefore combines information from all responses observed for the same node–month. Although the factor-regression density is Gaussian, the response likelihood is a non-Gaussian function of $f_n$, so the resulting full conditional does not have a standard conjugate form.

## D.2 Random-intercept rank-one specification

This section derives the full conditional posterior distributions associated with the random-intercept extension introduced in Section 5.2.2. The response likelihoods and their parameter priors remain unchanged from the baseline specification. Consequently, the corresponding full conditional kernels retain the forms derived in the preceding sections, conditional on the latent factors.

The derivations below therefore focus on the factor-regression coefficients, latent factors, node-specific intercepts, and their common scale parameter.

Let $N_{\text{node}}$ denote the number of distinct nodes included in estimation, and let $N_{\text{obs}} = |\mathcal{O}|$ denote the number of node–month observations. For each observation $n \in \mathcal{O}$, let $i(n)$ identify its corresponding node.

The extended factor regression is

$$f_n \mid \boldsymbol{\beta}, \mu_{i(n)}, \mathbf{x}_n \sim \mathcal{N}\left(\mathbf{x}_n^{\mathsf{T}}\boldsymbol{\beta} + \mu_{i(n)}, 1\right),$$

with hierarchical priors

$$\mu_i \mid \sigma_\mu \overset{\text{iid}}{\sim} \mathcal{N}(0, \sigma_\mu^2), \qquad \sigma_\mu \sim \text{HalfNormal}(1).$$

The derivations are expressed in terms of the node-specific intercepts $\mu_i$, corresponding to the hierarchical model formulation.

### D.2.1 Factor-regression coefficients

Conditional on the latent factors and node-specific intercepts, the response likelihood does not depend on $\boldsymbol{\beta}$. Therefore,

$$p(\boldsymbol{\beta} \mid \text{rest}) \propto p(\mathbf{f} \mid \boldsymbol{\beta}, \boldsymbol{\mu}, \mathbf{X}) p(\boldsymbol{\beta}).$$

Combining the Gaussian factor-regression density with the independent $\mathcal{N}(0, 0.5^2)$ coefficient priors gives

$$p(\boldsymbol{\beta} \mid \text{rest}) \propto \prod_{n \in \mathcal{O}} \exp\left[-\frac{1}{2}\left(f_n - \mathbf{x}_n^{\mathsf{T}}\boldsymbol{\beta} - \mu_{i(n)}\right)^2\right] \prod_{r=1}^{q} \exp\left(-\frac{\beta_r^2}{2(0.5)^2}\right).$$

Let $\boldsymbol{\mu}_{\mathcal{O}}$ denote the $N_{\text{obs}}$-dimensional vector whose $n^{nt}$ element is $\mu_{i(n)}$. Combining the exponential terms gives

$$p(\boldsymbol{\beta} \mid \text{rest}) \propto \exp\left\{ -\frac{1}{2}\Big[(\mathbf{f} - \boldsymbol{\mu}_{\mathcal{O}} - \mathbf{X}\boldsymbol{\beta})^{\mathsf{T}}(\mathbf{f} - \boldsymbol{\mu}_{\mathcal{O}} - \mathbf{X}\boldsymbol{\beta}) + 4\boldsymbol{\beta}^{\mathsf{T}}\boldsymbol{\beta}\Big] \right\}.$$

Expanding the quadratic term,

$$(\mathbf{f} - \boldsymbol{\mu}_{\mathcal{O}} - \mathbf{X}\boldsymbol{\beta})^{\mathsf{T}}(\mathbf{f} - \boldsymbol{\mu}_{\mathcal{O}} - \mathbf{X}\boldsymbol{\beta}) = (\mathbf{f} - \boldsymbol{\mu}_{\mathcal{O}})^{\mathsf{T}}(\mathbf{f} - \boldsymbol{\mu}_{\mathcal{O}}) - 2\boldsymbol{\beta}^{\mathsf{T}}\mathbf{X}^{\mathsf{T}}(\mathbf{f} - \boldsymbol{\mu}_{\mathcal{O}}) + \boldsymbol{\beta}^{\mathsf{T}}\mathbf{X}^{\mathsf{T}}\mathbf{X}\boldsymbol{\beta}.$$

The first term does not depend on $\boldsymbol{\beta}$ and can be absorbed into the normalising constant. The resulting kernel is

$$p(\boldsymbol{\beta} \mid \text{rest}) \exp\left\{ -\frac{1}{2}\Big[\boldsymbol{\beta}^{\mathsf{T}}(\mathbf{X}^{\mathsf{T}}\mathbf{X} + 4\mathbf{I}_q)\boldsymbol{\beta} - 2\boldsymbol{\beta}^{\mathsf{T}}\mathbf{X}^{\mathsf{T}}(\mathbf{f} - \boldsymbol{\mu}_{\mathcal{O}})\Big] \right\}.$$

Define

$$\mathbf{V}_\beta = (\mathbf{X}^{\mathsf{T}}\mathbf{X} + 4\mathbf{I}_q)^{-1}, \qquad \mathbf{m}_\beta = \mathbf{V}_\beta \mathbf{X}^{\mathsf{T}}(\mathbf{f} - \boldsymbol{\mu}_{\mathcal{O}}).$$

Since

$$\mathbf{V}_\beta^{-1}\mathbf{m}_\beta = \mathbf{X}^{\mathsf{T}}(\mathbf{f} - \boldsymbol{\mu}_{\mathcal{O}}),$$

completing the square gives

$$\boldsymbol{\beta}^{\mathsf{T}}\mathbf{V}_\beta^{-1}\boldsymbol{\beta} - 2\boldsymbol{\beta}^{\mathsf{T}}\mathbf{X}^{\mathsf{T}}(\mathbf{f} - \boldsymbol{\mu}_{\mathcal{O}}) = (\boldsymbol{\beta} - \mathbf{m}_\beta)^{\mathsf{T}}\mathbf{V}_\beta^{-1}(\boldsymbol{\beta} - \mathbf{m}_\beta) - \mathbf{m}_\beta^{\mathsf{T}}\mathbf{V}_\beta^{-1}\mathbf{m}_\beta.$$

The final term is constant with respect to $\boldsymbol{\beta}$. Therefore,

$$p(\boldsymbol{\beta} \mid \text{rest}) \propto \exp\left[ -\frac{1}{2}(\boldsymbol{\beta} - \mathbf{m}_\beta)^{\mathsf{T}}\mathbf{V}_\beta^{-1}(\boldsymbol{\beta} - \mathbf{m}_\beta) \right],$$

which yields

$$\boldsymbol{\beta} \mid \text{rest} \sim \mathcal{N}_q\Big( (\mathbf{X}^{\mathsf{T}}\mathbf{X} + 4\mathbf{I}_q)^{-1}\mathbf{X}^{\mathsf{T}}(\mathbf{f} - \boldsymbol{\mu}_{\mathcal{O}}), (\mathbf{X}^{\mathsf{T}}\mathbf{X} + 4\mathbf{I}_q)^{-1} \Big).$$

### D.2.2 Latent factors

For each observation $n$, the full conditional distribution of $f_n$ combines the observation-specific response likelihood with the Gaussian factor regression:

$$p(f_n \mid \text{rest}) \propto L_n(f_n) p(f_n \mid \boldsymbol{\beta}, \mu_{i(n)}, \mathbf{x}_n).$$

Substituting the Gaussian density gives

$$p(f_n \mid \text{rest}) \propto L_n(f_n) \exp\left[ -\frac{1}{2}\left(f_n - \mathbf{x}_n^{\mathsf{T}}\boldsymbol{\beta} - \mu_{i(n)}\right)^2 \right].$$

The likelihood contribution $L_n(f_n)$ is identical to that of the baseline model. Consequently, the extension modifies only the centre of the Gaussian factor-regression term, replacing $\mathbf{x}_n^{\mathsf{T}}\boldsymbol{\beta}$ with $\mathbf{x}_n^{\mathsf{T}}\boldsymbol{\beta} + \mu_{i(n)}$.

Because the response likelihood contains Beta and Bernoulli components linked to $f_n$ through logistic transformations, the resulting conditional distribution is not generally conjugate.

### D.2.3 Node-specific intercepts

For each node $i$, define $\mathcal{O}_i = \{n \in \mathcal{O} : i(n) = i\}, n_i = |\mathcal{O}_i|$. Conditional on the latent factors and the remaining parameters, only the factor-regression contributions associated with node $i$ and the Gaussian prior for $\mu_i$ depend on this parameter. Therefore,

$$p(\mu_i \mid \text{rest}) \propto p(\mu_i \mid \sigma_\mu) \prod_{n \in \mathcal{O}_i} p(f_n \mid \boldsymbol{\beta}, \mu_i, \mathbf{x}_n).$$

Writing the relevant densities explicitly,

$$p(\mu_i \mid \text{rest}) \propto \exp\left(-\frac{\mu_i^2}{2\sigma_\mu^2}\right) \prod_{n \in \mathcal{O}_i} \exp\left[-\frac{1}{2}\left(f_n - \mathbf{x}_n^\top \boldsymbol{\beta} - \mu_i\right)^2\right].$$

Define the factor-regression residuals $r_n = f_n - \mathbf{x}_n^\top \boldsymbol{\beta}$. The conditional kernel becomes

$$p(\mu_i \mid \text{rest}) \propto \exp\left\{-\frac{1}{2}\left[\frac{\mu_i^2}{\sigma_\mu^2} + \sum_{n \in \mathcal{O}_i} (r_n - \mu_i)^2\right]\right\}.$$

Expanding the sum of squared residuals,

$$\sum_{n \in \mathcal{O}_i} (r_n - \mu_i)^2 = \sum_{n \in \mathcal{O}_i} r_n^2 - 2\mu_i \sum_{n \in \mathcal{O}_i} r_n + n_i \mu_i^2.$$

Removing the term $\sum_{n \in \mathcal{O}_i} r_n^2$, which does not depend on $\mu_i$, gives

$$p(\mu_i \mid \text{rest}) \propto \exp\left\{-\frac{1}{2}\left[\left(n_i + \frac{1}{\sigma_\mu^2}\right)\mu_i^2 - 2\mu_i \sum_{n \in \mathcal{O}_i} r_n\right]\right\}.$$

Define

$$V_{\mu_i} = \left(n_i + \frac{1}{\sigma_\mu^2}\right)^{-1}, \qquad m_{\mu_i} = V_{\mu_i} \sum_{n \in \mathcal{O}_i} r_n.$$

Since

$$V_{\mu_i}^{-1} m_{\mu_i} = \sum_{n \in \mathcal{O}_i} r_n,$$

completing the square yields

$$V_{\mu_i}^{-1}\mu_i^2 - 2\mu_i \sum_{n \in \mathcal{O}_i} r_n = V_{\mu_i}^{-1}(\mu_i - m_{\mu_i})^2 - V_{\mu_i}^{-1} m_{\mu_i}^2.$$

The final term does not depend on $\mu_i$. Thus,

$$p(\mu_i \mid \text{rest}) \propto \exp\left[-\frac{(\mu_i - m_{\mu_i})^2}{2V_{\mu_i}}\right],$$

which gives the conjugate full conditional

$$\mu_i \mid \text{rest} \sim \mathcal{N}\left(V_{\mu_i} \sum_{n \in \mathcal{O}_i} (f_n - \mathbf{x}_n^\top \boldsymbol{\beta}),\ V_{\mu_i}\right),$$

where $V_{\mu_i} = \left(n_i + \sigma_\mu^{-2}\right)^{-1}$.

### D.2.4 Node-effect scale parameter

The full conditional distribution of $\sigma_\mu$ depends on its Half-Normal prior and the hierarchical Gaussian distribution of the node-specific intercepts. Conditional on $\boldsymbol{\mu}$, the factor-regression likelihood does not depend directly on $\sigma_\mu$. Therefore,

$$p(\sigma_\mu \mid \text{rest}) \propto p(\boldsymbol{\mu} \mid \sigma_\mu) p(\sigma_\mu).$$

The hierarchical prior contributes

$$p(\boldsymbol{\mu} \mid \sigma_\mu) \propto \prod_{i=1}^{N_{\text{node}}} \frac{1}{\sigma_\mu} \exp\left(-\frac{\mu_i^2}{2\sigma_\mu^2}\right),$$

for $\sigma_\mu > 0$. Combining the product gives

$$p(\boldsymbol{\mu} \mid \sigma_\mu) \propto \sigma_\mu^{-N_{\text{node}}} \exp\left(-\frac{\sum_{i=1}^{N_{\text{node}}} \mu_i^2}{2\sigma_\mu^2}\right).$$

The Half-Normal prior with unit scale has kernel

$$p(\sigma_\mu) \propto \exp\left(-\frac{\sigma_\mu^2}{2}\right) \mathbb{I}(\sigma_\mu > 0).$$

Multiplying these contributions yields

$$p(\sigma_\mu \mid \text{rest}) \propto \sigma_\mu^{-N_{\text{node}}} \exp\left[-\frac{1}{2}\left(\sigma_\mu^2 + \frac{\sum_{i=1}^{N_{\text{node}}} \mu_i^2}{\sigma_\mu^2}\right)\right] \mathbb{I}(\sigma_\mu > 0).$$

Equivalently, the log-kernel is

$$\log p(\sigma_\mu \mid \text{rest}) = -N_{\text{node}} \log \sigma_\mu - \frac{\sigma_\mu^2}{2} - \frac{\sum_{i=1}^{N_{\text{node}}} \mu_i^2}{2\sigma_\mu^2} + C,$$

where $C$ does not depend on $\sigma_\mu$.

Unlike the conditional distribution of the node-specific intercepts, this conditional distribution does not have a standard conjugate form for $\sigma_\mu$.

The expressions above correspond to the hierarchical formulation of the model. In the computational implementation, the node effects are represented using the equivalent non-centred transformation $\mu_i = \sigma_\mu u_i$, where $u_i \sim \mathcal{N}(0, 1)$. This changes the sampling coordinates, but not the posterior distribution of the model quantities of interest.

# Appendix E

# Additional posterior results

## E.1 Baseline rank-one specification

This section provides supplementary results for the baseline rank-one model, complementing the posterior analysis presented in Section 6.1.

### E.1.1 Terminals

#### Terminals residuals heterogeneity

The following table provides additional descriptive evidence of persistent terminal-level residual patterns under the baseline rank-one specification.

#### Terminals posterior predictive checks

Table E.2 reports marginal posterior predictive checks for the terminal baseline rank-one model, separately for all observations, lagged importers and lagged exporters. The reported intervals describe the distribution of each statistic across 200 conditional in-sample replicated datasets.

Table E.2: Marginal posterior predictive checks for terminal responses under the baseline rank-one model. All statistics are on the original response scale; intervals are based on 200 conditional in-sample replicates.

| Response | Statistic | Observed | Rep. mean | 95% lower | 95% upper |
|---|---|---|---|---|---|
| *All terminals* | | | | | |
| PageRank | Mean | 0.0185 | 0.0185 | 0.0183 | 0.0187 |
| PageRank | SD | 0.0133 | 0.0124 | 0.0121 | 0.0127 |
| PageRank | 5th percentile | 0.0089 | 0.0069 | 0.0067 | 0.0072 |
| PageRank | Median | 0.0147 | 0.0155 | 0.0152 | 0.0158 |
| PageRank | 95th percentile | 0.0391 | 0.0395 | 0.0384 | 0.0406 |

*Continued on next page*

*Table continued from previous page*

| Response | Statistic | Observed | Rep. mean | 95% lower | 95% upper |
|---|---|---|---|---|---|
| HHI | Mean | 0.5887 | 0.5899 | 0.5822 | 0.5975 |
| HHI | SD | 0.3254 | 0.3233 | 0.3193 | 0.3272 |
| HHI | 5th percentile | 0.1268 | 0.1320 | 0.1219 | 0.1437 |
| HHI | Median | 0.5045 | 0.4988 | 0.4863 | 0.5110 |
| HHI | 95th percentile | 1.0000 | 1.0000 | 1.0000 | 1.0000 |
| HHI | Proportion equal to one | 0.3409 | 0.3412 | 0.3275 | 0.3551 |
| Dependence | Mean | 0.4948 | 0.4953 | 0.4818 | 0.5116 |
| Dependence | SD | 0.3219 | 0.3179 | 0.3116 | 0.3253 |
| Dependence | 5th percentile | 0.0716 | 0.0796 | 0.0717 | 0.0879 |
| Dependence | Median | 0.4283 | 0.4175 | 0.4020 | 0.4374 |
| Dependence | 95th percentile | 1.0000 | 1.0000 | 1.0000 | 1.0000 |
| Dependence | Proportion equal to one | 0.2076 | 0.2071 | 0.1885 | 0.2248 |
| *Lagged importers* | | | | | |
| PageRank | Mean | 0.0145 | 0.0145 | 0.0143 | 0.0148 |
| PageRank | SD | 0.0045 | 0.0058 | 0.0056 | 0.0061 |
| PageRank | 5th percentile | 0.0089 | 0.0065 | 0.0062 | 0.0068 |
| PageRank | Median | 0.0136 | 0.0137 | 0.0135 | 0.0140 |
| PageRank | 95th percentile | 0.0233 | 0.0252 | 0.0246 | 0.0261 |
| HHI | Mean | 0.6745 | 0.6810 | 0.6718 | 0.6908 |
| HHI | SD | 0.2966 | 0.2981 | 0.2937 | 0.3028 |
| HHI | 5th percentile | 0.2446 | 0.2476 | 0.2342 | 0.2597 |
| HHI | Median | 0.5701 | 0.6060 | 0.5858 | 0.6257 |
| HHI | 95th percentile | 1.0000 | 1.0000 | 1.0000 | 1.0000 |
| HHI | Proportion equal to one | 0.4189 | 0.4346 | 0.4163 | 0.4521 |
| Dependence | Mean | 0.4812 | 0.4763 | 0.4601 | 0.4945 |
| Dependence | SD | 0.3574 | 0.3330 | 0.3256 | 0.3402 |
| Dependence | 5th percentile | 0.0614 | 0.0687 | 0.0603 | 0.0777 |
| Dependence | Median | 0.3472 | 0.3731 | 0.3545 | 0.3945 |
| Dependence | 95th percentile | 1.0000 | 1.0000 | 1.0000 | 1.0000 |
| Dependence | Proportion equal to one | 0.2645 | 0.2391 | 0.2172 | 0.2607 |
| *Lagged exporters* | | | | | |
| PageRank | Mean | 0.0269 | 0.0268 | 0.0264 | 0.0271 |
| PageRank | SD | 0.0198 | 0.0173 | 0.0166 | 0.0178 |
| PageRank | 5th percentile | 0.0088 | 0.0090 | 0.0085 | 0.0096 |
| PageRank | Median | 0.0210 | 0.0225 | 0.0219 | 0.0232 |
| PageRank | 95th percentile | 0.0714 | 0.0619 | 0.0587 | 0.0648 |
| HHI | Mean | 0.4119 | 0.4019 | 0.3917 | 0.4126 |
| HHI | SD | 0.3103 | 0.2905 | 0.2818 | 0.2993 |
| HHI | 5th percentile | 0.0877 | 0.0609 | 0.0516 | 0.0727 |

*Continued on next page*

*Table continued from previous page*

| Response | Statistic | Observed | Rep. mean | 95% lower | 95% upper |
|---|---|---|---|---|---|
| HHI | Median | 0.3333 | 0.3258 | 0.3144 | 0.3409 |
| HHI | 95th percentile | 1.0000 | 1.0000 | 1.0000 | 1.0000 |
| HHI | Proportion equal to one | 0.1800 | 0.1486 | 0.1348 | 0.1623 |
| Dependence | Mean | 0.5228 | 0.5343 | 0.5165 | 0.5550 |
| Dependence | SD | 0.2296 | 0.2800 | 0.2709 | 0.2882 |
| Dependence | 5th percentile | 0.1852 | 0.1216 | 0.1048 | 0.1382 |
| Dependence | Median | 0.4972 | 0.5010 | 0.4771 | 0.5286 |
| Dependence | 95th percentile | 1.0000 | 1.0000 | 1.0000 | 1.0000 |
| Dependence | Proportion equal to one | 0.0904 | 0.1411 | 0.1179 | 0.1614 |

Table E.3 presents the complete set of 45 pairwise Spearman correlation checks for the terminal baseline rank-one model, covering all 15 response pairs across the full estimation sample and the importer and exporter subsamples. For each comparison, the observed correlation is reported alongside the median and the 95% interval obtained from 200 conditional in-sample posterior predictive replicates. These results complement the selected discrepancies discussed in Section 6.1.1.

Table E.3: Complete joint posterior predictive checks for the terminal baseline rank-one model. Entries are Spearman correlations. Replicated intervals use 200 conditional in-sample replicates. PR = PageRank; AP = articulation-point status; GB = incident-global-bridge status; toAP = adjacency to an articulation point; DEP = dependence.

| Response pair | Observed | Rep. median | 95% lower | 95% upper |
|---|---|---|---|---|
| *All terminals* | | | | |
| PR–HHI | -0.639 | -0.712 | -0.730 | -0.695 |
| PR–DEP | 0.314 | 0.158 | 0.115 | 0.197 |
| PR–AP | 0.372 | 0.262 | 0.226 | 0.304 |
| PR–GB | -0.238 | -0.293 | -0.331 | -0.252 |
| PR–toAP | -0.015 | 0.024 | -0.026 | 0.070 |
| HHI–DEP | -0.171 | -0.158 | -0.196 | -0.114 |
| HHI–AP | -0.290 | -0.259 | -0.297 | -0.227 |
| HHI–GB | 0.533 | 0.295 | 0.248 | 0.335 |
| HHI–toAP | -0.038 | -0.025 | -0.065 | 0.021 |
| DEP–AP | 0.157 | 0.082 | 0.055 | 0.116 |
| DEP–GB | -0.066 | -0.070 | -0.113 | -0.037 |
| DEP–toAP | -0.082 | 0.005 | -0.027 | 0.037 |
| AP–GB | 0.290 | -0.124 | -0.162 | -0.095 |
| AP–toAP | -0.063 | 0.014 | -0.028 | 0.046 |

*Continued on next page*

*Table continued from previous page*

| Response pair | Observed | Rep. median | 95% lower | 95% upper |
|---|---|---|---|---|
| GB–toAP | 0.060 | -0.011 | -0.043 | 0.023 |
| *Importers* | | | | |
| PR–HHI | -0.416 | -0.576 | -0.602 | -0.543 |
| PR–DEP | 0.209 | 0.068 | 0.023 | 0.114 |
| PR–AP | 0.232 | 0.113 | 0.068 | 0.150 |
| PR–GB | -0.258 | -0.177 | -0.216 | -0.132 |
| PR–toAP | -0.043 | 0.015 | -0.027 | 0.057 |
| HHI–DEP | -0.163 | -0.075 | -0.121 | -0.027 |
| HHI–AP | -0.137 | -0.119 | -0.153 | -0.080 |
| HHI–GB | 0.644 | 0.189 | 0.137 | 0.226 |
| HHI–toAP | 0.014 | -0.016 | -0.060 | 0.030 |
| DEP–AP | 0.033 | 0.016 | -0.014 | 0.052 |
| DEP–GB | -0.154 | -0.020 | -0.062 | 0.016 |
| DEP–toAP | -0.045 | 0.000 | -0.036 | 0.040 |
| AP–GB | 0.137 | -0.037 | -0.082 | 0.003 |
| AP–toAP | 0.019 | 0.007 | -0.034 | 0.046 |
| GB–toAP | 0.131 | -0.005 | -0.046 | 0.037 |
| *Exporters* | | | | |
| PR–HHI | -0.818 | -0.779 | -0.802 | -0.754 |
| PR–DEP | 0.516 | 0.285 | 0.223 | 0.356 |
| PR–AP | 0.276 | 0.359 | 0.294 | 0.406 |
| PR–GB | -0.198 | -0.356 | -0.412 | -0.301 |
| PR–toAP | 0.166 | 0.039 | -0.037 | 0.093 |
| HHI–DEP | -0.102 | -0.260 | -0.314 | -0.199 |
| HHI–AP | -0.159 | -0.328 | -0.379 | -0.266 |
| HHI–GB | 0.398 | 0.336 | 0.268 | 0.397 |
| HHI–toAP | -0.230 | -0.033 | -0.089 | 0.033 |
| DEP–AP | 0.307 | 0.144 | 0.093 | 0.197 |
| DEP–GB | 0.195 | -0.118 | -0.186 | -0.065 |
| DEP–toAP | -0.106 | 0.015 | -0.043 | 0.067 |
| AP–GB | 0.546 | -0.158 | -0.209 | -0.102 |
| AP–toAP | -0.049 | 0.017 | -0.045 | 0.071 |
| GB–toAP | -0.047 | -0.013 | -0.081 | 0.042 |

Table E.1: Posterior residual summaries for selected terminals under the baseline rank-one model.

| Terminal | $N_i$ | Mean residual | Median residual | Positive residuals (%) |
|---|---|---|---|---|
| Rasgas LNG Terminal 3 | 59 | 3.134 | 3.119 | 100.0 |
| Fos Tonkin LNG Terminal | 50 | 1.426 | 1.451 | 100.0 |
| Yamal LNG Terminal | 59 | 0.958 | 0.723 | 96.6 |
| Sabine Pass LNG Terminal | 59 | 0.932 | 0.859 | 94.9 |
| Pluto LNG Terminal | 59 | 0.871 | 0.853 | 91.5 |
| Gate LNG Terminal | 59 | 0.706 | 0.776 | 89.8 |
| Corpus Christi LNG Terminal | 51 | 0.394 | 0.463 | 86.3 |
| Cameroon FLNG Terminal | 24 | -1.547 | -1.594 | 0.0 |
| Sakhalin II LNG Terminal | 21 | -1.459 | -1.645 | 0.0 |
| Punta Europa LNG Terminal | 20 | -1.234 | -1.455 | 0.0 |
| Elba Island LNG Terminal | 25 | -1.024 | -0.970 | 0.0 |
| Tiga Malaysia LNG Terminal | 30 | -0.931 | -0.817 | 3.3 |
| Oman Qalhat LNG Terminal | 25 | -1.553 | -1.805 | 4.0 |
| Queensland Curtis LNG Terminal | 47 | -0.928 | -0.911 | 6.4 |
| Senboku 2 LNG Terminal | 39 | -0.652 | -0.751 | 7.7 |
| Incheon LNG Terminal | 53 | -0.606 | -0.622 | 9.4 |
| Prelude FLNG Terminal | 24 | -1.002 | -1.001 | 12.5 |
| Wheatstone LNG Terminal | 50 | -0.542 | -0.537 | 14.0 |

*Notes:* $N_i$ denotes the number of terminal-month observations available for terminal $i$ in the estimation sample. Mean and median residuals are calculated across the posterior mean residuals $\widehat{z}_{it}$ for each terminal. Positive residuals (%) denotes the percentage of observations for which $\widehat{z}_{it} > 0$. The table includes terminals with at least 20 observations and a proportion of positive posterior mean residuals either above 85% or below 15%.

### E.1.2 Chokepoints

#### Chokepoints residuals heterogeneity

This section provides additional descriptive evidence of persistent chokepoint-level residual patterns under the baseline rank-one specification, complementing the discussion in the main text. The table includes chokepoints with at least 20 observations and positive posterior mean residuals in either more than 85% or less than 15% of their observed months.

Table E.4: Posterior residual summaries for selected chokepoints under the baseline rank-one model.

| Chokepoint | $N_i$ | Mean residual | Median residual | Positive residuals (%) |
|---|---|---|---|---|
| Gibraltar Strait | 59 | 1.294 | 1.301 | 96.6 |
| Taiwan Strait | 59 | 1.129 | 1.248 | 96.6 |
| Suez Canal | 59 | 1.066 | 1.225 | 96.6 |
| Malacca Strait | 59 | 0.980 | 1.180 | 91.5 |
| Bab el-Mandeb Strait | 59 | 0.915 | 1.089 | 89.8 |
| Strait of Hormuz | 59 | 0.771 | 0.767 | 98.3 |
| Lombok Strait | 59 | -0.409 | -0.438 | 11.9 |
| Makassar Strait | 59 | -0.441 | -0.420 | 5.1 |
| Bohai Strait | 52 | -0.564 | -0.513 | 3.8 |
| Ombai Strait | 59 | -0.720 | -0.763 | 3.4 |
| Tsugaru Strait | 56 | -0.793 | -0.789 | 7.1 |
| Cape of Good Hope | 57 | -1.083 | -1.130 | 8.8 |
| Sunda Strait | 49 | -1.179 | -1.254 | 4.1 |

*Notes:* $N_i$ denotes the number of chokepoint-month observations available for chokepoint $i$ in the estimation sample. Mean and median residuals are calculated across the posterior mean residuals $\widehat{z}_{it}$ for each chokepoint. Positive residuals (%) denotes the percentage of observations for which $\widehat{z}_{it} > 0$. The table reports chokepoints with at least 20 observations and more than 85% or less than 15% positive posterior mean residuals.

## E.2 Node-intercept rank-one specification

### E.2.1 Terminals

This subsection reports the complete marginal and joint posterior predictive checks for the terminal random-intercept rank-one specification. The checks use 200 conditional in-sample replicated datasets, as described in Section 6.2.1, and the reported intervals are the 2.5th and 97.5th percentiles of the replicated statistics. For the marginal checks, continuous responses are additionally examined separately for importers and exporters, using the lagged exporter indicator.

Table E.5: Marginal posterior predictive checks for terminal responses under the node-intercept rank-one model. All statistics are on the original response scale; intervals are based on 200 conditional in-sample replicates.

| Response | Statistic | Observed | Rep. mean | 95% lower | 95% upper |
|---|---|---|---|---|---|
| *All terminals* | | | | | |
| PageRank | Mean | 0.0185 | 0.0185 | 0.0184 | 0.0187 |
| PageRank | SD | 0.0133 | 0.0129 | 0.0126 | 0.0133 |
| PageRank | 5th percentile | 0.0089 | 0.0078 | 0.0075 | 0.0080 |
| PageRank | Median | 0.0147 | 0.0153 | 0.0151 | 0.0155 |
| PageRank | 95th percentile | 0.0391 | 0.0392 | 0.0381 | 0.0404 |
| HHI | Mean | 0.5887 | 0.5893 | 0.5804 | 0.5984 |
| HHI | SD | 0.3254 | 0.3229 | 0.3184 | 0.3268 |
| HHI | 5th percentile | 0.1268 | 0.1312 | 0.1205 | 0.1430 |
| HHI | Median | 0.5045 | 0.4974 | 0.4866 | 0.5092 |
| HHI | 95th percentile | 1.0000 | 1.0000 | 1.0000 | 1.0000 |
| HHI | Proportion equal to one | 0.3409 | 0.3403 | 0.3243 | 0.3522 |
| Dependence | Mean | 0.4948 | 0.4962 | 0.4834 | 0.5099 |
| Dependence | SD | 0.3219 | 0.3185 | 0.3120 | 0.3247 |
| Dependence | 5th percentile | 0.0716 | 0.0816 | 0.0733 | 0.0897 |
| Dependence | Median | 0.4283 | 0.4161 | 0.4012 | 0.4323 |
| Dependence | 95th percentile | 1.0000 | 1.0000 | 1.0000 | 1.0000 |
| Dependence | Proportion equal to one | 0.2076 | 0.2086 | 0.1919 | 0.2259 |
| AP | Proportion | 0.1057 | 0.1064 | 0.0929 | 0.1196 |
| GB | Proportion | 0.5850 | 0.5851 | 0.5670 | 0.6085 |
| toAP | Proportion | 0.9456 | 0.9457 | 0.9354 | 0.9551 |
| *Lagged importers* | | | | | |
| PageRank | Mean | 0.0145 | 0.0144 | 0.0143 | 0.0146 |
| PageRank | SD | 0.0045 | 0.0050 | 0.0048 | 0.0052 |
| PageRank | 5th percentile | 0.0089 | 0.0074 | 0.0071 | 0.0077 |
| PageRank | Median | 0.0136 | 0.0138 | 0.0136 | 0.0141 |
| PageRank | 95th percentile | 0.0233 | 0.0236 | 0.0230 | 0.0241 |
| HHI | Mean | 0.6745 | 0.6775 | 0.6667 | 0.6880 |
| HHI | SD | 0.2966 | 0.2963 | 0.2925 | 0.3005 |
| HHI | 5th percentile | 0.2446 | 0.2534 | 0.2422 | 0.2648 |
| HHI | Median | 0.5701 | 0.5953 | 0.5731 | 0.6184 |
| HHI | 95th percentile | 1.0000 | 1.0000 | 1.0000 | 1.0000 |
| HHI | Proportion equal to one | 0.4189 | 0.4270 | 0.4090 | 0.4439 |
| Dependence | Mean | 0.4812 | 0.4761 | 0.4596 | 0.4943 |
| Dependence | SD | 0.3574 | 0.3319 | 0.3238 | 0.3399 |
| Dependence | 5th percentile | 0.0614 | 0.0721 | 0.0628 | 0.0816 |
| Dependence | Median | 0.3472 | 0.3728 | 0.3567 | 0.3920 |

*Continued on next page*

*Table continued from previous page*

| Response | Statistic | Observed | Rep. mean | 95% lower | 95% upper |
|---|---|---|---|---|---|
| Dependence | 95th percentile | 1.0000 | 1.0000 | 1.0000 | 1.0000 |
| Dependence | Proportion equal to one | 0.2645 | 0.2390 | 0.2185 | 0.2602 |
| *Lagged exporters* | | | | | |
| PageRank | Mean | 0.0269 | 0.0270 | 0.0266 | 0.0274 |
| PageRank | SD | 0.0198 | 0.0188 | 0.0182 | 0.0194 |
| PageRank | 5th percentile | 0.0088 | 0.0092 | 0.0087 | 0.0097 |
| PageRank | Median | 0.0210 | 0.0220 | 0.0214 | 0.0225 |
| PageRank | 95th percentile | 0.0714 | 0.0669 | 0.0629 | 0.0719 |
| HHI | Mean | 0.4119 | 0.4075 | 0.3956 | 0.4177 |
| HHI | SD | 0.3103 | 0.2986 | 0.2882 | 0.3079 |
| HHI | 5th percentile | 0.0877 | 0.0564 | 0.0457 | 0.0688 |
| HHI | Median | 0.3333 | 0.3268 | 0.3151 | 0.3413 |
| HHI | 95th percentile | 1.0000 | 1.0000 | 1.0000 | 1.0000 |
| HHI | Proportion equal to one | 0.1800 | 0.1614 | 0.1445 | 0.1765 |
| Dependence | Mean | 0.5228 | 0.5376 | 0.5206 | 0.5533 |
| Dependence | SD | 0.2296 | 0.2842 | 0.2758 | 0.2915 |
| Dependence | 5th percentile | 0.1852 | 0.1176 | 0.1010 | 0.1332 |
| Dependence | Median | 0.4972 | 0.5033 | 0.4799 | 0.5260 |
| Dependence | 95th percentile | 1.0000 | 1.0000 | 1.0000 | 1.0000 |
| Dependence | Proportion equal to one | 0.0904 | 0.1458 | 0.1232 | 0.1658 |

Table E.6: Joint posterior predictive checks for terminal responses under the node-intercept rank-one model. Entries are Spearman rank correlations; replicated intervals are 95% equal-tailed intervals based on 200 conditional in-sample replicates.

| Response pair | Subset | Observed | Rep. median | 95% lower | 95% upper |
|---|---|---|---|---|---|
| *All terminals* | | | | | |
| PageRank–HHI | All | -0.6394 | -0.6858 | -0.7041 | -0.6641 |
| PageRank–Dependence | All | 0.3145 | 0.1528 | 0.1115 | 0.1904 |
| PageRank–AP | All | 0.3724 | 0.2479 | 0.2135 | 0.2873 |
| PageRank–GB | All | -0.2380 | -0.2512 | -0.2954 | -0.2122 |
| PageRank–toAP | All | -0.0150 | 0.0385 | 0.0011 | 0.0753 |
| HHI–Dependence | All | -0.1707 | -0.1490 | -0.1853 | -0.1022 |
| HHI–AP | All | -0.2904 | -0.2387 | -0.2764 | -0.1977 |
| HHI–GB | All | 0.5326 | 0.2420 | 0.2040 | 0.2888 |
| HHI–toAP | All | -0.0376 | -0.0370 | -0.0716 | -0.0017 |
| Dependence–AP | All | 0.1566 | 0.0796 | 0.0463 | 0.1129 |
| Dependence–GB | All | -0.0664 | -0.0637 | -0.0974 | -0.0270 |

*Continued on next page*

*Table continued from previous page*

| Response pair | Subset | Observed | Rep. median | 95% lower | 95% upper |
|---|---|---|---|---|---|
| Dependence–toAP | All | -0.0815 | 0.0090 | -0.0241 | 0.0437 |
| AP–GB | All | 0.2896 | -0.1078 | -0.1426 | -0.0696 |
| AP–toAP | All | -0.0628 | 0.0178 | -0.0150 | 0.0420 |
| GB–toAP | All | 0.0595 | -0.0156 | -0.0518 | 0.0233 |
| *Lagged importers* | | | | | |
| PageRank–HHI | Importer | -0.4163 | -0.5153 | -0.5496 | -0.4802 |
| PageRank–Dependence | Importer | 0.2088 | 0.0576 | 0.0111 | 0.0986 |
| PageRank–AP | Importer | 0.2320 | 0.0896 | 0.0467 | 0.1231 |
| PageRank–GB | Importer | -0.2576 | -0.1304 | -0.1744 | -0.0850 |
| PageRank–toAP | Importer | -0.0425 | 0.0196 | -0.0226 | 0.0596 |
| HHI–Dependence | Importer | -0.1625 | -0.0593 | -0.0996 | -0.0150 |
| HHI–AP | Importer | -0.1374 | -0.0896 | -0.1247 | -0.0396 |
| HHI–GB | Importer | 0.6437 | 0.1336 | 0.0864 | 0.1857 |
| HHI–toAP | Importer | 0.0136 | -0.0198 | -0.0598 | 0.0255 |
| Dependence–AP | Importer | 0.0331 | 0.0114 | -0.0302 | 0.0512 |
| Dependence–GB | Importer | -0.1541 | -0.0150 | -0.0554 | 0.0236 |
| Dependence–toAP | Importer | -0.0447 | 0.0018 | -0.0362 | 0.0475 |
| AP–GB | Importer | 0.1368 | -0.0217 | -0.0637 | 0.0150 |
| AP–toAP | Importer | 0.0193 | 0.0077 | -0.0375 | 0.0400 |
| GB–toAP | Importer | 0.1314 | -0.0068 | -0.0494 | 0.0313 |
| *Lagged exporters* | | | | | |
| PageRank–HHI | Exporter | -0.8178 | -0.7917 | -0.8137 | -0.7681 |
| PageRank–Dependence | Exporter | 0.5164 | 0.2822 | 0.2223 | 0.3538 |
| PageRank–AP | Exporter | 0.2764 | 0.3608 | 0.2864 | 0.4116 |
| PageRank–GB | Exporter | -0.1978 | -0.3406 | -0.4056 | -0.2838 |
| PageRank–toAP | Exporter | 0.1661 | 0.0555 | -0.0097 | 0.1130 |
| HHI–Dependence | Exporter | -0.1019 | -0.2515 | -0.3193 | -0.1852 |
| HHI–AP | Exporter | -0.1593 | -0.3272 | -0.3788 | -0.2574 |
| HHI–GB | Exporter | 0.3984 | 0.3096 | 0.2513 | 0.3783 |
| HHI–toAP | Exporter | -0.2304 | -0.0539 | -0.1079 | 0.0132 |
| Dependence–AP | Exporter | 0.3066 | 0.1400 | 0.0856 | 0.1866 |
| Dependence–GB | Exporter | 0.1946 | -0.1110 | -0.1647 | -0.0594 |
| Dependence–toAP | Exporter | -0.1061 | 0.0203 | -0.0355 | 0.0849 |
| AP–GB | Exporter | 0.5461 | -0.1497 | -0.1977 | -0.0832 |
| AP–toAP | Exporter | -0.0495 | 0.0236 | -0.0322 | 0.0728 |
| GB–toAP | Exporter | -0.0467 | -0.0203 | -0.0862 | 0.0338 |

*Notes.* AP denotes articulation-point status; GB denotes incident-global-bridge status; toAP denotes connection to an articulation point. Binary-response means are proportions.

### E.2.2 Chokepoints

Tables E.7 and E.8 report the complete marginal and pairwise joint posterior predictive checks for the chokepoint random-intercept rank-one model. Results are based on 200 conditional in-sample posterior predictive replicates.

Table E.7: Marginal posterior predictive checks for chokepoint responses under the random-intercept rank-one model. All statistics are on the original response scale; 95% intervals are based on 200 conditional in-sample replicates.

| Response | Statistic | Observed | Rep. mean | 95% lower | 95% upper |
|---|---|---|---|---|---|
| PageRank | Mean | 0.0176 | 0.0177 | 0.0173 | 0.0181 |
| PageRank | SD | 0.0069 | 0.0065 | 0.0061 | 0.0069 |
| PageRank | 5th percentile | 0.0084 | 0.0083 | 0.0077 | 0.0089 |
| PageRank | Median | 0.0166 | 0.0169 | 0.0164 | 0.0174 |
| PageRank | 95th percentile | 0.0302 | 0.0295 | 0.0286 | 0.0306 |
| Betweenness | Mean | 0.0132 | 0.0132 | 0.0126 | 0.0140 |
| Betweenness | SD | 0.0135 | 0.0121 | 0.0113 | 0.0130 |
| Betweenness | 5th percentile | 0.0006 | 0.0006 | 0.0004 | 0.0008 |
| Betweenness | Median | 0.0077 | 0.0097 | 0.0090 | 0.0106 |
| Betweenness | 95th percentile | 0.0413 | 0.0376 | 0.0348 | 0.0402 |
| Betweenness | Proportion equal to zero | 0.0052 | 0.0058 | 0.0009 | 0.0129 |
| Flow share | Mean | 0.0234 | 0.0234 | 0.0230 | 0.0238 |
| Flow share | SD | 0.0201 | 0.0200 | 0.0195 | 0.0206 |
| Flow share | 5th percentile | 0.0030 | 0.0035 | 0.0031 | 0.0038 |
| Flow share | Median | 0.0159 | 0.0157 | 0.0150 | 0.0164 |
| Flow share | 95th percentile | 0.0643 | 0.0641 | 0.0618 | 0.0662 |
| Articulation point (AP) | Proportion | 0.7360 | 0.7356 | 0.7008 | 0.7756 |
| Connected to AP | Proportion | 0.8960 | 0.8960 | 0.8761 | 0.9157 |

Table E.8: Joint posterior predictive checks for all pairwise Spearman correlations between chokepoint responses under the random-intercept rank-one model. Intervals are 95% equal-tailed posterior predictive intervals based on 200 conditional in-sample replicates.

| Response pair | Observed | Rep. median | 95% lower | 95% upper |
|---|---|---|---|---|
| PageRank–Betweenness | 0.6846 | 0.4731 | 0.4168 | 0.5162 |
| PageRank–Flow share | 0.6718 | 0.6090 | 0.5618 | 0.6444 |
| PageRank–Articulation point | 0.2867 | 0.2014 | 0.1402 | 0.2707 |
| PageRank–Connected to AP | 0.2157 | 0.1359 | 0.0777 | 0.1948 |
| Betweenness–Flow share | 0.8000 | 0.7005 | 0.6622 | 0.7364 |
| Betweenness–Articulation point | 0.2521 | 0.2306 | 0.1647 | 0.2952 |

*Continued on next page*

*Table continued from previous page*

| Response pair | Observed | Rep. median | 95% lower | 95% upper |
|---|---|---|---|---|
| Betweenness–Connected to AP | 0.3204 | 0.1558 | 0.0937 | 0.2125 |
| Flow share–Articulation point | 0.2910 | 0.3001 | 0.2257 | 0.3609 |
| Flow share–Connected to AP | 0.1818 | 0.1999 | 0.1368 | 0.2654 |
| Articulation point–Connected to AP | 0.1665 | 0.0701 | 0.0103 | 0.1435 |

# Appendix F

# Model diagnostics

## F.1 Prior sensitivity analysis

### F.1.1 Prior predictive checks

Table F.1 reports the observed proportions of observations equal to one for the two one-inflated terminal responses, together with the corresponding mean proportions across prior predictive replications under the three prior specifications.

Table F.1: Observed and mean prior predictive proportions of values equal to one for terminal responses under the three prior specifications.

| Response | Observed | ORG | CON | RS |
|---|---|---|---|---|
| Counterparty HHI | 0.3409 | 0.5063 | 0.4922 | 0.4920 |
| Role-specific dependence | 0.2076 | 0.4960 | 0.5042 | 0.5039 |

### F.1.2 MCMC sampling diagnostics

Table F.2 reports the maximum rank-normalised $\widehat{R}$ and minimum bulk and tail effective sample sizes for the original (ORG) and response-specific precision (RS) prior specifications of the rank-one models. All four fits were obtained using four chains, each with 1,500 burn-in iterations and 2,000 retained draws, providing comparable sampling configurations.

For terminals, the minimum bulk ESS was associated with the lagged exporter coefficient under both specifications, while the minimum tail ESS concerned the HHI boundary loading under ORG and a latent factor mean under RS. For chokepoints, the betweenness precision parameter exhibited the lowest bulk ESS under both specifications, decreasing from 472 under ORG to 359 under RS.

The maximum reported $\widehat{R}$ was 1.01 across all four fits, indicating no substantial between-chain discrepancies according to this diagnostic. Nevertheless, the lower effective sample sizes for selected parameters, particularly the betweenness precision under

RS, indicate reduced Monte Carlo precision for those posterior estimates.

Table F.2: MCMC convergence diagnostics for the original (ORG) and response-specific precision (RS) prior specifications of the rank-one models.

| Node type | Prior | Diagnostic | Value | Parameter |
|---|---|---|---|---|
| Terminal | ORG | Max $\widehat{R}$ | 1.01 | Multiple parameters |
| | | Min bulk ESS | 817 | $\beta_{\text{exporter}}$ |
| | | Min tail ESS | 1958 | HHI boundary loading |
| | RS | Max $\widehat{R}$ | 1.01 | HHI interior intercept |
| | | Min bulk ESS | 782 | $\beta_{\text{exporter}}$ |
| | | Min tail ESS | 2037 | Factor mean (obs. 1927) |
| Chokepoint | ORG | Max $\widehat{R}$ | 1.01 | Betweenness loading |
| | | Min bulk ESS | 472 | Betweenness precision |
| | | Min tail ESS | 905 | Betweenness precision |
| | RS | Max $\widehat{R}$ | 1.01 | Multiple parameters |
| | | Min bulk ESS | 359 | Betweenness precision |
| | | Min tail ESS | 780 | Betweenness intercept |

*Note:* All models were estimated using four chains, with 1,500 warm-up iterations and 2,000 retained draws per chain. Maximum $\widehat{R}$ values are reported to two decimal places.

## F.2 Rank-one MCMC convergence diagnostics

### F.2.1 R-hat and Effective Sample Sizes

Table F.3 reports the maximum rank-normalised $\widehat{R}$ and minimum bulk and tail effective sample sizes for the baseline rank-one terminal and chokepoint models, together with the parameters associated with the minimum ESS values. For terminals, the minimum bulk ESS was associated with the lagged exporter coefficient, while the minimum tail ESS concerned the loading governing the probability of counterparty HHI equal to one. For chokepoints, both minimum ESS values were associated with the betweenness precision parameter, indicating comparatively less efficient sampling for this parameter.

### F.2.2 Geweke diagnostic

The Geweke diagnostic compares estimates obtained from the first 10% and last 50% of each chain. The reference criterion was $|Z| \leq 1.96$, corresponding to a two-sided significance level of 0.05. For terminals, 98 out of 104 parameter-chain tests (94.2%) satisfied this criterion, while 6 tests (5.8%) exceeded the reference value. For chokepoints, 73 out of 76 tests (96.1%) satisfied the criterion, with three exceedances (3.9%).

Table F.3: Detailed MCMC convergence and mixing diagnostics for the baseline rank-one terminal and chokepoint models.

| Node type | Diagnostic | Value | Parameter |
|---|---|---|---|
| Terminal | Maximum $\widehat{R}$ | 1.001 | Multiple parameters |
| | Minimum bulk ESS | 8,602 | Lagged exporter coefficient |
| | Minimum tail ESS | 20,487 | HHI boundary loading |
| Chokepoint | Maximum $\widehat{R}$ | 1.001 | Multiple parameters |
| | Minimum bulk ESS | 4,808 | Betweenness precision |
| | Minimum tail ESS | 9,772 | Betweenness precision |

*Note:* Both models were estimated using four chains, each with 20,000 burn-in iterations and 30,000 retained draws. Maximum $\widehat{R}$ values are reported to three decimal places.

Table F.4 reports the parameter-chain combinations associated with these exceedances.

Table F.4: Parameter-chain combinations with absolute Geweke Z-scores exceeding 1.96 in the baseline rank-one models.

| Node type | Parameter | Chain | $Z$ |
|---|---|---|---|
| Terminal | Neighbour activity coefficient | 1 | 2.400 |
| | Voyage distance coefficient | 2 | 2.589 |
| | HHI interior loading | 1 | -2.013 |
| | HHI interior intercept | 1 | 1.988 |
| | HHI precision | 1 | 2.132 |
| | Dependence precision | 1 | -2.363 |
| Chokepoint | Betweenness interior intercept | 1 | 2.308 |
| | Articulation-point intercept | 1 | 2.145 |
| | PageRank precision | 2 | -2.472 |

### F.2.3 Heidelberger–Welch diagnostic

The Heidelberger–Welch procedure assesses stationarity and the Monte Carlo precision of posterior mean estimates. All 104 terminal and 76 chokepoint parameter-chain tests satisfied the stationarity criterion. For terminals, 103 tests passed without excluding any additional draws, while one required the exclusion of 6,000 initial draws. For chokepoints, 72 tests passed without additional exclusions, while the remaining four required the exclusion of between 3,000 and 15,000 draws. Table F.5 reports the parameter-chain combinations requiring additional exclusions.

The half-width test assesses the precision of posterior mean estimates by requiring the half-width of the 95% Monte Carlo confidence interval to be less than 10% of the absolute estimated mean. All 76 chokepoint tests satisfied this criterion. For

Table F.5: Parameter-chain combinations requiring additional initial-draw exclusions in the Heidelberger–Welch stationarity diagnostic.

| Node type | Parameter | Chain | Excluded draws |
|---|---|---|---|
| Terminal | Articulation-point binary loading | 4 | 6,000 |
| Chokepoint | Voyage distance coefficient | 2 | 6,000 |
| | Betweenness interior loading | 4 | 15,000 |
| | Flow-share interior loading | 4 | 15,000 |
| | Flow-share interior intercept | 1 | 3,000 |

terminals, 100 out of 104 tests passed, while the remaining four exceedances were associated with the counterparty HHI interior intercept (half-width values for each chain were respectively 0.375, 0.512, 0.578, 0.562).

### F.2.4 Trace plots

The following figures present posterior densities and trace plots for the global parameters of the baseline terminal and chokepoint models, estimated using four chains with 20,000 warm-up iterations and 30,000 retained draws per chain. For both models, the posterior density estimates of the four chains exhibit substantial overlap, suggesting that the chains explored similar regions of the parameter space. The corresponding trace plots show fluctuations around relatively stable levels, without visible systematic trends or persistent shifts over the retained iterations. These patterns are observed across regression coefficients, response loadings, intercepts, and precision parameters, including those associated with comparatively lower effective sample sizes. Overall, the visual diagnostics are consistent with satisfactory mixing and support the convergence assessment based on the rank-normalised $\widehat{R}$ and effective sample size statistics.




Figure 12: Posterior densities and trace plots for terminal regression coefficients (part 1).




Figure 13: Posterior densities and trace plots for terminal regression coefficients and for the PageRank response loading (part 2).




Figure 14: Posterior densities and trace plots for terminal response loadings (part 3).




Figure 15: Posterior densities and trace plots for terminal response loadings and intercepts (part 4).




Figure 16: Posterior densities and trace plots for terminal response intercepts and the PageRank precision parameter (part 5).




Figure 17: Posterior densities and trace plots for terminal precision parameters and one-mass intercepts (part 6).




Figure 18: Posterior densities and trace plots for terminal one-mass loadings (part 7).




Figure 19: Posterior densities and trace plots for chokepoint regression coefficients (part 1).




Figure 20: Posterior densities and trace plots for chokepoint response loadings (part 2).




Figure 21: Posterior densities and trace plots for chokepoint response loadings and intercepts (part 3).




Figure 22: Posterior densities and trace plots for chokepoint intercepts and precision parameters (part 4).




Figure 23: Posterior densities and trace plots for the remaining chokepoint precision and zero-mass parameters (part 5).

### F.2.5 Autocorrelation diagnostics

The figures presented in this section show within-chain autocorrelation functions for selected global parameters from the baseline models estimated using four chains, each with 20,000 burn-in iterations and 30,000 retained draws. Parameters were selected based on the lowest bulk ESS among regression coefficients and response loadings, the lowest bulk and tail ESS among all global parameters, and the PageRank anchor loading. For terminals, the selected parameters are the lagged exporter coefficient, the PageRank loading, and the loading for the probability of HHI equal to one. All three exhibit a progressive decline in autocorrelation, approaching zero within approximately 30–40 lags. The lagged exporter coefficient shows slightly more persistent autocorrelation, while the PageRank and HHI loadings follow similar decay patterns across chains. The minimum bulk ESS among these parameters is 8,602, associated with the lagged exporter coefficient. For chokepoints, the selected parameters are the lagged neighbour-activity coefficient, the betweenness loading and precision, and the PageRank loading. The lagged neighbour-activity coefficient exhibits a rapid autocorrelation decay, reaching values close to zero within the first few lags. The PageRank loading also shows a relatively rapid decline, whereas the betweenness parameters exhibit more persistent autocorrelation, particularly the precision parameter, whose autocorrelation approaches zero around lag 60–70. The latter also has the lowest bulk ESS among global parameters (4,808).

Overall, the autocorrelation functions show consistent decay patterns across chains, with no substantial persistent autocorrelation at longer lags. The slower decay observed for some parameters is reflected in their comparatively lower, but still substantial, effective sample sizes.

## F.3 Random-intercept rank-one MCMC convergence diagnostics

This subsection provides additional convergence and mixing diagnostics for the random-intercept rank-one specifications complementing the discussion in Chapter 6. The tables report rank-normalised $\widehat{R}$ statistics and bulk and tail effective sample sizes (ESS) for selected global parameters and node-specific intercepts. For the terminal model, the reported global parameters include those exhibiting the least favourable mixing diagnostics, together with the standard deviation of the node-specific intercepts. For the chokepoint model, particular attention is given to the response intercepts and node-specific effects, for which convergence difficulties were identified.

The additional diagnostics confirm substantial differences in sampling performance between the two specifications. For terminals, the least favourable diagnostics are associated primarily with response intercepts, while the node-specific effects exhibit comparatively satisfactory mixing. For chokepoints, the continuous-response and several




Figure 24: Within-chain autocorrelation functions for the lagged exporter coefficient, counterparty HHI (mass at 1) and PageRank loadings in the baseline rank-one terminal model.




Figure 25: Within-chain autocorrelation functions for the lagged neighbour activity coefficient, betweenness loading, Beta precision and PageRank loading in the baseline rank-one chokepoint model.

node-specific intercepts exhibit low effective sample sizes, whereas the node-effect standard deviation exhibits substantially better mixing diagnostics. These results indicate that the posterior summaries of individual chokepoint effects should be interpreted cautiously, as the available samples may not adequately characterise their posterior distributions.

Table F.6: Selected global-parameter sampling diagnostics for the terminal random-intercept model.

| Parameter | $\widehat{R}$ | Bulk ESS | Tail ESS |
|---|---|---|---|
| PageRank intercept | 1.003 | 795 | 1,734 |
| HHI interior-mean intercept | 1.003 | 883 | 2,031 |
| HHI boundary-component intercept | 1.003 | 802 | 1,855 |
| Dependence interior-mean intercept | 1.002 | 963 | 2,426 |
| Global-bridge binary intercept | 1.002 | 1,378 | 4,098 |
| Dependence boundary-component intercept | 1.002 | 1,433 | 4,221 |
| Node-effect standard deviation ($\sigma_\mu$) | 1.000 | 14,489 | 32,125 |

Table F.7: Selected global-parameter sampling diagnostics for the chokepoint random-intercept model.

| Parameter | $\widehat{R}$ | Bulk ESS | Tail ESS |
|---|---|---|---|
| PageRank intercept | 1.039 | 114 | 205 |
| Betweenness intercept | 1.040 | 113 | 202 |
| Flow-share intercept | 1.039 | 112 | 206 |
| Node-effect standard deviation ($\sigma_\mu$) | 1.001 | 2,801 | 11,759 |

Table F.8: Sampling diagnostics for the 24 node-specific intercepts in the chokepoint random-intercept model.

| Node | $\widehat{R}$ | Bulk ESS | Tail ESS |
|---|---|---|---|
| CP001 | 1.038 | 118 | 231 |
| CP002 | 1.036 | 122 | 241 |
| CP004 | 1.038 | 118 | 225 |
| CP005 | 1.037 | 118 | 219 |
| CP006 | 1.037 | 118 | 225 |
| CP007 | 1.032 | 138 | 304 |
| CP008 | 1.037 | 119 | 227 |
| CP009 | 1.037 | 119 | 227 |
| CP010 | 1.035 | 128 | 274 |
| CP011 | 1.038 | 118 | 229 |
| CP012 | 1.036 | 124 | 254 |
| CP013 | 1.034 | 130 | 282 |
| CP014 | 1.035 | 127 | 269 |
| CP015 | 1.033 | 132 | 290 |
| CP016 | 1.033 | 130 | 287 |
| CP017 | 1.035 | 128 | 277 |
| CP018 | 1.017 | 272 | 835 |
| CP019 | 1.031 | 144 | 331 |
| CP020 | 1.033 | 133 | 297 |
| CP021 | 1.006 | 763 | 3,297 |
| CP022 | 1.035 | 125 | 254 |
| CP023 | 1.015 | 312 | 1,006 |
| CP024 | 1.006 | 767 | 3,164 |
| CP027 | 1.036 | 124 | 253 |




Figure 26: Selected autocorrelation functions for the terminal random-intercept rank-one model. The upper plot corresponds to the lagged exporter regression coefficient, while the lower plot shows the PageRank continuous-response intercept. The four curves in each plot represent the individual MCMC chains.




Figure 27: Posterior density estimates and trace plots for selected factor-regression coefficients in the terminal random-intercept rank-one model. Different colours represent the four MCMC chains.




Figure 28: Posterior density estimates and trace plots for selected response-specific parameters in the terminal random-intercept rank-one model, including the connected-to-articulation-point loading and continuous-response intercepts.




Figure 29: Selected autocorrelation functions for the chokepoint random-intercept rank-one model. From top to bottom, the plots correspond to the monthly flow-share loading, the monthly flow-share intercept, and the betweenness centrality intercept. The four curves in each plot represent the individual MCMC chains. The response intercepts exhibit particularly persistent autocorrelation, indicating slow mixing.




Figure 30: Posterior density estimates and trace plots for the four factor-regression coefficients in the chokepoint random-intercept rank-one model.




Figure 31: Posterior density estimates and trace plots for the PageRank, betweenness centrality, and monthly flow-share intercepts in the chokepoint random-intercept rank-one model. The traces illustrate persistent fluctuations and slow mixing.

# Bibliography


Abril-Pla, O., Andreani, V., Carroll, C., Dong, L., Fonnesbeck, C. J., et al. (2023). Pymc: A modern, and comprehensive probabilistic programming framework in python. *PeerJ Computer Science*, 9:e1516.

Betancourt, M. (2017). A conceptual introduction to hamiltonian monte carlo.

Brandes, U. (2001). A faster algorithm for betweenness centrality. *Journal of Mathematical Sociology*, 25(2):163–177.

Brin, S. and Page, L. (1998). The anatomy of a large-scale hypertextual web search engine. *Computer Networks and ISDN Systems*, 30(1–7):107–117.

Dijkstra, E. W. (1959). A note on two problems in connexion with graphs. *Numerische Mathematik*, 1:269–271.

Eurostat (2021). SeaRoute. Computer software. Available at: `https://github.com/eurostat/searoute`. Accessed: 2026-08-16.

Flanders Marine Institute (n.d.). Marine Regions. Online geospatial database. Available at: `https://www.marineregions.org/`. Accessed: 2026-08-16.

Gelman, A., Carlin, J. B., Stern, H. S., Dunson, D. B., Vehtari, A., and Rubin, D. B. (2013). *Bayesian Data Analysis*. Chapman & Hall/CRC, Boca Raton, FL, 3 edition.

Geweke, J. (1992). Evaluating the accuracy of sampling-based approaches to the calculation of posterior moments. In Bernardo, J. M., Berger, J. O., Dawid, A. P., and Smith, A. F. M., editors, *Bayesian Statistics 4*, pages 169–193. Clarendon Press, Oxford.

Heidelberger, P. and Welch, P. D. (1983). Simulation run length control in the presence of an initial transient. *Operations Research*, 31(6):1109–1144.

Hirschman, A. O. (1945). *National Power and the Structure of Foreign Trade*. University of California Press, Berkeley.

Hoffman, M. D. and Gelman, A. (2014). The no-u-turn sampler: Adaptively setting path lengths in hamiltonian monte carlo. *Journal of Machine Learning Research*, 15(47):1593–1623.

International Maritime Organization (1998). Adoption, Designation and Substitution of Archipelagic Sea Lanes. Resolution MSC.72(69), International Maritime Organization. Adopted 19 May 1998, MSC 69/22/Add.1, Annex 9. Available at: https://navcen.uscg.gov/sites/default/files/pdf/marcomms/imo/msc_resolutions/MSC69-22a1-9.pdf. Accessed: 2026-08-16.

International Maritime Organization (2014). Routeing Measures Other Than Traffic Separation Schemes. SN Circular SN.1/Circ.326, International Maritime Organization. Issued 23 May 2014. Available at: https://mantamaritime.com/downloads/flag_news/sn.1_circ.326.pdf. Accessed: 2026-08-16.

International Maritime Organization (2019). New Traffic Separation Schemes. COLREG Circular COLREG.2/Circ.74, International Maritime Organization. Issued 14 June 2019. Available at: https://www.bimco.org/media/gusfhkrc/colreg2circ74-new-traffic-separation-schemes-secretariat.pdf. Accessed: 2026-08-16.

International Monetary Fund (2026). PortWatch Chokepoints Database. Available at: https://services9.arcgis.com/weJ1QsnbMYJlCHdG/ArcGIS/rest/services/PortWatch_chokepoints_database/FeatureServer/0. Accessed: 2026-09-04.

National Centers for Environmental Information (n.d.). Report for H01600A: Yucatan Channel. Online hydrographic survey record. Available at: https://www.ngdc.noaa.gov/nos/H00001-H02000/H01600A.html. National Oceanic and Atmospheric Administration. Accessed: 2026-08-16.

National Geospatial-Intelligence Agency (n.d.). NGA GeoNames: Bohai Strait. Online geographic database. Record UFI -1899028. Available at: https://geonames.nga.mil/geon-ags/rest/services/RESEARCH/GIS_OUTPUT/MapServer/0. Accessed: 2026-08-16.

Neal, R. M. (2011). Mcmc using hamiltonian dynamics. In Brooks, S., Gelman, A., Jones, G. L., and Meng, X.-L., editors, *Handbook of Markov Chain Monte Carlo*, pages 113–162. Chapman and Hall/CRC.

Page, L., Brin, S., Motwani, R., and Winograd, T. (1999). The PageRank citation ranking: Bringing order to the web. Technical Report 1999-66, Stanford InfoLab. Available at: http://ilpubs.stanford.edu:8090/422/.

Papaspiliopoulos, O., Roberts, G. O., and Sköld, M. (2007). A general framework for the parametrization of hierarchical models. *Statistical Science*, 22(1):59–73. Available at: https://arxiv.org/abs/math/0703048.

Russian Federation and United States (2017). Establishment of Two-Way Routes and Precautionary Areas in the Bering Sea and Bering Strait. NCSR Submission NCSR 5/3/7, International Maritime Organization. Submitted 17 November 2017. Available at: https://safety4sea.com/wp-content/uploads/2018/05/IMO-Est

ablishment-of-two-way-routes-and-precautionary-areas-in-the-Bering-Sea-and-Bering-Strait-2017_11.pdf. Accessed: 2026-08-16.

Vehtari, A., Gelman, A., Simpson, D., Carpenter, B., and Bürkner, P.-C. (2021). Rank-normalization, folding, and localization: An improved $\widehat{R}$ for assessing convergence of mcmc. *Bayesian Analysis*, 16(2):667–718.

World Bank Data Lab (2025). Sea routes. Jupyter notebook. *Red Sea Monitoring*. Available at: https://datapartnership.org/red-sea-monitoring/notebooks/routes/2-prepare-sea-routes.html. Updated: 2025-09-30. Accessed: 2026-08-16.

Zhou, C., Ciais, P., Mittakola, R. T., et al. (2026). Global marine LNG terminals, tankers & trade: A high-resolution AIS-based dataset of LNG trade (2020–2024). *Scientific Data*, 13:1153.