

# Basic Calculus for Greeks

November 4, 2022

# Outline

- 1 Definition and Notations
- 2 Derivatives of common functions
- 3 Some Basic Rules
- 4 Taylor Approximation

# Definition and Notations

- The first-order derivative can be defined by

$$f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$$

- ▶ Second-order derivative  $f''(x)$  is the derivative of  $f'(x)$
- The partial derivative of  $f(x_1, x_2)$  w.r.t  $x_1$  can be defined by

$$f_1(x_1, x_2) = \lim_{h \rightarrow 0} \frac{f(x_1+h, x_2) - f(x_1, x_2)}{h}$$

- ▶ Second-order derivatives:  $f_{11}(x_1, x_2)$ ,  $f_{22}(x_1, x_2)$  and  $f_{12}(x_1, x_2) = f_{21}(x_1, x_2)$
- ▶ E.g.  
 $f_{12}(x_1, x_2)$  can be obtained by taking partial derivative of  $f_1(x_1, x_2)$  w.r.t  $x_2$

# Derivatives of common functions

- Power function:  $(x^a)' = ax^{a-1}$ 
  - ▶ E.g.  $(x^5)' = 5x^4$ ;  $(x)' = 1$
- Exponential function:  $(e^{cx})' = ce^{cx}$ 
  - ▶ E.g.  $(e^{2x})' = 2e^{2x}$
- Logarithm function:  $(\ln x)' = \frac{1}{x}$
- The derivative of CDF is its PDF

$$F(x) = \int_{-\infty}^x f(t)dt \quad \text{then} \quad F'(x) = f(x)$$

- ▶ E.g. Standard Normal:  $\Phi'(x) = \phi(x)$

# Some Basic Rules

- Linearity: if  $h(x) = c_1 f(x) + c_2 g(x)$ , then  $\frac{dh^n(x)}{dx^n} = c_1 \frac{df^n(x)}{dx^n} + c_2 \frac{dg^n(x)}{dx^n}$  for any order of derivatives  $n$  if exists
  - ▶ E.g.  $W(S) = \sum_i n_i C_i(S) + \sum_j n_j P_j(S)$   
then  $W'(S) = \sum_i n_i C'_i(S) + \sum_j n_j P'_j(S)$
- Multiplication Rule: if  $h(x) = f(x)g(x)$ , then  $h'(x) = f'(x)g(x) + f(x)g'(x)$ 
  - ▶ E.g.  $h(x) = x^3 \Phi(x)$ , then  $h'(x) = 3x^2 \Phi(x) + x^3 \phi(x)$
- Quotient Rule: if  $h(x) = \frac{f(x)}{g(x)}$ , then  $h'(x) = \frac{f'(x)g(x) - f(x)g'(x)}{g(x)^2}$ 
  - ▶ E.g.  $h(x) = \frac{1}{x}$ , then  $h'(x) = -\frac{1}{x^2}$
- Chain Rule: if  $h(x) = f(g(x))$ , then  $h'(x) = f'(g(x))g'(x)$ 
  - ▶ E.g.  $h(x) = \Phi(\ln(x))$ , then  $h'(x) = \phi(\ln(x))\frac{1}{x}$

# Taylor Approximation

Second-order Taylor expansion for univariate function:

$$f(x) \approx f(x^*) + \frac{df(x^*)}{dx} (x - x^*) + \frac{1}{2} \frac{d^2f(x^*)}{dx^2} (x - x^*)^2$$

$$\Delta f(x^*) \approx \frac{df(x^*)}{dx} \Delta x^* + \frac{1}{2} \frac{d^2f(x^*)}{dx^2} \Delta x^{*2}$$

Second-order Taylor expansion for multivariate function:

$$\begin{aligned} \Delta f(x_1^*, x_2^*) &\approx f_1(x_1^*, x_2^*) \Delta x_1^* + f_2(x_1^*, x_2^*) \Delta x_2^* \\ &+ \frac{1}{2} [f_{11}(x_1^*, x_2^*) \Delta x_1^{*2} + 2f_{12}(x_1^*, x_2^*) \Delta x_1^* \Delta x_2^* + f_{22}(x_1^*, x_2^*) \Delta x_2^{*2}] \end{aligned}$$

For the first-order approximation, drop all second-order terms