

High-spin states of the $N = 82$ isotones ^{136}Xe , ^{137}Cs , and ^{138}Ba : Monopole-driven competition of neutron core-excitations with two-proton excitations to the $h_{11/2}$ high- j orbit

Hua Jin^{1, 2} Shigeru Tazaki³ Kazunari Kaneko⁴
Han-Kui Wang^{2, 5} Yang Sun²

¹Shanghai Dianji University, People's Republic of China

²Shanghai Jiao Tong University, People's Republic of China

³Fukuoka University, Japan


⁴Kyushu Sangyo University, Japan

⁵Zhoukou Normal University, People's Republic of China

Workshop on nuclear structural theory (Mianyang 2020)


- 1 Motivation
- 2 The Model
- 3 Calculated Results
- 4 Conclusion and Discussion

^{132}Sn mass region



taken from website


^{132}Sn mass region




taken from website

- Magic property of ^{132}Sn ^[1, 2]
- Nature 465, 454 (2010), Phys. Rev. Lett. 112, 172701 (2014)
- Nucleon-nucleon interactions^[3-5]
- Phys. Rev. Lett. 110, 192501 (2013); 113, 132502 (2014); 118, 092503 (2017)
- Shell evolutions^[6-9] when far away from the stability valley
- Phys. Rev. Lett. 109, 032501 (2012); 111, 152501 (2013); 113, 042502 (2014); 112, 132501 (2014)
- Nucleosynthesis^[10-14] relating to the astrophysical r process
- Phys. Rev. Lett. 109, 172501 (2012); 111, 061102 (2013); 114, 192501 (2015); 115, 232501 (2015); 118, 202502 (2017)

Neutron core excitations across the $N = 82$ closed shell




taken from website




Phys. Rev. C 65, 017302 (2001)

Neutron core excitations across the $N = 82$ closed shell




taken from website

- The medium- or high-spin states with high-excitation energies in some few-valence-nucleon $N = 82$ and $N = 83$ isotones beyond ^{132}Sn have been detected by analyzing different reaction product gamma ray using large gamma-ray detector arrays ^[15–22].
- Most of them are derived from neutron core excitations across the $N = 82$ closed shell coupling to the valence nucleons. Such core-excited states carry certain structure information.




Phys. Rev. C 65, 017302 (2001)

Shell model description



Shell model description



- 1 Motivation
- 2 The Model**
- 3 Calculated Results
- 4 Conclusion and Discussion

proton-neutron (pn) representation:


$$\begin{aligned}
 H &= H_{\text{sp}} + H_{P_0} + H_{P_2} + H_{QQ} + H_{OO} + H_{HH} + H_{\text{mc}} \\
 &= \sum_{\alpha,i} \varepsilon_a^i c_{\alpha,i}^\dagger c_{\alpha,i} - \frac{1}{2} \sum_{J=0,2} \sum_{ii'} g_{J,ii'} \sum_M P_{JM,ii'}^\dagger P_{JM,ii'} \\
 &\quad - \frac{1}{2} \sum_{\lambda=2,3,4} \sum_{ii'} \frac{\chi_{\lambda,ii'}}{b_0^{2\lambda}} \sum_M : Q_{\lambda M,ii'}^\dagger Q_{\lambda M,ii'} : \\
 &\quad + \sum_{j_a \leq j_b, ii'} k_{mc}(ia, i'b) \sum_{JM} A_{JM}^\dagger(ij_a, i'j_b) A_{JM}(ij_a, i'j_b), \tag{1}
 \end{aligned}$$


proton-neutron (pn) representation:

$$\begin{aligned}
 H &= H_{\text{sp}} + H_{P_0} + H_{P_2} + H_{QQ} + H_{OO} + H_{HH} + H_{\text{mc}} \\
 &= \sum_{\alpha,i} \varepsilon_a^i c_{\alpha,i}^\dagger c_{\alpha,i} - \frac{1}{2} \sum_{J=0,2} \sum_{ii'} g_{J,ii'} \sum_M P_{JM,ii'}^\dagger P_{JM,ii'} \\
 &\quad - \frac{1}{2} \sum_{\lambda=2,3,4} \sum_{ii'} \frac{\chi_{\lambda,ii'}}{b_0^{2\lambda}} \sum_M : Q_{\lambda M,ii'}^\dagger Q_{\lambda M,ii'} : \\
 &\quad + \sum_{j_a \leq j_b, ii'} k_{\text{mc}}(ia, i'b) \sum_{JM} A_{JM}^\dagger(ij_a, i'j_b) A_{JM}(ij_a, i'j_b), \quad (1)
 \end{aligned}$$

Here, the indices i and i' stand for proton(π) or neutron(ν), and b_0 is the harmonic-oscillator range parameter. The separable forces which include the $J = 0$ and $J = 2$ pairing (P_0 and P_2) terms, the multipole-multipole terms($\lambda = 2, 3, 4$), and **the monopole corrections (H_{mc})** are considered.

Model sapce





$$\varepsilon_a^i = \varepsilon_a'^i - \frac{1}{2j_a + 1} \sum_J \sum_{i', b=\text{hole}} (2J + 1) \langle i' j_b, i j_a | V | i' j_b, i j_a \rangle_J \quad (2)$$

- $\varepsilon_a'^i$: single particle or single hole energy from experiments
- ε_a^i : single particle energy for the present model space

Monopole Hamiltonian

$$H_m = \sum_{a,i} \varepsilon_a^i \hat{n}_{ai} + \sum_{ab,ii'} V_{ab}^{ii'} \frac{\hat{n}_{ai}(\hat{n}_{bi'} - \delta_{ab}\delta_{ii'})}{1 + \delta_{ab}\delta_{ii'}}. \quad (3)$$

$V_{ab}^{ii'}$ is the monopole component of the two-body interaction

$$V_{ab}^{ii'} = \frac{\sum_J \langle i j_a, i' j_b | V | i j_a, i' j_b \rangle_J (2J+1)(1+(-1)^J \delta_{ii'} \delta_{ab})}{(2j_a+1)(2j_b+1 - \delta_{ii'} \delta_{ab})}, \quad (4)$$

Effective single-particle energies (ESPEs)

Monopole Hamiltonian

$$H_m = \sum_{a,i} \varepsilon_a^i \hat{n}_{ai} + \sum_{ab,ii'} V_{ab}^{ii'} \frac{\hat{n}_{ai}(\hat{n}_{bi'} - \delta_{ab}\delta_{ii'})}{1 + \delta_{ab}\delta_{ii'}}. \quad (3)$$

$V_{ab}^{ii'}$ is the monopole component of the two-body interaction

$$V_{ab}^{ii'} = \frac{\sum_J \langle i j_a, i' j_b | V | i j_a, i' j_b \rangle_J (2J+1)(1+(-1)^J \delta_{ii'} \delta_{ab})}{(2j_a+1)(2j_b+1 - \delta_{ii'} \delta_{ab})}, \quad (4)$$

ESPE

The ESPE is defined as one-nucleon separation energy for an occupied orbital (or extra binding gained by the addition of a nucleon to an unoccupied orbital) evaluated from the monopole Hamiltonian^[24].

Effective single-particle energies (ESPEs)

Monopole Hamiltonian

$$H_m = \sum_{a,i} \varepsilon_a^i \hat{n}_{ai} + \sum_{ab,ii'} V_{ab}^{ii'} \frac{\hat{n}_{ai}(\hat{n}_{bi'} - \delta_{ab}\delta_{ii'})}{1 + \delta_{ab}\delta_{ii'}}. \quad (3)$$

$V_{ab}^{ii'}$ is the monopole component of the two-body interaction

$$V_{ab}^{ii'} = \frac{\sum_J \langle i j_a, i' j_b | V | i j_a, i' j_b \rangle_J (2J+1)(1+(-1)^J \delta_{ii'} \delta_{ab})}{(2j_a+1)(2j_b+1 - \delta_{ii'} \delta_{ab})}, \quad (4)$$

ESPE

The ESPE is defined as one-nucleon separation energy for an occupied orbital (or extra binding gained by the addition of a nucleon to an unoccupied orbital) evaluated from the monopole Hamiltonian^[24].



Monopole corrections :

$$\begin{aligned} k_{mc}(\nu h_{11/2}, \nu f_{7/2}) &= 0.04, \\ k_{mc}(\nu d_{3/2}, \nu f_{7/2}) &= 0.15, \\ k_{mc}(\pi g_{7/2}, \pi h_{11/2}) &= -0.15, \\ k_{mc}(\pi g_{7/2}, \nu h_{9/2}) &= -0.6, \\ k_{mc}(\pi h_{11/2}, \nu f_{7/2}) &= -1.0, \\ k_{mc}(\pi h_{11/2}, \nu h_{9/2}) &= -0.8. \quad (\text{in MeV}) \end{aligned}$$

- 1 Motivation
- 2 The Model
- 3 Calculated Results**
- 4 Conclusion and Discussion



$N = 82$ isotones ^{134}Te , ^{135}I

Jin, Hasegawa, Tazaki, Kaneko, and Sun, Phys. Rev. C 84, 044324 (2011)




$$k_{mc}(\nu h_{11/2}, \nu f_{7/2}) = 0.04,$$

$$k_{mc}(\nu d_{3/2}, \nu f_{7/2}) = 0.15. \quad (\text{in MeV})$$




Magnetic rotation band in ^{135}Te



Calculated $B(E2)$ and $B(M1)$ values for the core-excitation states in ^{135}Te


Heavier $N = 82$ isotones ^{136}Xe , ^{137}Cs , ^{138}Ba

Jin, Tazaki, Kaneko, Wang and Sun, Phys. Rev. C 100, 064316 (2019)




Most of the high-spin states with **negative-parity** have the predominant configuration of $(\pi g_{7/2} \pi d_{5/2})^{n-1} (\pi h_{11/2})^1$ except the highest 16_1^- and 17_1^- states in ^{138}Ba .

Occupation numbers




Configurations



Competitions between the configurations of **two-proton excitation to the $\pi h_{11/2}$ orbit** and those of **neutron core excitation** across the $N = 82$ closed shell.


Analysis of ESPEs




$$k_{mc}(\pi g_{7/2}, \pi h_{11/2}) = -0.15,$$

$$k_{mc}(\pi g_{7/2}, \nu h_{9/2}) = -0.6, \quad (\text{in MeV})$$

Quasiparticles' excitation based on the BCS theory




Quasiparticles' excitation based on the BCS theory



$$\begin{aligned}
 (\pi g_{7/2})^{n-2}(\pi h_{11/2})^2 &\rightarrow E_{2p} (2p2h) \\
 (\pi g_{7/2})^n(\nu h_{11/2})^{-1}(\nu f_{7/2})^1 &\rightarrow E_{1n} (1p1h) \\
 (\pi g_{7/2})^n(\nu h_{11/2})^{-1}(\nu h_{9/2})^1 &\rightarrow E'_{1n} (1p1h)
 \end{aligned}$$

Quasiparticles' excitation based on the BCS theory



$$\begin{aligned}
 (\pi g_{7/2})^{n-2}(\pi h_{11/2})^2 &\rightarrow E_{2p} (2p2h) \\
 (\pi g_{7/2})^n(\nu h_{11/2})^{-1}(\nu f_{7/2})^1 &\rightarrow E_{1n} (1p1h) \\
 (\pi g_{7/2})^n(\nu h_{11/2})^{-1}(\nu h_{9/2})^1 &\rightarrow E'_{1n} (1p1h)
 \end{aligned}$$

- Quasiparticle energy

$$E_{ik} = \sqrt{(\varepsilon_{ik} - \lambda_i)^2 + \Delta_i^2}, \quad i = \pi \text{ or } \nu$$

- Valence-particle number

$$N_i = \sum_{k>0} \{1 - (\varepsilon_{ik} - \lambda_i) / [(\varepsilon_{ik} - \lambda_i)^2 + \Delta_i^2]^{1/2}\}$$

- 1 Motivation
- 2 The Model
- 3 Calculated Results
- 4 Conclusion and Discussion**

Conclusion

- 1 The effective interaction derived from the EPQQM model has been used to carry out the shell-model calculation to describe the high-spin states in the heavier $N = 82$ isotones beyond ^{132}Sn .
- 2 The competition between neutron core excitation and two-proton excitation to the $\pi h_{11/2}$ orbit has been found in the high-spin states for the heavier $N = 82$ isotones.
- 3 Such competition has been analyzed by the theoretical ESPEs. The adopted monopole corrections contribute to the variation of ESPEs and lead to the structural evolution.

Conclusion

- 1 The effective interaction derived from the EPQQM model has been used to carry out the shell-model calculation to describe the high-spin states in the heavier $N = 82$ isotones beyond ^{132}Sn .
- 2 The competition between neutron core excitation and two-proton excitation to the $\pi h_{11/2}$ orbit has been found in the high-spin states for the heavier $N = 82$ isotones.
- 3 Such competition has been analyzed by the theoretical ESPEs. The adopted monopole corrections contribute to the variation of ESPEs and lead to the structural evolution.

Discussion

- 1 Spurious center-of-mass motion
- 2 To construct the monopole terms starting from the monopole-based universal force $V_{\text{MU}}^{[25, 26]}$
- 3 To extend the present effective interaction to the application in the heavier $N = 83$ isotones

Spurious center-of-mass motion

$$H' = H_{\text{SM}} + H_{\beta} = H_{\text{SM}} + \beta \left\{ \frac{\left(\sum_{i=1}^A \vec{p}_i \right)^2}{2Am} + \frac{1}{2} \frac{m\omega^2}{A} \left(\sum_{i=1}^A \vec{r}_i \right)^2 - \frac{3}{2} \hbar\omega \right\}$$

Spurious center-of-mass motion

$$H' = H_{\text{SM}} + H_{\beta} = H_{\text{SM}} + \beta \left\{ \frac{\left(\sum_{i=1}^A \vec{p}_i \right)^2}{2Am} + \frac{1}{2} \frac{m\omega^2}{A} \left(\sum_{i=1}^A \vec{r}_i \right)^2 - \frac{3}{2} \hbar\omega \right\}$$

Two-body potential^[27] $V_{\beta} = \frac{\beta}{A} \left\{ \frac{\vec{p}_1 \cdot \vec{p}_2}{m} + \frac{1}{2} m\omega^2 \vec{r}_1 \cdot \vec{r}_2 \right\}$

$$\begin{aligned} E_{JT}(j_1 j_2; j_3 j_4) &= \langle n_1 l_1 j_1, n_2 l_2 j_2 | V_{\beta} | n_3 l_3 j_3, n_4 l_4 j_4 \rangle \\ &= \frac{\beta \hbar \omega}{A} \sum_{\lambda S n l N L} \frac{\{1 - (-1)^{S+T+I}\}}{\sqrt{(1 + \delta_{n_1 n_2} \delta_{l_1 l_2} \delta_{j_1 j_2})(1 + \delta_{n_3 n_4} \delta_{l_3 l_4} \delta_{j_3 j_4})}} \\ &\quad \times (2\lambda + 1)(2S + 1) \sqrt{(2j_1 + 1)(2j_2 + 1)(2j_3 + 1)(2j_4 + 1)} \\ &\quad \times \left\{ \begin{matrix} l_1 & l_2 & \lambda \\ \frac{1}{2} & \frac{1}{2} & S \\ j_1 & j_2 & J \end{matrix} \right\} \left\{ \begin{matrix} l_3 & l_4 & \lambda \\ \frac{1}{2} & \frac{1}{2} & S \\ j_3 & j_4 & J \end{matrix} \right\} [(2N + L) - (2n + I)] \\ &\quad \times M_{\lambda}(n l N L : n_1 l_1 n_2 l_2) M_{\lambda}(n l N L : n_3 l_3 n_4 l_4) \end{aligned}$$

Spurious center-of-mass motion

$$H' = H_{\text{SM}} + H_{\beta} = H_{\text{SM}} + \beta \left\{ \frac{\left(\sum_{i=1}^A \vec{p}_i \right)^2}{2Am} + \frac{1}{2} \frac{m\omega^2}{A} \left(\sum_{i=1}^A \vec{r}_i \right)^2 - \frac{3}{2} \hbar\omega \right\}$$

Two-body potential^[27] $V_{\beta} = \frac{\beta}{A} \left\{ \frac{\vec{p}_1 \cdot \vec{p}_2}{m} + \frac{1}{2} m\omega^2 \vec{r}_1 \cdot \vec{r}_2 \right\}$

$$\begin{aligned} E_{JT}(j_1 j_2; j_3 j_4) &= \langle n_1 l_1 j_1, n_2 l_2 j_2 | V_{\beta} | n_3 l_3 j_3, n_4 l_4 j_4 \rangle \\ &= \frac{\beta \hbar \omega}{A} \sum_{\lambda S n l N L} \frac{\{1 - (-1)^{S+T+I}\}}{\sqrt{(1 + \delta_{n_1 n_2} \delta_{l_1 l_2} \delta_{j_1 j_2})(1 + \delta_{n_3 n_4} \delta_{l_3 l_4} \delta_{j_3 j_4})}} \\ &\quad \times (2\lambda + 1)(2S + 1) \sqrt{(2j_1 + 1)(2j_2 + 1)(2j_3 + 1)(2j_4 + 1)} \\ &\quad \times \left\{ \begin{matrix} l_1 & l_2 & \lambda \\ \frac{1}{2} & \frac{1}{2} & S \\ j_1 & j_2 & J \end{matrix} \right\} \left\{ \begin{matrix} l_3 & l_4 & \lambda \\ \frac{1}{2} & \frac{1}{2} & S \\ j_3 & j_4 & J \end{matrix} \right\} [(2N + L) - (2n + I)] \\ &\quad \times M_{\lambda}(n l N L : n_1 l_1 n_2 l_2) M_{\lambda}(n l N L : n_3 l_3 n_4 l_4) \end{aligned}$$

Modification of the single-particle energy^[27]

$$\tilde{\epsilon}_j = \epsilon_j + (2n + I + \frac{3}{2}) \frac{\beta \hbar \omega}{A} + \frac{1}{2(2j + 1)} \sum_{JTj_c} (2J + 1)(2T + 1) E_{JT}(jj_c; jj_c)$$

Spurious center-of-mass motion

$$H' = H_{\text{SM}} + H_{\beta} = H_{\text{SM}} + \beta \left\{ \frac{\left(\sum_{i=1}^A \vec{p}_i \right)^2}{2Am} + \frac{1}{2} \frac{m\omega^2}{A} \left(\sum_{i=1}^A \vec{r}_i \right)^2 - \frac{3}{2} \hbar\omega \right\}$$

Two-body potential^[27] $V_{\beta} = \frac{\beta}{A} \left\{ \frac{\vec{p}_1 \cdot \vec{p}_2}{m} + \frac{1}{2} m\omega^2 \vec{r}_1 \cdot \vec{r}_2 \right\}$


$$\begin{aligned} E_{JT}(j_1 j_2; j_3 j_4) &= \langle n_1 l_1 j_1, n_2 l_2 j_2 | V_{\beta} | n_3 l_3 j_3, n_4 l_4 j_4 \rangle \\ &= \frac{\beta \hbar \omega}{A} \sum_{\lambda} \frac{\{1 - (-1)^{S+T+I}\}}{S n l N L \sqrt{(1 + \delta_{n_1 n_2} \delta_{l_1 l_2} \delta_{j_1 j_2})(1 + \delta_{n_3 n_4} \delta_{l_3 l_4} \delta_{j_3 j_4})}} \\ &\quad \times (2\lambda + 1)(2S + 1) \sqrt{(2j_1 + 1)(2j_2 + 1)(2j_3 + 1)(2j_4 + 1)} \\ &\quad \times \left\{ \begin{matrix} l_1 & l_2 & \lambda \\ \frac{1}{2} & \frac{1}{2} & S \\ j_1 & j_2 & J \end{matrix} \right\} \left\{ \begin{matrix} l_3 & l_4 & \lambda \\ \frac{1}{2} & \frac{1}{2} & S \\ j_3 & j_4 & J \end{matrix} \right\} [(2N + L) - (2n + I)] \\ &\quad \times M_{\lambda}(n l N L : n_1 l_1 n_2 l_2) M_{\lambda}(n l N L : n_3 l_3 n_4 l_4) \end{aligned}$$

Modification of the single-particle energy^[27]


$$\tilde{\epsilon}_j = \epsilon_j + (2n + I + \frac{3}{2}) \frac{\beta \hbar \omega}{A} + \frac{1}{2(2j + 1)} \sum_{JTj_c} (2J + 1)(2T + 1) E_{JT}(jj_c; jj_c)$$

D. H. Gloeckner and R. D. Lawson, Phys. Lett. 53B, 313 (1974)


1⁻ spuriousities




1⁻ spuriousities



1⁻ spuriousities



1⁻ spuriousities



- Spurious center-of-mass motion has an effect of the order $1/A$ ^[28]
- Such spuriousities could be neglected for heavy nuclei

Monopole interaction


Monopole-based universal interaction V_{MU}

Phys. Rev. Lett. 104, 012501 (2010)


$$V_{MU} \rightarrow V_C (\text{Gaussian}) + V_T (\pi + \rho)$$

• Central Gaussian force $V_C = \sum_{S,T} f_{S,T} P_{S,T} \exp \left[-(r/\mu)^2 \right]$

• Tensor force $V_T = (\vec{\tau}_1 \cdot \vec{\tau}_2) \left([\vec{s}_1 \vec{s}_2]^{(2)} \cdot Y^{(2)} \right) f(r)$



Neutron $\nu i_{13/2}$ excitation in $N = 83$ isotones



Phys. Rev. C 63, 024322 (2001)

Thank you for your attention!!



- [1] K. L. Jones, *et al.* Nature **465**, 454 (2010).
- [2] J. M. Allmond, *et al.* Phys. Rev. Lett. **112**, 172701 (2014).
- [3] D. T. Yordanov, *et al.* Phys. Rev. Lett. **110**, 192501 (2013).
- [4] G. S. Simpson, *et al.* Phys. Rev. Lett. **113**, 132502 (2014).
- [5] J. M. Allmond, *et al.* Phys. Rev. Lett. **118**, 092503 (2017).
- [6] J. Hakala, *et al.* Phys. Rev. Lett. **109**, 032501 (2012).
- [7] H. Watanabe, *et al.* Phys. Rev. Lett. **111**, 152501 (2013).
- [8] H. Watanabe, *et al.* Phys. Rev. Lett. **113**, 042502 (2014).
- [9] J. Taprogge, *et al.* Phys. Rev. Lett. **112**, 132501 (2014).
- [10] R. L. Kozub, *et al.* Phys. Rev. Lett. **109**, 172501 (2012).
- [11] J. Van Schelt, *et al.* Phys. Rev. Lett. **111**, 061102 (2013).
- [12] G. Lorusso, *et al.* Phys. Rev. Lett. **114**, 192501 (2015).
- [13] D. Atanasov, *et al.* Phys. Rev. Lett. **115**, 232501 (2015).

- [14] V. Vaquero, *et al.* Phys. Rev. Lett. **118**, 202502 (2017).
- [15] P. J. Daly, *et al.* Phys. Rev. C **59**, 3066 (1999).
- [16] R. Broda, *et al.* Phys. Rev. C **59**, 3071 (1999).
- [17] C. T. Zhang, *et al.* Phys. Rev. Lett. **77**, 3743 (1996).
- [18] Urban, *et al.* Phys. Rev. C **62**, 027301 (2000).
- [19] B. Fornal, *et al.* Phys. Rev. C **63**, 024322 (2001).
- [20] S. K. Saha, *et al.* Phys. Rev. C **65**, 017302 (2001).
- [21] K. Li, *et al.* Phys. Rev. C **75**, 044314 (2007).
- [22] A. Astier, *et al.* Phys. Rev. C **85**, 064316 (2012).
- [23] L. Coraggio, A. Covello, A. Gargano, N. Itaco, and T. T. S. Kuo, Phys. Rev. C **80**, 044320 (2009).
- [24] N. A. Smirnova, B. Ballya, K. Heyde, F. Nowacki, and K. Sieja, Phys. Lett. B **686**, 109 (2010).

- [25] K. Kaneko, T. Mizusaki, Y. Sun, and S. Tazaki, Phys. Rev. C **89**, 011302(R) (2014).
- [26] T. Otsuka, *et al.* Phys. Rev. Lett. **104**, 012501 (2010)
- [27] D. H. Gloeckner and R. D. Lawson, Phys. Lett. **53B**, 313 (1974).
- [28] P. Ring and P. Schuck, *The Nuclear Many-Body Problem* (Spring-Verlag, New York, 1980).