




曲阜师范大学
QUFU NORMAL UNIVERSITY

跃迁求和规则作为近似方法的判据

Yi Lu (路毅), Qufu Normal University,
MianYang 2020.1.13





曲阜师范大学
QUFU NORMAL UNIVERSITY

曲阜师范大学





粒子物理： 邵凤兰（教授）
王瑞芹（副教授）
强子物理： 李刚（教授）
李翠（讲师）
高能天体物理： 张志彬（教授）
仪双喜（副教授）
杨玉鹏（副教授）
原子核物理： 路毅（讲师）
房建军（讲师）



Transition sum rules as a test stone

1. Derivation of transition sum rules
2. Code Implementation
3. Application: test of Brink-Axel hypothesis, calculation of TRK sum rules.
4. (New) Shell model v.s. HF&PHF v.s. NPA: sum rules from ground states.

Collaborators:

Calvin W. Johnson, San Diego State University

Ken Luu, San Diego State University



单体跃迁算符

$$\hat{O}_{KM} = \sum_{ab} F_{ab}[K]^{-1} [c_a^\dagger \otimes \tilde{c}_b]_{KM}$$

$$\begin{aligned} B(J_i \rightarrow J_f) &= \frac{1}{2J_i + 1} \sum_{M_f M_i M} |\langle J_f M_f | O_{KM} | J_i M_i \rangle|^2 \\ &= \frac{1}{2J_i + 1} \sum_{M_f} \sum_{M_i M} (J_i M_i K M | J_f M_f)^2 |\langle J_f || O_K || J_i \rangle|^2 \\ &= \frac{2J_f + 1}{2J_i + 1} |\langle J_f || O_K || J_i \rangle|^2 \end{aligned}$$



$$S(E_i, E_x) \equiv \sum_f \delta(E_x - E_f + E_i) B(F; i \rightarrow f)$$

$$S_k(E_i) = \int (E_x)^k S(E_i, E_x) dE_x.$$

Non-energy weighted sum rule $S_0(E_i) = \int S(E_i, E_x) dE_x$

e.g. **Ikeda sum rule** of *Gamow-Teller transitions*

$$S_0(GT_-) - S_0(GT_+) = 3(N - Z)g_A^2$$

Energy-weighted sum rule $S_1(E_i) = \int E_x S(E_i, E_x) dE_x$

e.g. **Thomas-Reiche-Kuhn** sum rule of *electric dipole transitions*

In atomic physics: $S_1 = \frac{3e^2\hbar^2}{2m_e} N_e$ (for a multi-electron atom)

In nuclear physics: $S_1 \sim \frac{NZe^2\hbar^2}{Am_N}$



0阶求和规则：总跃迁强度

We can define $\mathcal{F}_{ba} \equiv (-1)^{1+j_a+j_b} F_{ab}^*$, and

$$\mathcal{F}_{K-M} \equiv (-1)^M \hat{F}_{KM}^\dagger = [K]^{-1} \sum_{ab} \mathcal{F}_{ba} (\hat{b}^\dagger \otimes \tilde{a})_{K-M}.$$

$$B(F) = \frac{1}{2J_i + 1} \sum_{M_i M M_f} |\langle J_f M_f | \hat{F}_{KM} | J_i M_i \rangle|^2,$$

$$\begin{aligned} S_0(F) &= \frac{1}{2J_i + 1} \sum_{M_i M M_f J_f \alpha} \langle J_i M_i | \hat{F}_{KM}^\dagger | J_f M_f, \alpha \rangle \langle J_f M_f, \alpha | \hat{F}_{KM} | J_i M_i \rangle \\ &= \frac{1}{2J_i + 1} \sum_{M_i} \langle J_i M_i | \sum_M \hat{F}_{KM}^\dagger \hat{F}_{KM} | J_i M_i \rangle \\ &= \langle J_i M_i | (-1)^K [K] (\mathcal{F}_K \otimes \hat{F}_K)_{00} | J_i M_i \rangle \end{aligned}$$



1阶求和规则：能量权重求和规则

$$\begin{aligned}
 S_1 &= \frac{1}{2J_i + 1} \sum_{M_f M_i M} |\langle J_f M_f | O_{KM} | J_i M_i \rangle|^2 (E_f - E_i) \\
 &= \frac{1}{2J_i + 1} \sum_{M_f M_i M} \langle J_i M_i | O_{KM}^\dagger | J_f M_f \rangle \langle J_f M_f | O_{KM} | J_i M_i \rangle (E_f - E_i) \\
 &= \frac{1}{2(2J_i + 1)} \sum_{M_f M_i M} (\langle J_i M_i | O_{KM}^\dagger H | J_f M_f \rangle \langle J_f M_f | O_{KM} | J_i M_i \rangle + \langle J_i M_i | O_{KM}^\dagger | J_f M_f \rangle \langle J_f M_f | H O_{KM} | J_i M_i \rangle \\
 &\quad - \langle J_i M_i | H O_{KM}^\dagger | J_f M_f \rangle \langle J_f M_f | O_{KM} | J_i M_i \rangle - \langle J_i M_i | O_{KM}^\dagger | J_f M_f \rangle \langle J_f M_f | O_{KM} H | J_i M_i \rangle) \\
 &= \frac{1}{2} \langle J_i M_i | \sum_M (2O_{KM}^\dagger H O_{KM} - H O_{KM}^\dagger O_{KM} - O_{KM}^\dagger O_{KM} H) | J_i M_i \rangle. \\
 S_1 &= \frac{1}{2} \langle J_i M_i | \sum_M (-1)^M (2O_{K-M} H O_{KM} - H O_{K-M} O_{KM} - O_{K-M} O_{KM} H) | J_i M_i \rangle \\
 &= \frac{1}{2} \langle J_i M_i | \sum_M (-1)^{1+M} [(H, O)_{KM}, O_{K-M}] | J_i M_i \rangle \\
 &= \frac{1}{2} \langle J_i M_i | \sum_M (-1)^{1+M} [(H, O)_{KM}, O_{K-M}] | J_i M_i \rangle \\
 &= \langle J_i M_i | \frac{(-1)^{1+K}}{2} \sqrt{2K+1} [(H, O), O]_{00} | J_i M_i \rangle
 \end{aligned}$$



sum rule = 求和算符的期望值

$$\begin{aligned}
 S_0(\hat{F}_K, i \rightarrow all) &= \frac{1}{2J_i + 1} \sum_{M_i M M_f J_f} \langle J_i M_i | \hat{F}_{KM}^\dagger | J_f M_f \rangle \langle J_f M_f | \hat{F}_{KM} | J_i M_i \rangle \\
 &= \frac{1}{2J_i + 1} \sum_{M_i} \langle J_i M_i | \sum_M \hat{F}_{KM}^\dagger \hat{F}_{KM} | J_i M_i \rangle \\
 &= \langle J_i M_i | (-1)^K [K] (\hat{F}_K \otimes \hat{F}_K)_{00} | J_i M_i \rangle
 \end{aligned}$$

That is to say, the NEWSR equals the expectation value of an operator

$$\hat{O}_{NEWSR} = (-1)^K [K] (\hat{F}_K \otimes \hat{F}_K)_{00}.$$

$$\hat{O}_{NEWSR} = \sum_{ab} \frac{\sum_c \mathbf{F}_{ca}^* \mathbf{F}_{cb}}{[j_a]} \delta_{j_a j_b} (\hat{\mathbf{a}}^\dagger \otimes \tilde{\mathbf{b}})_{00} + \sum_{abcd} \mathbf{F}_{cb}^* \mathbf{F}_{ad} \left\{ \begin{matrix} j_a & j_d & K \\ j_c & j_b & I \end{matrix} \right\} [\mathbf{I}] [(\hat{\mathbf{a}}^\dagger \otimes \hat{\mathbf{b}}^\dagger)_{\mathbf{I}} \otimes (\tilde{\mathbf{c}} \otimes \tilde{\mathbf{d}})_{\mathbf{I}}]_{00}$$



$$S_1 = \sum_f (E_f - E_i) |\langle f | T | i \rangle|^2 = \langle i | O_{\text{d.c.}} | i \rangle$$

$$\begin{aligned} O_{\text{d.c.}} &= \frac{(-1)^K}{2} \sqrt{2K+1} [T_K, (H, T_K)]_0 \\ &= \sum_{ab} g_{ab} \hat{a} (a^\dagger \times \tilde{b})_0 + \frac{1}{4} \sum_{abcd} \sqrt{(1+\delta_{ab})(1+\delta_{cd})} \\ &\quad \times \sum_I \hat{I} W(abcd; I) \left[A_I^\dagger(ab) \times \tilde{A}_I(cd) \right]_0, \end{aligned}$$



$$g_{ab} = \frac{\delta_{j_a j_b}}{2(2j_a + 1)} \sum_{cd} (-e_{ac} F_{cd} F_{bd}^* + F_{ac} e_{cd} F_{bd}^* + F_{ca}^* e_{cd} F_{db} - F_{ca}^* F_{cd} e_{db}), \quad (23)$$

$$W^1(abcd; J) = -\frac{1}{2}(1 + \mathcal{P}_{cdJ}) \sum_{efJ'} (-1)^{J+J'} (2J' + 1) \pi_{de}^{J'} \zeta_{ef} \zeta_{cd}^{-1} V_J(ab, ef) \\ \times F_{ec} F_{fd} \left\{ \begin{matrix} J & K & J' \\ j_d & j_e & j_f \end{matrix} \right\} \left\{ \begin{matrix} J & K & J' \\ j_e & j_d & j_c \end{matrix} \right\}, \quad (25)$$

$$W^2(abcd; J) = -\frac{1}{2}(1 + \mathcal{P}_{cdJ}) \sum_{efJ'} (2J' + 1) \pi_{cf}^{J'} \zeta_{ce} \zeta_{cd}^{-1} V_J(ab, ce) \\ \times F_{ef} F_{df}^* \left\{ \begin{matrix} J & K & J' \\ j_f & j_c & j_e \end{matrix} \right\} \left\{ \begin{matrix} J & K & J' \\ j_f & j_c & j_d \end{matrix} \right\}, \quad (26)$$

$$W^3(abcd; J) = (1 + \mathcal{P}_{abJ})(1 + \mathcal{P}_{cdJ}) \sum_{efJ'} (2J' + 1) \zeta_{be} \zeta_{df} \zeta_{ab}^{-1} \zeta_{cd}^{-1} V_{J'}(be, df) \\ \times F_{ea}^* F_{fc} \left\{ \begin{matrix} J & K & J' \\ j_e & j_b & j_a \end{matrix} \right\} \left\{ \begin{matrix} J & K & J' \\ j_f & j_d & j_c \end{matrix} \right\}, \quad (27)$$

$$W^4(abcd; J) = P_{ac} P_{bd} W^{1*}(abcd; J), \quad (28)$$

$$W^5(abcd; J) = P_{ac} P_{bd} W^{2*}(abcd; J), \quad (29)$$

Y. Lu, C. W. Johnson, PRC 97, 034320(2018),
“Transition sum rules in the Shell Model”.



luyi07/PandasCommute: This is a small code to be used in nuclear structure calculations. It helps evaluate energy-weighted-sum rules of one-body transitions in the shell model.

33 commits 3 branches 0 releases 1 contributor

Branch: master New pull request Create new file Upload files Find file Clone or download

luyi07	deleted big file warehouse/9hbaromega.int	Latest commit c9b1e50 on 23 May
README	fixed the bug in omp parallelization	4 months ago
code	fixed the bug in read_F_new() in code/input.cpp	2 months ago
examples	fixed the bug in omp parallelization	4 months ago
input	added new function to generate random interactions	2 months ago
output	9 hbar-omega input	2 months ago
warehouse	deleted big file warehouse/9hbaromega.int	2 months ago
README.md	added acknowledgement to README	4 months ago
compile.sh	Upgraded the algorithm to be applicable to 2-body core-shell models	5 months ago


Capable to do 15 $\hbar\omega$ shells on a small workstation

<https://github.com/luyi07/PandasCommute>



Test the Brink-Axel hypothesis

Energy-weighted sum rules (EWSR) and transition strength function centroids as a function of initial energy E_i . Results are put into 2-MeV bins with the average and root-mean-square fluctuation shown; the fluctuations are not sensitive to the size of the bins. (a) EWSRs for isoscalar $E2$ for ^{34}Cl in the sd shell. The (red) solid line is the secular behavior predicted by spectral distribution theory, as described in Ref. [7]. (b) Centroids for $M1$ transitions in ^{21}Ne in the sd shell. (c) EWSR for $E1$ transitions in ^{10}B in $0p-1s-0d_{5/2}$ space. (d) Centroids for GT transitions, sum of β_{\pm} , for ^{27}Ne in the sd shell.




Brink-Axel hypothesis: $S(E_i, E_x) \equiv \sum_f \delta(E_x - E_f + E_i) B(F; i \rightarrow f)$ is independent on E_i . If that's true, **EWSR should be independent on E_i** .

Y. Lu, C. W. Johnson, PRC 97, 034320(2018),
"Transition sum rules in the Shell Model".



Calculate the TRK sum rule

PandasCommute
+
BigStick



The **Thomas-Reiche-Kuhn** sum rule: $S_1 \sim \frac{NZe^2\hbar^2}{Am_N}$

Our calculation: For $A=8-18$ nuclei, $S_1/\frac{NZe^2\hbar^2}{Am_N} \in (1.4, 2.3)$, which is in consistence with expr. Data, although we use a modest valence space: spd5.



Validity of Nucleon pair approximation

$$\hat{A}_r^\dagger = \sum_{ab} y(abr)(\hat{a}^\dagger \otimes \hat{b}^\dagger)_r$$

$$\hat{A}_{J_N M_N}^\dagger(\alpha_N) = |r_1 r_2 \cdots r_N; J_1 J_2 \cdots J_N; M_N\rangle = ((\hat{A}_{r_1}^\dagger \otimes \hat{A}_{r_2}^\dagger)_{J_2} \cdots)_{J_N M_N}$$

Intrinsic structure of pairs are frozen ($y(abr)$ are fixed)

	⁴⁵ Ca			
7/2 ₁ ⁻	0.990	0.997	0.991	0.999
5/2 ₁ ⁻	0.975	0.986	0.990	0.999
3/2 ₁ ⁻	0.984	0.985	0.985	0.997
11/2 ₁ ⁻	0.977	0.987	0.987	1.000
9/2 ₁ ⁻	0.983	0.986	0.992	0.999
15/2 ₁ ⁻	0.979	0.996	0.995	1.000
1/2 ₁ ⁻	0.714	0.935	0.814	0.975
13/2 ₁ ⁻	0.058	0.937	0.711	0.977
17/2 ₁ ⁻		0.982	0.898	0.995
19/2 ₁ ⁻		0.931	0.528	0.998
21/2 ₁ ⁻		0.955	0.996	1.000
3/2 ₂ ⁻	0.566	0.847	0.670	0.925
5/2 ₂ ⁻	0.743	0.846	0.890	0.970
7/2 ₂ ⁻	0.434	0.842	0.697	0.884
9/2 ₂ ⁻	0.896	0.966	0.976	0.988
11/2 ₂ ⁻	0.643	0.957	0.873	0.981
13/2 ₂ ⁻		0.909	0.668	0.972
15/2 ₂ ⁻		0.570	0.098	0.987
17/2 ₂ ⁻		0.891	0.337	0.985
19/2 ₂ ⁻		0.870	0.540	0.970

spin ^{parity}	SM	SD	FP
0 ⁺	8 316	4	64
2 ⁺	37 219	7	213
4 ⁺	54 190	4	307
6 ⁺	57 309		336
8 ⁺	50 319		313
10 ⁺	38 242		257

Truncations prove to be effective at low-lying states



$$\hat{A}_r^\dagger = \sum_{ab} y(abr)(\hat{a}^\dagger \otimes \hat{b}^\dagger)_r$$

$$\hat{A}_{J_N M_N}^\dagger(\alpha_N) = |r_1 r_2 \cdots r_N; J_1 J_2 \cdots J_N; M_N\rangle = ((\hat{A}_{r_1}^\dagger \otimes \hat{A}_{r_2}^\dagger)_{J_2} \cdots)_{J_N} M_N$$

变分法：调节集体对结构系数，使得哈密顿量期望值取极小值

$$\epsilon_{NPA} = \langle NPA | \hat{H} | NPA \rangle.$$

共轭梯度法是一种梯度下降算法，对结构系数一般有几十个自由参数，所以需要几十次一维优化，每次一维优化需要几十次迭代，所以一个核需要计算几百上千次，才能得到“最优参数”。适用于价核子数较少的核。



Comparison

In the same valence space, use the same effective interactions:

Shell Model

arXiv:1801.08432 [pdf, ps, other] [physics.comp-ph](#) [nucl-th](#)

BIGSTICK: A flexible configuration-interaction shell-model code

Authors: Calvin W. Johnson, W. Erich Ormand, Kenneth S. McElvain, Hongzhang Shan

NPA: PandaWarrior



Physics Reports

Volume 545, Issue 1, 1 December 2014, Pages 1-45



Nucleon-pair approximation to the nuclear shell model



Y.M. Zhao ^a ✉, A. Arima ^{a, b}

Hartree-Fock、Projected Hartree-Fock in the occupation space

- “The random phase approximation vs. exact shell-model correlation energies,” I. Stetcu and **C. W. Johnson**, Phys. Rev. C **66** 034301 (2002).
- “SU(3) vs. projected Hartree-Fock state,” **C. W. Johnson**, I. Stetcu, and J. P. Draayer, Phys. Rev. C **66**, 034312 (2002).
- “Scalar ground-state observables in the random phase approximations,” **C. W. Johnson** and I. Stetcu, Phys. Rev. C **66**, 064304 (2002).
- “Tests of the random phase approximation for transitions strengths,” I. Stetcu and **C. W. Johnson**, Phys. Rev. C. **67**, 043315 (2003).
- “Gamow-Teller transitions and deformation in the proton-neutron random phase approximation,” I. Stetcu and **C. W. Johnson**, Phys. Rev. C. **69**, 024311 (2004).
- “Collapse of the random phase approximation: examples and counter-examples from the shell model,” **C. W. Johnson** and I. Stetcu, Phys. Rev C **80**, 024320 (2009).
- “Projection of angular momentum by linear algebra,” **C. W. Johnson** and K. D. O'Mara, Phys. Rev C **96**, 064304 (2017)
- “Convergence and efficiency of angular momentum projection for many-body systems,” **C.W. Johnson** and C.-F. Jiao, J. Phys. G **46**, 015101-015115 (2019).



Preliminary results





sd壳所有 $Z \leq N$ 的原子核基态
usdb, HF v.s. PHF v.s. NPA(SDG)



M1: non-energy-weighted sum rules & energy-weighted sum rules

Preliminary results




sd壳所有 $Z \leq N$ 的原子核基态
usdb, HF v.s. PHF v.s. NPA(SDG)




GT: non-energy-weighted sum rules & energy-weighted sum rules


GTminusnewsr



GTplusnewsr




GTewsr






Preliminary results

pf shell NPA v.s. SM: g.s. E2 NEWSR



pf shell NPA v.s. SM: g.s. E2 EWSR




pf壳部分原子核基态
gx1a, HF v.s. PHF v.s. NPA(SDGI)



Preliminary results

pf shell NPA v.s. SM: g.s. M1 NEWSR



pf壳部分原子核基态
gx1a, HF v.s. PHF v.s. NPA(SDGI)



Summary

1. 对于sd壳、pf壳一些开壳核，SDG/SDGI可以描述基态。
2. Transition sum rules 是Model-independent的判据，可以用来测试近似方法，配对截断与严格对角化结果非常接近。
3. 原则上，Configuration-Interaction 的方法都可以通过sum rule 检验波函数是否互相一致！



曲阜师范大学
QUFU NORMAL UNIVERSITY

谢谢!

