Module 2.2 | 15 candidate ASVs at actual G4 sampling nodes

C01 3cae56af0b082bba9c858d1844913fc3
Candidatus_Competibacter

C02 d1725130559823512a63bbc2c1bab523
Incertae_Sedis

C03 78c2842bcb5d089b057f231ddd53b8e1
Defluviicoccus

C04 be9af24f8e82484720783bcb4df4f6de
Chryseotalea

C05 54089b19ce1b3690c1e03d4d34a0162c
Aridibacter

C06 3b91d8fd7dc57a941d49bed1b084064d
Acinetobacter

C07 9d94363a8ab81a79ecf766644aa45b23
Acinetobacter

C08 82c70888e3f2436874633b4285a188e9
Defluviicoccus

C09 a578dfb348f923c17e372f94df5716ef
Unclassified genus

C10 c32a06324874a524e422813afbe86397
Chryseosolibacter

C11 f0e953dd2a7892838f44fea2495a408f
Candidatus_Competibacter

C12 b815668a1e362dc6a897d2895c351449
Acinetobacter

C13 f23e2ebe64cf70326440deb743f4bf85
Candidatus_Competibacter

C14 3e1088366bf860effa27b1322906df56
Candidatus_Competibacter

C15 f23982138e4d5ed3fcb068cbac95af41
Acinetobacter

Relative abundance (% of all ASV reads; independent panel scales)

Recorded G4 day (day 1 = recovered pre−switch baseline)

Sampling position ○ Pre−switch baseline ● Post−adaptation

Reactor R1 R2 R3 R4

R1: 1,76,145; R2/R3/R4: 1,30,76,145. Lines connect observations only; no imputed nodes.
G4 operating labels: R1/R3 = 30 C; R2/R4 = ambient, unmeasured. Different initial sludge states.
Closed compositional data; persistence is detection at sampled nodes, not viability, function or continuous residence.