
G5 | R1 size fractions | Phylum
One mother sample; 145 + 18 d (derived); DO stage setting 0.5+/-0.2 mg/L; no biological replication
Relative abundance
100%
75%
50%
25%
0%
S1
<100 um
S2
100-200 um
S3
200-500 um
S4
>500 um
Recorded sampling order within each run (equal bar spacing)
Phylum
Pseudomonadota
Bacteroidota
Chloroflexota--other
Chlorobiota
Acidobacteriota
Nitrospirota
Myxococcota
Thermomicrobiota
Planctomycetota
Minisyncoccota
Others
Unclassified
Original counts; no rarefaction/filtering. Stage DO settings are not measured DO. G4 baseline setting is uncertain.


G5 | R1 size fractions | Class
One mother sample; 145 + 18 d (derived); DO stage setting 0.5+/-0.2 mg/L; no biological replication
Relative abundance
100%
75%
50%
25%
0%
S1
<100 um
S2
100-200 um
S3
200-500 um
S4
>500 um
Recorded sampling order within each run (equal bar spacing)
Class
Gammaproteobacteria
Alphaproteobacteria
Anaerolineae
Cytophagia
Saprospiria
Blastocatellia
Ignavibacteria--other
Sphingobacteriia
Chitinophagia
Candidatus_Kapaibacteriia
Others
Unclassified
Original counts; no rarefaction/filtering. Stage DO settings are not measured DO. G4 baseline setting is uncertain.

G5 | R1 size fractions | Genus

One mother sample; 145 + 18 d (derived); DO stage setting 0.5+/-0.2 mg/L; no biological replication



Original counts; no rarefaction/filtering. Stage DO settings are not measured DO. G4 baseline setting is uncertain.