


Original counts; no rarefaction/filtering. Stage DO settings are not measured DO. G4 baseline setting is uncertain.




Original counts; no rarefaction/filtering. Stage DO settings are not measured DO. G4 baseline setting is uncertain.


Batch_01 main composition | Genus
29 samples (G1-G4); run-specific day axes; empty positions are unsampled observations
1 Startup
2 After mixing / splitting
3 After storage / recovery
R1
R2
R3
R4
Relative abundance
100%
75%
50%
25%
0%
A0 G1 | d1
A1 G1 | d~120
B0 G1 | d1
B1 G1 | d~120
C0 G1 | d1
C1 G1 | d~120
D0 G1 | d1
D1 G1 | d~120
A2 G3 | d56
B2 G2 | d21
B3 G3 | d56
C2 G3 | d56
D2 G2 | d21
D3 G3 | d56
E1 G4 | baseline DO setting not set: 0.5+/-0.2
E2 G4 | d76 DO set: 0.5+/-0.2
E3 G4 | d145 DO set: 0.5+/-0.2
F1 G4 | baseline DO setting not set: 0.5+/-0.2
F2 G4 | d30 DO set: 0.5+/-0.2
F3 G4 | d76 DO set: 0.5+/-0.2
F4 G4 | d145 DO set: 0.5+/-0.2
G1 G4 | baseline DO setting not set: 0.5+/-0.2
G2 G4 | d30 DO set: 0.5+/-0.2
G3 G4 | d76 DO set: 0.5+/-0.2
G4 G4 | d145 DO set: 0.5+/-0.2
H1 G4 | baseline DO setting not set: 0.5+/-0.2
H2 G4 | d30 DO set: 0.5+/-0.2
H3 G4 | d76 DO set: 0.5+/-0.2
H4 G4 | d145 DO set: 0.5+/-0.2
Recorded sampling order within each run (equal bar spacing)
Acinetobacter
Candidatus_Tepidaquicella
eudomonadati--other;p__Bacteroidota;c__Sphingobacteriia;o__Sphingobacteriales;f__Sphingobacteriaceae;g__Incertae_Sedis]
Incertae_Sedis [d__Bacteria;k__Pseudomonadati--other;p__Bacteroidota;c__Saprospiria;o__Saprospirales;f__Lewinellaceae;
Hassallia
Chryseosolibacter
Unclassified
Original counts; no rarefaction/filtering. Stage DO settings are not measured DO. G4 baseline setting is uncertain.