

**Akash Institute of Medical Sciences &
Research Centre (AIMSRC)**

Patient Name:	HANS RAJ	Age/Gender:	33Y/M	Patient ID:	3049906
Study Date:	20250918	Ref. Doctor:	john	Accession No:	53344
Reported Date:	19/2/2026, 11:01:22 am	Modality:	CR	Body Part:	PELVIS

CR PELVIS REPORT

History:

Patient presents with pelvic pain / history of trauma / difficulty in walking. (Modify as needed)The bony pelvis shows normal alignment. The iliac bones, sacrum, pubic rami, and acetabula appear intact with no evidence of fracture or lytic or sclerotic lesions. Both hip joints are well aligned. The femoral heads are normally positioned within the acetabula. The joint spaces are maintained. No evidence of dislocation is seen. The sacroiliac joints appear symmetrical and unremarkable. The pubic symphysis is normal in width. No obvious soft tissue abnormality or abnormal calcification is noted. Normal study. - No abnormality detected.

Findings:

The bony pelvis shows normal alignment. The iliac bones, sacrum, pubic rami, and acetabula appear intact with no evidence of fracture or lytic or sclerotic lesions.

Both hip joints are well aligned. The femoral heads are normally positioned within the acetabula. The joint spaces are maintained. No evidence of dislocation is seen.

The sacroiliac joints appear symmetrical and unremarkable. The pubic symphysis is normal in width.

No obvious soft tissue abnormality or abnormal calcification is noted. Normal study.

- No abnormality detected.

The pelvic ring is intact with normal alignment. The iliac wings, ischial bones, and superior and inferior pubic rami appear normal in contour and trabecular pattern. No evidence of cortical break, fracture line, or focal bony destruction is identified.

Both acetabula are well formed with preserved cortical margins. The femoral heads are spherical in contour and are well seated within the acetabula bilaterally. No evidence of subluxation or dislocation is seen. The hip joint spaces are maintained symmetrically without significant narrowing.

The sacroiliac joints are symmetrical with maintained joint spaces and no erosions or sclerosis. The sacrum appears intact without obvious fracture or lytic lesions. The pubic symphysis is normal in width without diastasis.

There is no radiographic evidence of osteolytic or osteoblastic lesions. No periosteal reaction is noted. The visualized proximal femoral shafts are unremarkable.

The surrounding soft tissues appear normal. No abnormal calcification or soft tissue swelling is appreciated.

The pelvic ring is intact with normal alignment. The iliac wings, ischial bones, and superior and inferior pubic rami appear normal in contour and trabecular pattern. No evidence of cortical break, fracture line, or focal bony destruction is identified.

Key Images:

Conclusion:

Within normal limits. No radiographic evidence of acute fracture or dislocation. Pelvic bones and bilateral hip joints appear normal. The bony pelvis shows normal alignment. The iliac bones, sacrum, pubic rami, and acetabula appear intact with no evidence of fracture or lytic or sclerotic lesions. Both hip joints are well aligned. The femoral heads are normally positioned within the acetabula. The joint spaces are maintained. No evidence of dislocation is seen. The sacroiliac joints appear symmetrical and unremarkable. The pubic symphysis is normal in width. No obvious soft tissue abnormality or abnormal calcification is noted. Normal study.

No abnormality detected.

No radiographic evidence of acute fracture or dislocation.

No significant bony abnormality detected in the pelvis and bilateral hip joints.

Reported By:**ANJALI**

anjali
MBBS
Signed on 17/2/2026, 11:28:43 am

Approved By:**Lekhana**

Lekhana M N
MBBS
Signed on 17/2/2026, 11:28:50 am