

Literature review on non-spherical particles

Alessio Raimondi

October 3, 2019

Literature review (Drag coefficient)

Before

Correlation for arbitrary shaped particles

1. Ganser [1993]
2. **Haider and Levenspiel [1989]**
3. Hartman [1994]
4. Chien [1994] (
5. Swamme and Ojha [1991]

Performance

Correlation for specific shaped particles

- Brenner [1963]
 - Bowen and Masliyah [1973]
 - Tripathi et al [1994]
 - Militzer et al [1989]
 - Huner and Hussey [1977]
 - Ui et al [1984]
 - Michael [1966]
 - Shail and Norton [1969]
 - Davis [1990]
- } Spheroid
 } Cylinder
 } Disc

C
H
A
B
R
A
1
9
9
9

After

Correlation for arbitrary shaped particles

1. Holzer and Sommerfeld [2008]
2. Tran-Cong et al. [2004]

Correlation for specific shaped particles

- Zastawny et al. [2012]
- Loth [2008]
- Yow [1994]
- Vakil and Green [2011]
- Mando and Rosendahl [2010]

The correlations must take into account:

- ☐ Particle shape.
- ☐ Particle orientation.
- ☐ Particle rotation.

Holzer and Sommerfeld - 2008

- ▶ Huge literature and data review (2061 values)
- ▶ Interpolation of different previous models
- ▶ Arbitrary shape (and Orientation!)
- ▶ Valid for all the Subcritical Regime

$$c_D = \frac{8}{Re} \frac{1}{\sqrt{\Phi_{//}}} + \frac{16}{Re} \frac{1}{\Phi} + \frac{3}{\sqrt{Re}} \frac{1}{\Phi^{\frac{3}{4}}} + 0.4210^{0.4(-\log \Phi)^{0.2}} \frac{1}{\Phi_{\perp}}$$


Holzer and Sommerfeld - 2008

	Haider and Levenspiel [1]	Ganser incl. Leith [2]	Present Eq. (9)	Present Eq. (10)
<i>Mean relative deviation</i>				
Sphere (683 values)	6.59%	10.9%	9.17%	9.17%
Isometric particles (655 values)	6.65%	6.46%	10.5%	10.9%
Cuboids and cylinders (337 values)	42.3%	38.4%	27.2%	29%
Disks and plates (386 values)	$2 \cdot 10^3\%$	$1.8 \cdot 10^3\%$	17.7%	16.8%
All Data (2061 values)	383%	348%	14.1%	14.4%
<i>Maximum relative deviation</i>				
Sphere (683 values)	44%	43%	45%	45%
Isometric particles (655 values)	50%	55%	68%	68%
Cuboids and cylinders (337 values)	$1.1 \cdot 10^3\%$	$1.1 \cdot 10^3\%$	88%	88%
Disks and plates (386 values)	$2.1 \cdot 10^4\%$	$2.4 \cdot 10^4\%$	75%	75%

- ▶ Works better than the other for disks and plates
- ▶ Ganser works better for isometric particles
- ▶ Works well overall

Sanjeevi, Padding and Kuipers – 2015

DNS of a prolate Ellipsoid with $\Phi = 0.886$



Stokes Regime

$$c_D = \frac{8}{Re} \frac{1}{\sqrt{\Phi_{\perp}}} + \frac{16}{Re} \frac{1}{\Phi}$$

- ▶ Leith's (or Ganser's) model – 1993
- ▶ Valid for low Re (≤ 10)
- ▶ For sphere ($\Phi_{\perp} = \Phi_{//} = \Phi$) degenerates to Stokes' analytical solution
- ▶ Modification: $\Phi_{\perp} \rightarrow \Phi_{//}$ (for better approximation of the c_D as a function of the particle orientation)

Newton Regime

Blasius - 1908

Friction drag for plates and disks (small cross-sectional area)

$$c_D = 1.327 \cdot 2 \left(\frac{8}{9}\right)^{\frac{1}{4}} \pi^{\frac{1}{4}} \left(\frac{\text{depth}}{\text{length}}\right)^{\frac{1}{4}} \frac{1}{\Phi^{\frac{3}{4}}} \frac{1}{\sqrt{Re}}$$

for square plates: $c_D = 3.43/(\Phi^{\frac{3}{4}} \sqrt{Re})$

Tran-Cong (2004) and Ganser (1993)

Term for high Re proportional to the projected cross-sectional area (Tran-Cong) with the same factor of proportionality of Ganser.

$$c_D = 0.4210^{0.4(-\log \Phi)^{0.2}} \frac{1}{\Phi_{\perp}}$$

Symbols

Sphericity

$$\Phi = \frac{A_{\text{eq sphere}}}{A_{\text{particle}}}$$

Ratio between the surface area of the volume-equivalent sphere and the area of the actual particle

Crosswise Sphericity

$$\Phi_{\perp} = \frac{A_{\text{eq sphere, cross}}}{A_{\text{particle}, \perp}}$$

Ratio between the cross-sectional area of the volume-equivalent sphere and the projected cross-sectional area of the actual particle

Symbols

Lengthwise Sphericity

$$\Phi_{//} = \frac{A_{\text{eq sphere, cross}}}{\Delta A}$$

$$\Delta A = \frac{A_{\text{particle}}}{2} - \bar{L}_{//}$$

Ratio between the cross-sectional area of the volume-equivalent sphere and the difference between half the surface area and the *average* mean longitudinal projected cross-sectional area of the actual particle.

Since $L_{//}$ depends on the angle of view, an arithmetic average over an entire revolution is used

Recent work – camera-based methods

China - 2017



Correlation	Prediction error		
	r	s_1	s_2
Haider and Levenspiel [23]	13.84%	9.85	2.47
Swamee and Ojha [24]	21.27%	22.22	7.07
Ganser [25]	8.41%	3.46	0.73
Chien [1]	17.27%	13.21	3.23
Yow et al. [29]	47.77%	130.24	/
Holzer and Sommerfeld [36]	11.07%	5.75	1.32
Our equation	4.91%	1.45	0.30

Limitations:

- Only three major regular particle shapes investigated (sphere, cube, cylinder)

Recent work – camera-based methods

Italy (Bari) - 2015



Limitations:

- ▶ Small data sample
- ▶ No comparison with Holzer and Sommerfeld (although good agreement with Ganser)

To do list

- ▶ Read Loth-2008 (Snow??)
- ▶ Replicate the work of the workshop on SU2
- ▶ (Try the H&S formula on the Italian data)