

طراحی الگوریتم ها

ملکی مجد

مباحث

حل مساله بیشترین جریان

Maximum Flow

Max-flow min-cut theorem

Ford-Fulkerson method


Edmonds-Karp algorithm

Maximum bipartite matching

مبحث *Maximum Flow* از فصل ۲۶ کتاب CLRS تدریس می شود.

Maximum Flow

+a graph-theoretic definition of flow networks



A **flow network** $G = (V, E)$:

is a directed graph in which each edge $(u, v) \in E$ has a nonnegative **capacity** $c(u, v) \geq 0$.

If $(u, v) \notin E$, we assume that $c(u, v) = 0$.

In Flow networks


We distinguish : a **source** s and a **sink** t

we assume :

every vertex lies on some path from the source to the sink

The graph is therefore connected, and $|E| \geq |V| - 1$.

Example of network



flow


A flow in G is a **real-valued function f** : $V \times V \rightarrow R$ that satisfies the following three properties:

- **Capacity constraint:** For all $u, v \in V$, we require $f(u, v) \leq c(u, v)$.
- **Skew symmetry:** For all $u, v \in V$, we require $f(u, v) = -f(v, u)$.
- **Flow conservation:** For all $u \in V - \{s, t\}$, we require $\sum_{v \in V} f(u, v) = 0$
flow in equals **flow out** for vertex other than source and sink

The **value** of a flow f :

total flow out of the source ($|f| = \sum_{v \in V} f(s, v)$)

A sample flow



maximum-flow problem

In the ***maximum-flow problem***, we are given a flow network G with source s and sink t , and we wish to **find a flow of maximum value**.

Networks with multiple sources and sinks

- This problem is no harder than ordinary maximum flow
 - We can **reduce the problem** of determining a maximum flow in a network with multiple sources and multiple sinks to an ordinary maximum-flow problem


- **Add following:**

a ***supersource*** s

add a directed edge (s, s_i) with capacity $c(s, s_i) = \infty$ for each source s_i

a ***supersink*** t

and add a directed edge (t_i, t) with capacity $c(t_i, t) = \infty$ for each sink t_i



lemma

$$f(X, Y) = \sum_{x \in X} \sum_{y \in Y} f(x, y)$$

Let $G = (V, E)$ be a flow network, and let f be a flow in G . Then the following equalities hold:

1. For all $X \subseteq V$, we have $f(X, X) = 0$.
2. For all $X, Y \subseteq V$, we have $f(X, Y) = -f(Y, X)$.
3. For all $X, Y, Z \subseteq V$ with $X \cap Y = \emptyset$, we have the sums $f(X \cup Y, Z) = f(X, Z) + f(Y, Z)$ and $f(Z, X \cup Y) = f(Z, X) + f(Z, Y)$.

The value of flow

$$\begin{aligned}|f| &= f(s, V) \\&= f(V, V) - f(V - s, V) \\&= -f(V - s, V) \\&= f(V, V - s) \\&= f(V, t) + f(V, V - s - t) \\&= f(V, t)\end{aligned}$$



The Ford-Fulkerson method
+for solving the maximum-flow problem

Residual networks

- The amount of *additional flow* we can push from u to v before exceeding the capacity $c(u, v)$ is the **residual capacity** of (u, v) , given by

$$c_f(u, v) = c(u, v) - f(u, v).$$

given a flow network and a flow, the residual network consists of edges that can admit more flow




how a flow in a residual network relates to a flow in the original flow network

Let $G = (V, E)$ be a flow network with source s and sink t , and let f be a flow in G . Let G_f be the residual network of G induced by f , and let f' be a flow in G_f . Then the flow sum $f + f'$ defined by equation (26.4) is a flow in G with value $|f + f'| = |f| + |f'|$.

Augmenting paths

Augmenting path is

a path from the source s to the sink t along which we can send more flow, and then augmenting the flow along this path



- ❖ A flow is maximum if and only if its residual network contains no augmenting path.

Residual capacity of an augmenting path

- The maximum amount by which we can increase the flow on each edge in an augmenting path p

$$c_f(p) = \min \{c_f(u, v) : (u, v) \text{ is on } p\}$$

residual capacity of an augmenting path can add to the value of flow

Let $G = (V, E)$ be a flow network, let f be a flow in G , and let p be an augmenting path in G_f . Define a function $f_p : V \times V \rightarrow \mathbf{R}$ by

$$f_p(u, v) = \begin{cases} c_f(p) & \text{if } (u, v) \text{ is on } p , \\ -c_f(p) & \text{if } (v, u) \text{ is on } p , \\ 0 & \text{otherwise .} \end{cases} \quad (26.6)$$

Then, f_p is a flow in G_f with value $|f_p| = c_f(p) > 0$.

Corollary

Let $G = (V, E)$ be a flow network, let f be a flow in G , and let p be an augmenting path in G_f . Let f_p be defined as in equation (26.6). Define a function $f' : V \times V \rightarrow \mathbf{R}$ by $f' = f + f_p$. Then f' is a flow in G with value $|f'| = |f| + |f_p| > |f|$.

General method

How increase the value of flow


FORD-FULKERSON-METHOD(G, s, t)

- 1 initialize flow f to 0
- 2 **while** there exists an augmenting path p
- 3 **do** augment flow f along p
- 4 **return** f

Definition of cut

- A ***cut*** (S, T) of flow network $G = (V, E)$ is a partition of V into S and $T = V - S$ such that $s \in S$ and $t \in T$.
- If f is a flow, then the ***net flow*** across the cut (S, T) is defined to be $f(S, T)$.
- The ***capacity*** of the cut (S, T) is $c(S, T)$.
- A ***minimum cut*** of a network is a cut whose capacity is minimum over all cuts of the network.

A cut with
the net flow $f(S, T) = 19$, and
the capacity $c(S, T) = 26$.




- The value of any flow f in a flow network G is bounded from above by the capacity of any cut of G .
(can proved by definition of flow and cut)

Max-flow min-cut theorem

If f is a flow in a flow network $G = (V, E)$ with source s and sink t , then the following conditions are equivalent:

1. f is a maximum flow in G .
2. The residual network G_f contains no augmenting paths.
3. $|f| = c(S, T)$ for some cut (S, T) of G .



Max-flow min-cut theorem

If f is a flow in a flow network $G = (V, E)$ with source s and sink t , then the following conditions are equivalent:

1. f is a maximum flow in G .
2. The residual network G_f contains no augmenting paths.
3. $|f| = c(S, T)$ for some cut (S, T) of G .

Corollary


The value of any flow f in a flow network G is bounded from above by the capacity of any cut of G .


basic Ford-Fulkerson algorithm


expands on the FORD-FULKERSON METHOD


FORD-FULKERSON(G, s, t)


- 1 **for** each edge $(u, v) \in E[G]$
- 2 **do** $f[u, v] \leftarrow 0$
- 3 $f[v, u] \leftarrow 0$
- 4 **while** there exists a path p from s to t in the residual network G_f
- 5 **do** $c_f(p) \leftarrow \min \{c_f(u, v) : (u, v) \text{ is in } p\}$
- 6 **for** each edge (u, v) in p
- 7 **do** $f[u, v] \leftarrow f[u, v] + c_f(p)$
- 8 $f[v, u] \leftarrow -f[u, v]$



















Time complexity


- The running time of FORD-FULKERSON depends on how the augmenting path p in line 4 is determined.
- where is the maximum flow found by the algorithm:
a straightforward implementation runs in time $O(E |f^*|)$
- Prove Hint:
the flow value increases by at least one unit in each iteration

$O(E |f^*|)$ can be bad!


- continue, choosing the augmenting path $s \rightarrow u \rightarrow v \rightarrow t$ in the odd-numbered iterations and the augmenting path $s \rightarrow v \rightarrow u \rightarrow t$ in the even-numbered iterations.



(a)



(b)



(c)

The Edmonds-Karp algorithm

- The bound on FORD-FULKERSON can be improved if we implement the computation of the augmenting path p in line 4 **with a breadth-first search** (each edge has unit distance (weight))
the augmenting path is a *shortest* path from s to t in the residual network

the running time of the Edmonds-Karp algorithm is $O(V E^2)$

the running time of the Edmonds-Karp algorithm is $O(V E^2)$


Lemma 26.8

If the Edmonds-Karp algorithm is run on a flow network $G = (V, E)$ with source s and sink t , then for all vertices $v \in V - \{s, t\}$, the shortest-path distance $\delta_f(s, v)$ in the residual network G_f increases monotonically with each flow augmentation.

Theorem 26.9

If the Edmonds-Karp algorithm is run on a flow network $G = (V, E)$ with source s and sink t , then the total number of flow augmentations performed by the algorithm is $O(VE)$.

Show the execution of the Edmonds-Karp algorithm on the flow network



Sample Problem

Suppose you are given a flow network G with *integer* edge capacities and an *integer* maximum flow f^* in G . Describe algorithms for the following operations:

- (a) INCREMENT(e): Increase the capacity of edge e by 1 and update the maximum flow.
- (b) DECREMENT(e): Decrease the capacity of edge e by 1 and update the maximum flow.

Both algorithms should modify f^* so that it is still a maximum flow, more quickly than recomputing a maximum flow from scratch.


Maximum bipartite matching
+solve using flow network

The maximum matching problem

Given an undirected graph $G = (V, E)$,

Matching is a subset of edges $M \subseteq E$ such that for all vertices $v \in V$, at most one edge of M is incident on v .


Maximum matching is a matching of maximum cardinality



a vertex $v \in V$ is **matched** by matching M if some edge in M is incident on v ; otherwise, v is **unmatched**.

The maximum-bipartite-matching problem

- **Bipartite graphs:** the vertex set can be partitioned into $V = L \cup R$, where L and R are disjoint and all edges in E go between L and R .
(further assume that every vertex in V has at least one incident edge.)



Finding a maximum bipartite matching

- We can use the **Ford-Fulkerson method** to find a maximum matching in an **undirected bipartite graph** $G = (V, E)$ in time **polynomial in $|V|$ and $|E|$** .
- First:
define the **corresponding flow network** $G' = (V', E')$ for the bipartite graph G



corresponding flow network

- Add source s and sink t as new vertices
 - let $V' = V \cup \{s, t\}$.
- The directed edges of G' are :
 - the edges of E , directed from L to R ($V = L \cup R$), along with V new edges:

$$\begin{aligned} E' &= \{(s, u) : u \in L\} \\ &\quad \cup \{(u, v) : u \in L, v \in R, \text{ and } (u, v) \in E\} \\ &\quad \cup \{(v, t) : v \in R\} . \end{aligned}$$

- assign unit capacity to each edge in E' .

The flow network corresponding to a bipartite graph



The order of E' ?

$$|E| \leq |E'|$$

$$|E'| = |E| + |V| \leq 3|E|, (|E| \geq |V|/2)$$

$$so |E'| = \Theta(E)$$

a matching in G corresponds directly to
a flow in G 's corresponding flow network

Lemma 26.10

Let $G = (V, E)$ be a bipartite graph with vertex partition $V = L \cup R$, and let $G' = (V', E')$ be its corresponding flow network. If M is a matching in G , then there is an integer-valued flow f in G' with value $|f| = |M|$. Conversely, if f is an integer-valued flow in G' , then there is a matching M in G with cardinality $|M| = |f|$.

maximum matching in a bipartite graph and
the value of a maximum flow

Theorem 26.11 (Integrality theorem)

If the capacity function c takes on only integral values, then the maximum flow f produced by the Ford-Fulkerson method has the property that $|f|$ is integer-valued. Moreover, for all vertices u and v , the value of $f(u, v)$ is an integer.


Corollary 26.12

The cardinality of a maximum matching M in a bipartite graph G is the value of a maximum flow f in its corresponding flow network G' .


Whole picture

- Given a bipartite undirected graph G ,
 - Create the flow network G'
 - Run the Ford-Fulkerson method
Ford-Fulkerson time complexity = $O(E |f^*|)$
 - Obtaining a maximum matching M from the integer-valued maximum flow
 - The value of the maximum flow in G' is $O(V)$.
 - Since any matching in a bipartite graph has cardinality at most $\min(L, R) = O(V)$,
- Time complexity of finding a maximum matching in a bipartite graph ($|E'| = \Theta(E)$) is $O(V E)$.


Compute the maximum bipartite matching




Compute the maximum bipartite matching




Compute the maximum bipartite matching




Compute the maximum bipartite matching



Compute the maximum bipartite matching



Compute the maximum bipartite matching



Some sample problems

- 1- If all edges in a graph have distinct capacities, is there a unique maximum flow?
- 2- Let $G = (V, E)$ be a bipartite graph with vertex partition $V = L \cup R$, and let G' be its corresponding flow network. Give a good upper bound on the length of any augmenting path found in G' during the execution of FORD-FULKERSON.
- 3- A ***perfect matching*** is a matching in which every vertex is matched.
Let $G = (V, E)$ be an undirected bipartite graph with vertex partition $V = L \cup R$, where $|L| = |R|$. For any $X \subseteq V$, define the ***neighborhood*** of X as $N(X) = \{y \in V : (x, y) \in E \text{ for some } x \in X\}$, that is, the set of vertices adjacent to some member of X .
Prove ***Hall's theorem***: there exists a perfect matching in G if and only if $|A| \leq |N(A)|$ for every subset $A \subseteq L$.