

The bilaterian forebrain: an evolutionary chimaera

Maria Antonietta Tosches and Detlev Arendt

The insect, annelid and vertebrate forebrains harbour two major centres of output control, a sensory-neurosecretory centre releasing hormones and a primordial locomotor centre that controls the initiation of muscular body movements. In vertebrates, both reside in the hypothalamus. Here, we review recent comparative neurodevelopmental evidence indicating that these centres evolved from separate condensations of neurons on opposite body sides ('apical nervous system' versus 'blastoporal nervous system') and that their developmental specification involved distinct regulatory networks (apical *six3* and *rx* versus mediolateral *nk* and *pax* gene-dependent patterning). In bilaterian ancestors, both systems approached each other and became closely intermingled, physically, functionally and developmentally. Our 'chimeric brain hypothesis' sheds new light on the vast success and rapid diversification of bilaterian animals in the Cambrian and revises our understanding of brain architecture.

Addresses

European Molecular Biology Laboratory, Developmental Biology Unit, Meyerhofstrasse 1, 69012 Heidelberg, Germany

Corresponding author: Arendt, Detlev (arendt@embl.de)

Current Opinion in Neurobiology 2013, **23**:1080–1089

This review comes from a themed issue on **Development of neurons and glia**

Edited by **Samuel Pfaff** and **Shai Shaham**

For a complete overview see the [Issue](#) and the [Editorial](#)

Available online 27th September 2013

0959-4388/\$ – see front matter, Published by Elsevier Ltd.

<http://dx.doi.org/10.1016/j.conb.2013.09.005>

Introduction

Vertebrates, insects and annelid worms share the presence of a centralized nervous system (CNS) with an anterior brain. The brain integrates sensory information to control the animal body through distinct effector systems [1]. First, neurosecretory centres in the brain release hormones that control the general behavioural state, homeostasis and life cycle transitions. Second, the brain is the higher-order control centre for the hierarchically organized somatic motor system, that controls muscular movements. This review proposes a scenario for the evolutionary origin of these fundamentally distinct control centres.

Most instructive in this context is the analysis of regionalization of the brain neuroectoderm, which is generated by the intersection of the 'apical' and 'mediolateral'

patterning systems (Figure 1). We will reason that these systems were initially separate, specifying 'apical' (Figure 1a) versus 'blastoporal' (Figure 1b) identities in prebilaterian ancestors, and started to overlap in the bilaterian lineage only. We will then review evidence linking the apical system to the specification of an 'apical nervous system' in several bilaterians, comprising sensory-neurosecretory protoneurons, that perceive light and other stimuli to control the release of neuropeptides and hormones. The circadian and hormonal release centre in the bilaterian forebrain can thus be regarded as the 'legacy' of an ancient apical nervous system. In contrast, the mediolateral system governs the differential specification of various other types of sensory neurons, interneurons and motoneurons that constitute the 'blastoporal nervous system'. The latter is composed of axonal circuits for the control of body movements and referred to as 'blastoporal legacy'.


In the course of bilaterian evolution, the most anterior part of the blastoporal nervous system, the primordial locomotor control centre, approached, and finally merged with the sensory-neurosecretory centre of the apical nervous system. This way, an evolutionary chimaera of neural tissues evolved, combining two distinct effector systems of different origin.

A fusion of apical nervous system and anterior blastoporal nervous system components has been previously put forward for the evolution of the protostome brain, based on morphological evidence [2]. Taking into account new molecular data, we propose here that this fusion occurred earlier in animal evolution, predating the bilaterian ancestors.

The apical and mediolateral patterning systems evolved at opposite body sides

The apical system, also referred to as 'head patterning system', involves antagonistic Wnt and FGF signalling. It specifies the tissue around the apical pole (i.e., the non-blastoporal side of the body) in insect [3,4], annelid [3,5], brachiopod [6] echinoderm [7], hemichordate, [8,9], amphioxus [10] and vertebrate development [11] (Figure 1a). The apical system controls the expression of the region-specific transcription factors *six3*, *rx*, *foxQ2* and *fezf* in radial and partially overlapping domains (Figure 1a). Recent data from a bilaterian outgroup, the cnidarians (comprising, e.g. sea anemone and jellyfish), indicates that this patterning system predates bilaterians [12,13]. In insects, vertebrates and annelids, the *six3*+ apical region gives rise to the most anterior part of the

Figure 1



The apical and the blastoporal patterning systems. Expression of nervous system regionalization genes in Cnidaria, annelids, vertebrates and insects. **(a)** The apical patterning system comprises concentric expression of *six3*, *rx*, *fezf* and *foxQ2*. **(b)** The blastoporal patterning system establishes the expression of transcription factors centred around the blastopore. *Dll* is expressed at the interface between the two systems. **(c)** The apical (blue) and blastoporal (red) patterning systems are active on opposite body sides in Cnidaria, but they fuse in bilaterians at the level of the anterior forebrain. The expression of *foxA* (grey) demarcates the blastopore in Cnidaria, and the neural midline in Bilateria (plus the blastopore lips in protostomes, see [20]). Cnidaria expression data are drawn from *Nematostella* [12^{**},13^{**},95,96] and *Acropora* [97] (lateral views), annelid expression data from *Platynereis* [5,17,98] (blastoporal view in (a) and (b), blastoporal-lateral view in (c)), vertebrate expression data from *Xenopus* [99–103] (blastoporal view in (a) and (b), blastoporal-lateral view in (c)), and insect expression data from *Tribolium* [3^{*},4,16,104] (blastoporal view). *FoxQ2* expression has never been characterized in vertebrates and insects.

brain [3^{*}]. In vertebrates, this region includes the telencephalon and anterior hypothalamus [14].

The mediolateral system likewise represents ancient bilaterian heritage, involving midline Hedgehog and lateral BMP signalling. In vertebrates, insects and annelids this system controls the staggered expression of medial *foxA*, *sim* and *nk*, and lateral *pax*, *msx* and *dll* genes (Figure 1b) [15–17]. In the vertebrate neuroectoderm, these transcription factors specify the floor, basal, alar and

roofplate subdivisions of the folded neural tube that extend into the forebrain [18]. In annelids, the same sequence of genes establishes a bilateral sequence of molecular regions that are centred along the neural midline (Figure 1b). Given that in annelids and other protostomes the neural midline emerges from the median fusion of the blastopore lips [19], it has been suggested that the mediolateral system was originally active alongside the blastopore (and that this spatial connection was lost in vertebrate ancestors due to changes in the mode of

gastrulation) [20]. Recent data from Cnidaria support this hypothesis, revealing that *dll*, *msx*, *pax3/7* and *nk2* genes are expressed in similar sequence around the blastopore [21], together with the medial signal *hedgehog* [22] (Figure 1b). Bilateral brain patterning thus involves two systems that were initially separate and started to overlap in bilaterian ancestors only (Figure 1c). In the subsequent paragraphs we will reason that both systems play conserved roles in the specification of fundamentally different nervous system components that evolved at opposite body sides. Comparative data allows sorting out the ‘original’ apical versus mediolateral neural cell fates and thus reconstructing the constituents that were involved in the formation of the chimeric bilaterian brain.

The apical nervous system: an assembly of sensory-neurosecretory protoneurons

Marine invertebrates are particularly useful to understand the apical nervous system component of the bilaterian brain. Their ciliated planktonic larvae develop nervous structures (including an ‘apical organ’) from the *six3*+ apical plate, downstream of the apical patterning system [3,6]. These larval structures are in place before the blastoporal mediolateral patterning system establishes the axonal circuits for the control of body musculature (that are fully developed only after metamorphosis). In species with a gradual, non-catastrophic metamorphosis, large part of the apical plate remains and becomes incorporated into the adult brain, referred to as cerebral ganglia. That means that in marine ciliated larvae we can study components of the apical nervous system in a more ‘puristic’ (meaning pre-fusion) form.

In these larvae, the most prevalent apical plate cell types are multimodal sensory-neurosecretory cells, also referred to as protoneurons [23]. These cells usually exhibit a flask-shaped morphology with ciliated dendritic endings [24]. Based on ultrastructural features but also molecular data, these cells closely resemble the cerebrospinal-fluid (CSF)-contacting neurons of the vertebrate brain [23,25] (Figure 2a). Ciliary-type photoreceptors employing c-opsins, orthologs of vertebrate rod and cone opsin, have also been identified in the annelid apical plate [31], where they develop under the control of *rx* (MAT and DA, unpublished). C-opsin expression in the apical plate of brachiopods [32] and echinoderms [33•] confirms that non-visual photoreception is a major task of the larval apical nervous system. Usually, an apical plexus forms underneath the apical organ [24,26], which functions at the same time as a neural integration site and a neuroendocrine release site [24] (Figure 2a). These data suggest that protoneuron-like cells, including ciliary photoreceptors, in conjunction with a neurosecretory plexus represent ancient apical nervous system-derived components. This notion finds support by a comparative survey of *six3* and *rx*-dependent cell types in the adult annelids and vertebrates.

The ‘apical legacy’ of the bilaterian forebrain: (1) non-visual light perception and the molecular clock


Rx and *six3*, key transcription factors of the apical patterning system, were initially identified as regulators of retinal development [14,27], but later studies highlighted their essential roles in postmitotic photoreceptors. *Rx* directly controls the expression of *opsins* and other photoreceptor-specific genes through binding to a photoreceptor-specific regulatory element (PCE-1) [28,29]. Similarly, *six3* directly controls *rhodopsin* expression [30]. These data corroborate our notion that *six3* and *rx* played ancient roles in ciliary photoreceptor cell specification and differentiation [31] and suggest that vertebrate rods and cones evolved from apical plate non-visual photoreceptors. Consistently, in amphioxus, the frontal eye, the proposed homolog of vertebrate retina, develops from *rx*+ *six3*+ progenitors; *six3* is expressed in the differentiated row 1 photoreceptors (and *rx* in frontal eye row 3 cells) [34••].

Rx and *six3* are also expressed in the developing vertebrate pineal [35,36], the proposed evolutionary ‘sister’ of the retina [37,38••]. In non-mammalian vertebrates, the pineal likewise contains ciliary-type photoreceptors employing c-opsins. Also, it entrains an endogenous clock and in all vertebrates produces melatonin, the main output of the circadian system. *Rx* expression in the pineal shows diurnal oscillations in adults, suggesting a possible role in the rhythmic expression of target genes [39]. Both the retina and the pineal represent evaginations of the forebrain and can thus be regarded part of the adult apical nervous system.

Finally, in the anterior hypothalamus the suprachiasmatic nucleus (the master clock that integrates inputs from the retina and the pineal) develops under the control of *six3*, its paralog *six6*, and *rx*, and rhythmically expresses *six3* and *six6* in adults [41••,42]. These data indicate that the anterior hypothalamus, together with the ciliary photoreceptors of retina and pineal, represents ‘apical legacy’ of the vertebrate forebrain.

Supporting our notion that luminance detection, the control of physiology and an endogenous central molecular clock were directly coupled in multifunctional cells in the urbilaterian forebrain, we have recently found that *rx*-dependent ciliary photoreceptor cells in the annelid forebrain (that persist from the larval apical plate) run a circadian clock and release melatonin as an effector. Moreover, specific markers of the vertebrate anterior hypothalamus are also expressed in the annelid apical plate ([25] and MAT and DA, unpublished), corroborating our hypothesis that vertebrate pineal, retina and anterior hypothalamus evolved from apical plate non-visual photoreceptors after ‘division of labour’ [40].

Figure 2



Apical and blastoporal 'legacies' in bilaterian brains. **(a)** Comparison of the annelid apical plate (left) and the vertebrate diencephalon (right). Annelid drawing based on [5,24,25] and unpublished data; vertebrate scheme redrawn from [105]. Hyp, hypothalamus; me, median eminence; Pi, pineal; Pp, parapineal; R, retina. **(b)** Comparison of the neural architecture that underlies the control of the trunk locomotor pattern generators by descending projections from ventral *nk2+* forebrain regions (yellow). Based on [76,77,106]. Cc, circumoesophageal connectives; sog, subesophageal ganglion; fe, frontal eye; ppn2, preinfundibular projection neuron 2; pinfn, postinfundibular neuropile; PMC, primary motor centre; Hyp/PTu, hypothalamus/posterior tuberculum.

The 'apical legacy' of the bilaterian forebrain: (2) the hormonal release centre

Expression of *rx* in the anterior neural plate also includes prospective neurosecretory areas of the vertebrate hypothalamus. Here, *rx* determines cell fates as *rx* mutant mice fail to develop the posterior pituitary [43] and conditional *rx* elimination under a *Six3::Cre* driver results in the loss of the arcuate and the ventromedial nuclei of the hypothalamus [44^{***}]. Consistently, *rx*⁺ stem cells are able to generate vasopressinergic and NPY⁺ neurons *in vitro*, but cannot express posterior hypothalamic markers [45]. In annelids *rx* and *six3* are likewise expressed in neurosecretory cells that project into the apical plexus (see

above); and in insects they are expressed in the brain neurosecretory centres (pars intercerebralis and pars lateralis, see [3^{*},46]). Another apical nervous system marker, *fezf*, also contributes to the specification of neuroendocrine fates: in zebrafish, *fezf* acts upstream of *otp* to control the differentiation of isotocinergic neurons [47]; similarly, vasotocin (the vasopressin/isotocin ortholog) is expressed in *otp*⁺ flask-shaped cells of the annelid forebrain [25].

The similarity of invertebrate and vertebrate neurohormonal sites [48] is highlighted by the conservation of some of the neuropeptides and hormones released in these structures, like the earlier mentioned vasotocin and

melatonin (the pineal is a neurohemal organ [49]). Moreover, RFamideergic neurons are characteristic of the apical plate and apical plexus of annelids, molluscs, hemichordates and cnidarians [25,50–52], of the insect pars intercerebralis [46] and of the vertebrate hypothalamus [53]. Finally, most of the neurohormones and neuropeptides of the vertebrate arcuate nucleus-median eminence complex, like GnRH, NPY and enkephalins, have ancient evolutionary origins [54] and might be present in the apical plate of various invertebrates, as suggested by cross-reactive antibodies [55,56]. We thus propose that the hormonal release centre represents a characteristic apical nervous system derivative of the bilaterian forebrain.


The blastoporal nervous system: an axonal system comprising lateral sensory-integrative and medial motor regions

In contrast to the apical nervous system, which, in essence, can be traced back to protoneuron-like sensory cells specialized on neurosecretory output, the blastoporal nervous system is functionally based on axonal connections between its components. It is best described as a sensory-contraction system, composed of axonal circuits that integrate sensory information to control body musculature. Across bilaterians, the blastoporal nervous system is specified by the mediolateral patterning system, which establishes a lateral sensory-integrative region

comprising sensory neurons and interneurons, and a medial motor region comprising interneurons and motoneurons, in annelids [17] as well as in vertebrates [15]. Importantly, recent comparative data indicate that a medial *foxA*+, *nk2*+ contractile-motor region alongside the blastopore and a more lateral *msx*+, *pax3/7*+, *Dll*+ sensory-integrative region in a distance to the blastopore also exist, albeit in a less centralized form, in the cnidarian outgroup (reviewed in [21]). This would imply that the functional link between mediolateral patterning and neural cell type specification predates the cnidarian-bilaterian ancestor (and that the blastoporal and apical nervous systems indeed evolved on opposite body sides, Figure 3).

Underscoring the essential nature of targeted axonal connections for information transfer in the blastoporal nervous system, transcription factors that play conserved roles in the control of cell fates have proven indispensable for the establishment of axonal connection between post-mitotic neurons. For example, *foxA*, *pax6* and *nk2* directly control the expression of guidance factors important for axonal outgrowth and the formation of major axonal tracts [57–62,63*,64*]. Also, in mice and fishes mutant for the transcription factor *sim*, medial neurons are generated correctly but fail to establish proper axonal projections [65,66*]; likewise, *Drosophila sim* is essential for axogenesis [67].

Figure 3



The chimeric brain hypothesis. In the scheme, the apical nervous system is in blue, and the motor and the sensory-integrative components of the blastoporal nervous system are represented in yellow and red, respectively. In the last common ancestor of Cnidaria and Bilateria (neuralian ancestor) the two systems were separate, and concentrated respectively around the apical pole and the blastopore. With the transition to the urbilaterian (the ancestor of all Bilateria) the apical nervous system and anterior side of the blastoporal nervous system merged to form the forebrain, as found in extant annelids and vertebrates.

A 'blastoporal legacy' of the bilaterian forebrain: the primordial locomotor control centre


Our focus is on the most anterior extension of the medial motor region that became part of the ventral bilaterian forebrain. The transcription factors *nk2.1*, *nk2.2* and *sim* demarcate this region in vertebrates, ascidians, amphioxus and annelids [25,68–71]; notably, an anterior ventromedian stripe of *nk2.1* expression is also found in other marine invertebrates including hemichordates, indicating broad evolutionary conservation [6,8,25,72]. In vertebrates, the *nk2+* forebrain region comprises the subprethalamal basal plate and the tuberal and caudal hypothalamus, including the mammillary bodies [70,73,74*] (that are absent in mouse *nkx2.1* mutants [1,73]). These regions are known to integrate rostral sensory inputs and initiate and modulate locomotion via axons projecting caudally into the brainstem and the spinal cord [1,75] (Figure 2b). Similarly, in amphioxus the motoneurons and interneurons of the primary motor centre are innervated by preinfundibular projection neurons and by fibres from the tegmental or postinfundibular neuropil (a site of integration of rostral sensory inputs), both located in the *nk2.1/nk2.2* region of the cerebral vesicle [76] (Figure 2b). In insects and annelids, large axons descend from the posterior brain to the trunk and contact directly or indirectly trunk motoneurons [77,78*]. In insects, these axons emerge from the central complex, a series of midline neuropils that represents the highest motor control centre. In annelids, the corresponding neurons appear to be part of the *nk2.1* domain and their precise molecular identity is currently investigated (MAT, unpublished).

In line with evolutionary conservation of primordial locomotor control centres in the ventromedian bilaterian forebrain, the *nk2.1+* brain regions contain dopaminergic neurons, in vertebrates, amphioxus, ascidians and annelids [79,80]. The role of dopaminergic neurons in locomotor control is well established in vertebrates and invertebrates [81], but further studies are needed to test the homology of these neuromodulatory neuronal circuits throughout Bilateria.

The chimeric brain hypothesis

So far, and for clarity, our effort has been to sort out the 'apical' and 'blastoporal' legacies of the bilaterian forebrain. Figure 4 summarizes and illustrates both hypothetical contributions on a fate map of the anterior vertebrate neural plate. Notably, as a result of the fusion of neurosecretory and locomotor control centres that took place more than six hundred million years ago in prebilaterian ancestors, these components are partially intermingled and closely interconnected in today's brains, to an extent that it becomes sometimes difficult to tell them apart and draw clear boundaries (striped areas in Figure 4).

Figure 4



The apical and blastoporal contributions to the vertebrate forebrain. Generalized fate map of the vertebrate neural plate, illustrating the apical and blastoporal legacies of the vertebrate brain as deduced from the expression of key transcription factors and the comparison of cell types. The fate map is based on [107,108]. Abbreviations: Ad, adenohypophysis; aHyp, anterior hypothalamus; cHyp, caudal hypothalamus; ETh, eminentia thalamica; Ha, habenula; Pi/PP, pineal/parapineal; PTu, posterior tuberculum; PT, pretectum; PTh, prethalamus; Tel, telencephalon; Th, thalamus; tHyp, tuberal hypothalamus; ZLI, zona limitans intrathalamica.

In line with this, characteristic apical and mediolateral system genes such as *six3*, *rx* versus *nk2.1*, *pax6* are broadly co-expressed in the developing bilaterian brain. For example, despite being necessary only for the development of the rostral forebrain [82], *six3* expression extends posteriorly into the thalamus and the midbrain during late embryonic development [83] (where deep brain photoreceptors have been described [84]). Also, *nk2.1* expression has expanded anteriorly into the former 'apical' territory. Such expansion might have changed (or complemented) the neurosecretory cell fate towards that of projection neuron, thus contributing to the functional fusion of the two effector systems. Intriguingly, the anterior limits of *nkx2.1* and *nkx2.2* expression are variable and appear to expand even within vertebrates into brain regions that had not expressed them in early vertebrates, insofar as the entire expression domain of *shh* and *nkx2.1* in the ventral telencephalon probably arose *de novo* after the separation of cyclostomes and gnathostomes [74*,85].

To track these variations and to reconstruct ancient brain subdivisions, a systems-level analysis can be more powerful than the analysis of single genes. On the basis of the combinatorial analysis of hundreds of transcription factors, a new interpretation of the vertebrate hypothalamus has been recently proposed [86**], and according to this study, only the caudal hypothalamus (mammillary region

and retromammillary areas) is considered a true basal plate derivative. This new interpretation is in agreement with our evolutionary scenario, which considers the caudal hypothalamus as the anterior part of the blastoporal nervous system basal plate (Figure 4).

Despite the paucity of comparative data, some examples of the chimeric nature of the bilaterian forebrain can be recognized both at the tissue and at the cell type level.

At the tissue level, the fusion of 'apical' and 'blastoporal' legacies would involve the functional integration of cell types of different origin, through the establishment of new synaptic connections. For instance, while the ciliary photoreceptors of the vertebrate retina constitute 'apical' legacy (see above), the ganglion cells appear to be of 'blastoporal' origin, as suggested by their dependence on 'blastoporal' transcription factors such as *pax6* and *brn3* [87] and by the presence of rhabdomeric photoreceptors along the entire trunk neuroectoderm of protostomes and deuterostomes [88].

At the cell type level, the interesting perspective emerges that neural cells that initially developed under the unique influence of one patterning system, started to receive signals from the other. Such newly received signals might have activated downstream target genes belonging to the respective other system and might have led, at least in part, to 'chimeric' cell fates, according to a cell type neofunctionalization scenario. Indicative of truly chimeric cell fates, neurosecretory and caudally projecting neurons of the same peptidergic type coexist in the tuberal hypothalamus [89,90]. In the same region, *otp* is responsible for the differentiation of caudally projecting dopaminergic neurons of the diencephalospinal tract, that control the maturation of locomotor patterns [90,91]; moreover, light-sensitive *otp+* neurons have been recently implicated in dark photokinesis [92]. Intriguingly, both 'blastoporal' and 'apical' transcription factors, like *sim* and *fezf*, act in parallel with *otp* to specify these neuronal types [47,90,93], suggesting that different cellular properties, like the neurosecretory fate or establishment of axonal connections, might be controlled by distinct gene regulatory modules acting within neighbouring or even in one and the same cell [94**].

Concluding remarks

If the apical neurosecretory versus blastoporal contractile-motor centres indeed evolved on opposite body sides, what was the driving force to establish contact and finally fuse at the new front end of bilaterian ancestors? We speculate that one selective advantage might have been that the neurosecretory release centre obtained more and more direct control on the newly evolving locomotor control centre. Also, sensory modalities specific of the apical system might have been secondarily recruited for the control of locomotion, via newly evolving direct

connections to the muscular motor circuits. It is conceivable that the combination of 'division of labour' and cell type neofunctionalization (with the parallel activation of 'apical' and 'blastoporal' differentiation programmes in the same cell) has boosted cell type diversity in the evolving bilaterian forebrain, and triggered the establishment of new neural circuits between previously non-related parts of the nervous system. These enabled an unprecedented diversification in the divergent bilaterian lineages and led to a true 'Cambrian explosion' for bilaterian brain evolution.

Acknowledgements

We thank all members of the Arendt lab for numerous valuable and inspiring discussions. Our work on *Platynereis* brain development has been partially funded by the ERCEA-AdG-2011-294810_BrainEvoDevo.

References and recommended reading

Papers of particular interest, published within the period of review, have been highlighted as:

- of special interest
- of outstanding interest

1. Swanson LW: **Cerebral hemisphere regulation of motivated behavior.** *Brain Res* 2000, **886**:113-164.
2. Nielsen C: **How to make a protostome.** *Invertebr. Syst.* 2012, **26**:25-40.
3. Steinmetz PRH, Urbach R, Posnien N, Eriksson J, Kostyuchenko RP, Brena C, Guy K, Akam M, Bucher G, Arendt D: **Six3 demarcates the anterior-most developing brain region in bilaterian animals.** *EvoDevo* 2010, **1**:14.
- In this paper, Steinmetz *et al.* demonstrate the conservation of anterior *six3* expression in a broad range of protostomes.
4. Posnien N, Koniszewski NDB, Hein HJ, Bucher G: **Candidate gene screen in the red flour beetle *Tribolium* reveals *six3* as an ancient regulator of anterior median head and central complex development.** *PLoS Genet* 2011, **7**:e1002416.
5. Marlow H, Tosches MA, Tomer R, Steinmetz PRH, Lauri A, Larsson T, Arendt D: **Larval body patterning and apical organs are conserved in animal evolution.** *BMC Biol* 2013. (submitted for publication).
6. Santagata S, Resh C, Hejnal A, Martindale MQ, Passamanek YJ: **Development of the larval anterior neurogenic domains of *Terebratalia transversa* (Brachiopoda) provides insights into the diversification of larval apical organs and the spiralian nervous system.** *EvoDevo* 2012, **3**:3.
7. Angerer LM, Yaguchi S, Angerer RC, Burke RD: **The evolution of nervous system patterning: insights from sea urchin development.** *Development* 2011, **138**:3613-3623.
8. Lowe CJ, Wu M, Salic A, Evans L, Lander E, Stange-Thomann N, Gruber CE, Gerhart J, Kirschner M: **Anteroposterior patterning in hemichordates and the origins of the chordate nervous system.** *Cell* 2003, **113**:853-865.
9. Pani AM, Mullarkey EE, Aronowicz J, Assimacopoulos S, Grove EA, Lowe CJ: **Ancient deuterostome origins of vertebrate brain signalling centres.** *Nature* 2012, **483**:289-294.
- In this paper, Pani *et al.* use gene expression and functional data to demonstrate the conservation of the vertebrate brain signalling centres in a basal deuterostome, the hemichordate worm *Saccoglossus kowalevskii*. In particular, they propose the homology of the invertebrate apical FGF signalling centre with the vertebrate anterior neural ridge, and they identify a ZLI-like centre at the boundary between the proboscis and the collar.
10. Yu J-K, Satou Y, Holland ND, Shin-I T, Kohara Y, Satoh N, Bronner-Fraser M, Holland LZ: **Axial patterning in cephalochordates and the evolution of the organizer.** *Nature* 2007, **445**:613-617.

11. Wilson SW, Houart C: **Early steps in the development of the forebrain.** *Dev Cell* 2004, **6**:167-181.
12. Sinigaglia C, Busengdal H, Leclère L, Technau U, Rentzsch F: **The bilaterian head patterning gene *six3/6* controls aboral domain development in a cnidarian.** *PLoS Biol* 2013, **11**:e1001488.
- This paper, together with [13**], demonstrates the conservation of the apical patterning system in Cnidaria. Sinigaglia *et al.* show that the bilaterian apical plate genes are expressed in the aboral region of *Nematostella*, and that *six3* is crucial for the specification of apical identity.
13. Marlow H, Matus DQ, Martindale MQ: **Ectopic activation of the canonical Wnt signaling pathway affects ectodermal patterning along the primary axis during larval development in the anthozoan *Nematostella vectensis*.** *Dev Biol* 2013 <http://dx.doi.org/10.1016/j.ydbio.2013.05.022>.
- This paper, together with [12**], demonstrates the conservation of the apical patterning system in Cnidaria. Marlow *et al.* show that in *Nematostella* canonical Wnt signalling is essential for ectodermal patterning along the apical-blastoporal axis.
14. Oliver G, Mailhos A, Wehr R, Copeland NG, Jenkins NA, Gruss P: ***Six3*, a murine homologue of the *sine oculis* gene, demarcates the most anterior border of the developing neural plate and is expressed during eye development.** *Development* 1995, **121**:4045-4055.
15. Jessell TM: **Neuronal specification in the spinal cord: inductive signals and transcriptional codes.** *Nature Rev Genet* 2000, **1**:20-29.
16. Wheeler SR, Carrico ML, Wilson BA, Skeath JB: **The *Tribolium* columnar genes reveal conservation and plasticity in neural precursor patterning along the embryonic dorsal-ventral axis.** *Dev Biol* 2005, **279**:491-500.
17. Denes AS, Jékely G, Steinmetz PRH, Raible F, Snyman H, Prud'homme B, Ferrier DEK, Balavoine G, Arendt D: **Molecular architecture of annelid nerve cord supports common origin of nervous system centralization in bilateria.** *Cell* 2007, **129**:277-288.
18. Puelles L, Rubenstein JLR: **Forebrain gene expression domains and the evolving prosomeric model.** *Trends Neurosci* 2003, **26**:469-476.
19. Steinmetz PRH, Zelada-González F, Burgdorf C, Wittbrodt J, Arendt D: **Polychaete trunk neuroectoderm converges and extends by mediolateral cell intercalation.** *Proc Natl Acad Sci U S A* 2007, **104**:2727-2732.
20. Arendt D, Nübler-Jung K: **Dorsal or ventral: similarities in fate maps and gastrulation patterns in annelids, arthropods and chordates.** *Mech Dev* 1997, **61**:7-21.
21. Arendt D, Tosches MA, Marlow H: **From nerve ring to nerve cord — evolution of the nervous system.** *Nat Rev Neurosci* 2013. (submitted for publication).
22. Matus DQ, Magie CR, Pang K, Martindale MQ, Thomsen GH: **The Hedgehog gene family of the cnidarian *Nematostella vectensis*, and implications for understanding metazoan Hedgehog pathway evolution.** *Dev Biol* 2008, **313**:501-518.
23. Vigh B, Vigh-Teichmann I: **Actual problems of the cerebrospinal fluid-contacting neurons.** *Microsc Res Tech* 1998, **41**:57-83.
24. Lacalli TC: **Structure and development of the apical organ in trochophores of *Spirobranchus polycer*. *Phyllodoce maculata* and *Phyllodoce mucosa* (Polychaeta).** *Proc R Soc B: Biol Sci* 1981, **212**:381-402.
25. Tessmar-Raible K, Raible F, Christodoulou F, Guy K, Rembold M, Hausen H, Arendt D: **Conserved sensory-neurosecretory cell types in annelid and fish forebrain: insights into hypothalamus evolution.** *Cell* 2007, **129**:1389-1400.
26. Byrne M, Nakajima Y, Chee FC, Burke RD: **Apical organs in echinoderm larvae: insights into larval evolution in the Ambulacraria.** *Evol Dev* 2007, **9**:432-445.
27. Mathers PH, Grinberg a, Mahon KA, Jamrich M: **The *Rx* homeobox gene is essential for vertebrate eye development.** *Nature* 1997, **387**:603-607.
28. Kimura A, Singh D, Wawrousek EF, Kikuchi M, Nakamura M, Shinohara T: **Both *PCE-1/RX* and *OTX/CRX* interactions are necessary for photoreceptor-specific gene expression.** *J Biol Chem* 2000, **275**:1152-1160.
29. Pan Y, Martinez-De Luna RI, Lou C-H, Nekkalapudi S, Kelly LE, Sater AK, El-Hodiri HM: **Regulation of photoreceptor gene expression by the retinal homeobox (*Rx*) gene product.** *Dev Biol* 2010, **339**:494-506.
30. Manavathi B, Peng S, Rayala SK, Talukder AH, Wang MH, Wang R-A, Balasenthil S, Agarwal N, Frishman LJ, Kumar R: **Repression of *Six3* by a corepressor regulates rhodopsin expression.** *Proc Natl Acad Sci U S A* 2007, **104**:13128-13133.
31. Arendt D, Tessmar-Raible K, Snyman H, Dorrestein AW, Wittbrodt J: **Ciliary photoreceptors with a vertebrate-type opsin in an invertebrate brain.** *Science* 2004, **306**:869-871.
32. Passamaneck YJ, Furchheim N, Hejnol A, Martindale MQ, Lütter C: **Ciliary photoreceptors in the cerebral eyes of a protostome larva.** *EvoDevo* 2011, **2**:6.
33. Ooka S, Katow T, Yaguchi S, Yaguchi J, Katow H: **Spatiotemporal expression pattern of an encephalopsin orthologue of the sea urchin *Hemicentrotus pulcherrimus* during early development, and its potential role in larval vertical migration.** *Dev Growth Differ* 2010, **52**:195-207.
- To date, this is the only paper showing the expression of c-opsins in the apical plate of a basal deuterostome larva, making the link with previous data from protostomes.
34. Vopalensky P, Pergner J, Liegertova M, Benito-Gutierrez E, Arendt D, Kozmik Z: **Molecular analysis of the amphioxus frontal eye unravels the evolutionary origin of the retina and pigment cells of the vertebrate eye.** *Proc Natl Acad Sci U S A* 2012, **109**:15383-15388.
- For a very long time the supposed homology of the amphioxus frontal eye and the vertebrate retina has relied exclusively on morphology. Here Vopalensky and colleagues show for the first time the expression of ciliary opsins in the amphioxus frontal eye photoreceptors. Moreover, they use expression data for transcription factors and projection data to corroborate the homology of frontal eye and vertebrate retina.
35. Casarosa S, Andreazzoli M, Simeone A, Barsacchi G: ***Xrx1*, a novel *Xenopus* homeobox gene expressed during eye and pineal gland development.** *Mech Dev* 1997, **61**:187-198.
36. Ghanbari H, Seo HC, Fjose A, Brändli AW: **Molecular cloning and embryonic expression of *Xenopus* *Six* homeobox genes.** *Mech Dev* 2001, **101**:271-277.
37. Mano H, Fukada Y: **A median third eye: pineal gland retraces evolution of vertebrate photoreceptive organs.** *Photochem Photobiol* 2007, **83**:11-18.
38. Lamb TD: **Evolution of phototransduction, vertebrate photoreceptors and retina.** *Prog Retinal Eye Res* 2013 <http://dx.doi.org/10.1016/j.preteyeres.2013.06.001>.
- In this comprehensive review, Trevor Lamb explores in great detail all the steps that led to the evolution of the sophisticated vertebrate eyes starting from the first metazoan opsins.
39. Rohde K, Klein DC, Möller M, Rath MF: ***Rax*: developmental and daily expression patterns in the rat pineal gland and retina.** *J Neurochem* 2011, **118**:999-1007.
40. Arendt D, Hausen H, Purschke G: **The “division of labour” model of eye evolution.** *Philos Trans R Soc Lond Ser B: Biol Sci* 2009, **364**:2809-2817.
41. VanDunk C, Hunter LA, Gray PA: **Development, maturation, and necessity of transcription factors in the mouse suprachiasmatic nucleus.** *J Neurosci* 2011, **31**:6457-6467.
- Here VanDunk *et al.* follow with spatio-temporal accuracy the development of the suprachiasmatic nucleus (SCN) in the mouse. They show that the SCN develops from *six3+ six6+ rx+* progenitors, and that the conditional knockout of *six3* is sufficient to completely eliminate the SCN.
42. Clark DD, Gorman MR, Hatori M, Meadows JD, Panda S, Mellon PL: **Aberrant development of the suprachiasmatic nucleus and circadian rhythms in mice lacking the homeodomain protein *Six6*.** *J Biol Rhythms* 2013, **28**:15-25.
43. Medina-Martinez O, Amaya-Manzanares F, Liu C, Mendoza M, Shah R, Zhang L, Behringer RR, Mahon KA, Jamrich M: **Cell-autonomous requirement for *rx* function in the mammalian retina and posterior pituitary.** *PLoS ONE* 2009, **4**:e4513.

44. Lu F, Kar D, Gruenig N, Zhang ZW, Cousins N, Rodgers HM, Swindell EC, Jamrich M, Schuurmans C, Mathers PH *et al.*: **Rax is a selector gene for mediobasal hypothalamic cell types.** *J Neurosci* 2013, **33**:259-272.
- This paper describes the role of rx (rax) in the development of the mammalian hypothalamus. Here they show that rax is essential for the specification of cell fates in two nuclei of the tuberal hypothalamus, the ARC and the VMH.
45. Wataya T, Ando S, Muguruma K, Ikeda H, Watanabe K, Eiraku M, Kawada M, Takahashi J, Hashimoto N, Sasai Y: **Minimization of exogenous signals in ES cell culture induces rostral hypothalamic differentiation.** *Proc Natl Acad Sci U S A* 2008, **105**:11796-11801.
 46. De Velasco B, Erclik T, Shy D, Sclafani J, Lipshitz H, McInnes R, Hartenstein V: **Specification and development of the pars intercerebralis and pars lateralis, neuroendocrine command centers in the Drosophila brain.** *Dev Biol* 2007, **302**:309-323.
 47. Blechman J, Borodovsky N, Eisenberg M, Nabel-Rosen H, Grimm J, Levkowitz G: **Specification of hypothalamic neurons by dual regulation of the homeodomain protein Orthopedia.** *Development* 2007, **134**:4417-4426.
 48. Hartenstein V: **The neuroendocrine system of invertebrates: a developmental and evolutionary perspective.** *J Endocrinol* 2006, **190**:555-570.
 49. Joly J-S, Osório J, Alunni A, Auger H, Kano S, Rétaux S: **Windows of the brain: towards a developmental biology of circumventricular and other neurohemal organs.** *Semin Cell Dev Biol* 2007, **18**:512-524.
 50. Dickinson AJG, Croll RP: **Development of the larval nervous system of the gastropod Ilyanassa obsoleta.** *J Comp Neurol* 2003, **466**:197-218.
 51. Nezhlin LP, Yushin VV: **Structure of the nervous system in the tornaria larva of Balanoglossus proterogonius (Hemichordata: Enteropneusta) and its phylogenetic implications.** *Zoomorphology* 2004, **123**:1-13.
 52. Nakanishi N, Yuan D, Jacobs DK, Hartenstein V: **Early development, pattern, and reorganization of the planula nervous system in Aurelia (Cnidaria, Scyphozoa).** *Dev Genes Evol* 2008, **218**:511-524.
 53. Ohtomi M, Fujii K, Kobayashi H: **Distribution of FMRFamide-like immunoreactivity in the brain and neurohypophysis of the lamprey, *Lampetra japonica*.** *Cell Tissue Res* 1989, **256**:581-584.
 54. Jékely G: **Global view of the evolution and diversity of metazoan neuropeptide signaling.** *Proc Natl Acad Sci U S A* 2013, **110**(21):8702-8707.
 55. Dickinson AJG, Croll RP: **Development of the larval nervous system of the gastropod Ilyanassa obsoleta.** *J Comp Neurol* 2003, **466**:197-218.
 56. Yuan D, Nakanishi N, Jacobs DK, Hartenstein V: **Embryonic development and metamorphosis of the scyphozoan Aurelia.** *Dev Genes Evol* 2008, **218**:525-539.
 57. Mastick GS, Davis NM, Andrew GL, Easter SS: **Pax-6 functions in boundary formation and axon guidance in the embryonic mouse forebrain.** *Development* 1997, **124**:1985-1997.
 58. Jones L, López-Bendito G, Gruss P, Stoykova A, Molnár Z: **Pax6 is required for the normal development of the forebrain axonal connections.** *Development* 2002, **129**:5041-5052.
 59. Nomura T, Holmberg J, Frisen J, Osumi N: **Pax6-dependent boundary defines alignment of migrating olfactory cortex neurons via the repulsive activity of ephrin A5.** *Development* 2006, **133**:1335-1345.
 60. Marín O, Baker J, Puelles L, Rubenstein JLR: **Patterning of the basal telencephalon and hypothalamus is essential for guidance of cortical projections.** *Development* 2002, **129**:761-773.
 61. Kawano H, Horie M, Honma S, Kawamura K, Takeuchi K, Kimura S: **Aberrant trajectory of ascending dopaminergic pathway in mice lacking Nkx2.1.** *Exp Neurol* 2003, **182**:103-112.
 62. Nóbrega-Pereira S, Kessaris N, Du T, Kimura S, Anderson SA, Marín O: **Postmitotic Nkx2-1 controls the migration of telencephalic interneurons by direct repression of guidance receptors.** *Neuron* 2008, **59**:733-745.
 63. Sebastián-Serrano A, Sandonis A, Cardozo M, Rodríguez-Tornos FM, Bovolenta P, Nieto M: **Pax6 expression in postmitotic neurons mediates the growth of axons in response to SFRP1.** *PLoS ONE* 2012, **7**:e31590.
- This study demonstrates that pax6 is necessary and sufficient to confer responsiveness of postmitotic neurons to a specific axon guidance cue.
64. Metzakopian E, Lin W, Salmon-Divon M, Dvinge H, Andersson E, Ericson J, Perlmann T, Whitsett JA, Bertone P, Ang S-L: **Genome-wide characterization of Foxa2 targets reveals upregulation of floor plate genes and repression of ventrolateral genes in midbrain dopaminergic progenitors.** *Development* 2012, **139**:2625-2634.
- This paper characterized the role of foxA2 in the mouse floorplate by looking at its targets with ChIP-seq. In particular, it is shown that foxA2 controls directly the expression of slit genes, which are essential for the guidance of axons across the ventral midline.
65. Marion J-F, Yang C, Caqueret A, Boucher F, Michaud JL: **Sim1 and Sim2 are required for the correct targeting of mammillary body axons.** *Development* 2005, **132**:5527-5537.
 66. Schweitzer J, Löhr H, Bonkowski JL, Hübscher K, Driever W: **Sim1a and Arnt2 contribute to rhombalamo-spinal axon guidance by regulating Robo2 activity via a Robo3-dependent mechanism.** *Development* 2013, **140**:93-106.
- This paper, together with [65], shows that one role of sim, conserved within vertebrates, is to regulate the expression of robo genes for the correct guidance of axons projecting from the posterior hypothalamus to the spinal cord.
67. Freer SM, Lau DC, Pearson JC, Talsky KB, Crews ST: **Molecular and functional analysis of Drosophila single-minded larval central brain expression.** *Gene Expr Patterns* 2011, **11**:533-546.
 68. Venkatesh TV, Holland ND, Holland LZ, Su MT, Bodmer R: **Sequence and developmental expression of amphioxus Amphioxus Nk2-1: insights into the evolutionary origin of the vertebrate thyroid gland and forebrain.** *Dev Genes Evol* 1999, **209**:254-259.
 69. Moret F, Christiaen L, Deyts C, Blin M, Vernier P, Joly J-S: **Regulatory gene expressions in the ascidian ventral sensory vesicle: evolutionary relationships with the vertebrate hypothalamus.** *Dev Biol* 2005, **277**:567-579.
 70. Ogasawara M, Shigetani Y, Suzuki S, Kuratani S, Satoh N: **Expression of thyroid transcription factor-1 (TTF-1) gene in the ventral forebrain and endostyle of the agnathan vertebrate, *Lampetra japonica*.** *Genesis* 2001, **30**:51-58.
 71. Holland LZ, Venkatesh TV, Gorlin A, Bodmer R, Holland ND: **Characterization and developmental expression of Amphioxus Nk2-2, an NK2 class homeobox gene from Amphioxus (Phylum Chordata; Subphylum Cephalochordata).** *Dev Genes Evol* 1998, **208**:100-105.
 72. Takacs CM, Amore G, Oliveri P, Poustka AJ, Wang D, Burke RD, Peterson KJ: **Expression of an NK2 homeodomain gene in the apical ectoderm defines a new territory in the early sea urchin embryo.** *Dev Biol* 2004, **269**:152-164.
 73. Kimura S, Hara Y, Pineau T, Fernandez-Salguero P, Fox CH, Ward JM, Gonzalez FJ: **The T/ebp null mouse: thyroid-specific enhancer-binding protein is essential for the organogenesis of the thyroid, lung, ventral forebrain, and pituitary.** *Genes Dev* 1996, **10**:60-69.
 74. Lauter GRO, Söll I, Hauptmann G: **Molecular characterization of prosomeric and intraprosomeric subdivisions of the embryonic zebrafish diencephalon.** *J Comp Neurol* 2013, **521**:1093-1118.
- Lauter *et al.* provide an high resolution map of gene expression in the developing zebrafish diencephalon. This study confirms the existence of conserved domains within the developing forebrains of vertebrates.
75. Ménard A, Grillner S: **Diencephalic locomotor region in the lamprey-afferents and efferent control.** *J Neurophysiol* 2008, **100**:1343-1353.
 76. Lacalli TC: **The dorsal compartment locomotory control system in amphioxus larvae.** *J Morphol* 2002, **252**:227-237.

77. Orrhage L, Müller MCM: **Morphology of the nervous system of Polychaeta (Annelida).** *Hydrobiologia* 2005, **535–536**:79–111.
 78. Mullins OJ, Brodfuehrer PD, Jusufovic S, Hackett JT, Fiesen WO: **Specialized brain regions and sensory inputs that control locomotion in leeches.** *J Comp Physiol A: Neuroethol Sens Neural Behav Physiol* 2012, **198**:97–108.
- Here Mullins *et al.* dissect the functional role of the most anterior part of the brain of an annelid (the supraesophageal ganglion of the leech *Hirudo verbana*) in the modulation of locomotion.
79. Tomer R, Denes AS, Tessmar-Raible K, Arendt D: **Profiling by image registration reveals common origin of annelid mushroom bodies and vertebrate pallium.** *Cell* 2010, **142**:800–809.
 80. Yamamoto K, Vernier P: **The evolution of dopamine systems in chordates.** *Front Neuroanat* 2011, **5**:21.
 81. Vidal-Gadea AG, Pierce-Shimomura JT: **Conserved role of dopamine in the modulation of behavior.** *Commun Integr Biol* 2012, **5**:440–447.
 82. Lavado A, Lagutin OV, Oliver G: **Six3 inactivation causes progressive caudalization and aberrant patterning of the mammalian diencephalon.** *Development* 2008, **135**:441–450.
 83. Conte I, Morcillo J, Bovolenta P: **Comparative analysis of Six3 and Six6 distribution in the developing and adult mouse brain.** *Dev Dyn* 2005, **234**:718–725.
 84. Fischer RM, Fontinha BM, Kirchmaier S, Steger J, Bloch S, Inoue D, Panda S, Rumpel S, Tessmar-Raible K: **Co-expression of VAL- and TMT-opsins uncovers ancient photosensory interneurons and motoneurons in the vertebrate brain.** *PLoS Biol* 2013, **11**:e1001585.
 85. Sugahara F, Aota S, Kuraku S, Murakami Y, Takio-Ogawa Y, Hirano S, Kuratani S: **Involvement of Hedgehog and FGF signalling in the lamprey telencephalon: evolution of regionalization and dorsoventral patterning of the vertebrate forebrain.** *Development* 2011, **138**:1217–1226.
 86. Diez-Roux G, Banfi S, Sultan M, Geffers L, Anand S, Rozado D, Magen A, Canidio E, Pagani M, Peluso I *et al.*: **A high-resolution anatomical atlas of the transcriptome in the mouse embryo.** *PLoS Biol* 2011, **9**:e1000582.
- On the basis of a high-throughput expression analysis, Diez-Roux and colleagues propose a new model for the organization of the mouse hypothalamus, and a new definition of the alar-basal boundary.
87. Nakanishi N, Yuan D, Hartenstein V, Jacobs DK: **Evolutionary origin of rhopalia: insights from cellular-level analyses of Otx and POU expression patterns in the developing rhopalian nervous system.** *Evol Dev* 2010, **12**:404–415.
 88. Backfisch B, Veedin Rajan VB, Fischer RM, Lohs C, Arboleda E, Tessmar-Raible K, Raible F: **Stable transgenesis in the marine annelid *Platynereis dumerilii* sheds new light on photoreceptor evolution.** *Proc Natl Acad Sci U S A* 2013, **110**:193–198.
 89. Hallbeck M, Blomqvist A: **Spinal cord-projecting vasopressinergic neurons in the rat paraventricular hypothalamus.** *J Comp Neurol* 1999, **411**:201–211.
 90. Löhr H, Ryu S, Driever W: **Zebrafish diencephalic A11-related dopaminergic neurons share a conserved transcriptional network with neuroendocrine cell lineages.** *Development* 2009, **136**:1007–1017.
 91. Lambert AM, Bonkowski JL, Masino MA: **The conserved dopaminergic diencephalospinal tract mediates vertebrate locomotor development in zebrafish larvae.** *J Neurosci* 2012, **32**:13488–13500.
 92. Fernandes AM, Fero K, Arrenberg AB, Bergeron SA, Driever W, Burgess HA: **Deep brain photoreceptors control light-seeking behavior in zebrafish larvae.** *Curr Biol* 2012, **22**:1–6.
 93. Yang N, Dong Z, Guo S: **Fezf2 regulates multilineage neuronal differentiation through activating basic helix-loop-helix and homeodomain genes in the zebrafish ventral forebrain.** *J Neurosci* 2012, **32**:10940–10948.
 94. Hobert O: **Regulation of terminal differentiation programs in the nervous system.** *Annu Rev Cell Dev Biol* 2011, **27**:681–696.
- In this review, Oliver Hobert illustrates how the combinatorial expression of 'terminal selector transcription factors' defines the different functional properties of differentiated neurons. The *de novo* expression of a terminal selector gene can explain mechanistically how new cell types evolve by neofunctionalization.
95. Matus DQ, Pang K, Daly M, Martindale MQ: **Expression of Pax gene family members in the anthozoan cnidarian, *Nematostella vectensis*.** *Evol Dev* 2007, **9**:25–38.
 96. Martindale MQ, Pang K, Finnerty JR: **Investigating the origins of triploblasty: "mesodermal" gene expression in a diploblastic animal, the sea anemone *Nematostella vectensis* (phylum, Cnidaria; class, Anthozoa).** *Development* 2004, **131**:2463–2474.
 97. De Jong DM, Hislop NR, Hayward DC, Reece-Hoyes JS, Pontynen PC, Ball EE, Miller DJ: **Components of both major axial patterning systems of the Bilateria are differentially expressed along the primary axis of a "radiate" animal, the anthozoan cnidarian *Acropora millepora*.** *Dev Biol* 2006, **298**:632–643.
 98. Christodoulou F, Raible F, Tomer R, Simakov O, Trachana K, Klaus S, Snyman H, Hannon GJ, Bork P, Arendt D: **Ancient animal microRNAs and the evolution of tissue identity.** *Nature* 2010, **463**:1084–1088.
 99. Zuber ME, Gestri G, Viczian AS, Barsacchi G, Harris WA: **Specification of the vertebrate eye by a network of eye field transcription factors.** *Development* 2003, **130**:5155–5167.
 100. Irimia M, Piñeiro C, Maeso I, Gómez-Skarmeta JL, Casares F, García-Fernández J: **Conserved developmental expression of Fezf in chordates and *Drosophila* and the origin of the Zona Limitans Intrathalamica (ZLI) brain organizer.** *EvoDevo* 2010, **1**:7.
 101. Small EM, Vokes SA, Garriock RJ, Li D, Krieg PA: **Developmental expression of the *Xenopus* Nkx2-1 and Nkx2-4 genes.** *Mech Dev* 2000, **96**:259–262.
 102. Bang AG, Papalopulu N, Kintner C, Goulding MD: **Expression of Pax-3 is initiated in the early neural plate by posteriorizing signals produced by the organizer and by posterior non-axial mesoderm.** *Development* 1997, **124**:2075–2085.
 103. Papalopulu N, Kintner C: ***Xenopus* Distal-less related homeobox genes are expressed in the developing forebrain and are induced by planar signals.** *Development* 1993, **117**:961–975.
 104. Schröder R, Eckert C, Wolff C, Tautz D: **Conserved and divergent aspects of terminal patterning in the beetle *Tribolium castaneum*.** *Proc Natl Acad Sci U S A* 2000, **97**:6591–6596.
 105. Vigh B, Manzano MJ, Zádori A, Frank CL, Lukáts A, Röhlisch P, Szél A, Dávid C: **Nonvisual photoreceptors of the deep brain, pineal organs and retina.** *Histol Histopathol* 2002, **17**:555–590.
 106. Butler AB, Hodos W: *Comparative Vertebrate Neuroanatomy – Evolution and Adaptation.* Wiley-Interscience; 2005.
 107. Eagleson GW, Harris WA: **Mapping of the presumptive brain regions in the neural plate of *Xenopus laevis*.** *J Neurobiol* 1990, **21**:427–440.
 108. Staudt N, Houart C: **The prethalamus is established during gastrulation and influences diencephalic regionalization.** *PLoS Biol* 2007, **5**:e69.