

Does climate limit arboreality in lungless salamanders?

GO SEE POSTER #32 FOR A PHYLOGENETIC APPROACH!

Lauren Mellenthin¹, Erica Baken¹, Dr. Dean Adams¹

¹Iowa State University, Ames, Iowa

Introduction


- Arboreality has evolved *at least* 5 times within Plethodontid salamanders. [1]
- Yet no morphological differences separate arboreal and terrestrial species. [1]
- There is minimal range overlap between the two microhabitat types. Preliminary results discovered that **71%** of the arboreal species distribution does not overlap with the terrestrial species distribution.

From these observations we tested the following hypotheses:

(*H1*) The climate models differ between arboreal & terrestrial microhabitats.

(*H2*) Arboreal species do not live in habitats suitable for terrestrial life.

Methods



Conclusions

Despite the broad scale of these climate data, we found:

- (*H1*) The arboreal niche model is significantly different from the terrestrial niche model with respect to climate and environment ($D = 0.35$, $P\text{-value} < 0.01$, $I = 0.62$, $P\text{-value} < 0.01$).
- (*H2*) About half of the terrestrial species distribution is suitable for arboreality (**51%**), while an extremely small portion of the arboreal species distribution is suitable for terrestrial life (**4%**).

Future Directions


- These results suggest climate may have influenced the macroevolution of *arboreality* in this family of salamanders.
- Understanding the climatic limits of different species can inform future climate projection models. [2]

Acknowledgments & Literature Cited

Thanks to the following: NSF for funding. Comments & feedback from Dr. Dean Adams' evolutionary morphology lab.

1. Baken and Adams. Accepted. Macroevolution of Arboreality in Salamanders. Ecology and Evolution.
2. McIntire, Kd. "Plant Climbing by Salamanders as a Compensatory Behavior in Relation to Climate." *Integrative and Comparative Biology*. 59 (2019)
3. Salamander silhouettes: Phylopic, Matt Reibold (modified by T. Michael Keesey), License link: <http://creativecommons.org/licenses/by-sa/3.0/>
4. Bonetti, R. M., and A. L. Blair. 2017. Evidence for complex life cycle constraints on salamander body form diversification. *Proc. Natl. Acad. Sci.* 114(57):9936-9941.
5. IUCN 2019. The IUCN Red List of Threatened Species. Version 2019-1. <http://www.iucnredlist.org>.
6. Pick, S.E. and R.J. Hijmans, 2017. WorldClim 2: New 1-km spatial resolution climate surfaces for global land areas. *International Journal of Climatology*.
7. IPCC, 2013. Climate Change 2013: The physical science basis. In: Stocker, T.F., Qin, D., Plattner, G.K., Tignor, M., Allen, S.K., Boschung, J., Nauels, A., Xia, Y., Bex, V., Midgley, P.M. (Eds.), Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, pp. 1535.


Terrestrial Suitability Model



4.1% Overlap

Arboreal Distribution

Terrestrial Distribution



50.7% Overlap

Arboreal Suitability Model

Figure 1. We used environmental niche models (ENM) to evaluate the overlap in species distributions and their corresponding suitable niche. Blue coloration representing > 0.5 suitability, red coloration representing 0.25-0.49 suitability, and white coloration representing 0-0.24 suitability.

