

**MATHEMATICS 2023 - INFINITE SERIES I (EXAMPLES)**  
**ACADIA UNIVERSITY**

1. QUESTIONS

**Exercise 2.** Compute the Taylor expansion of  $1/x$  about 1. Find the radius of convergence. Using Taylor Theorem, find a bound on  $|1/x - S_2(x)|$  where  $S_2$  is the sum of the first two terms over the interval  $[1, 2)$ . Can you bound the error over  $[1, 2)$  using another theorem? What result do you get?

**Exercise 3.** Will  $\sum_{k=0}^{\infty} \frac{e^k}{k!}$  converge???

**Exercise 4.** Will  $\sum_{k=0}^{\infty} \frac{k!}{(2k)!}$  converge???

**Exercise 5.** Will  $\sum_{k=2}^{\infty} \frac{(-1)^k}{\ln k}$  converge?

## 6. SOLUTIONS

**Example.**  $1/x = 1 - (x-1) + (x-1)^2 - (x-1)^3 + \dots$ . The ratio test gives us  $R = |x-1|$  and thus the ratio of convergence is 1. Taylor Theorem bounds the error after two terms by  $\frac{\frac{2}{\xi}(x-1)^2}{2} = \frac{(x-1)^2}{\xi} < \frac{(1)^2}{1} = 1$ . Using AST, we check that this is an alternating series (assuming  $1 > (x-1) > 0$  makes this easy), hence the error after two terms is given by  $(x-1)^2 < 1$ .

**Example.** We apply the ratio test,  $\frac{e^{k+1}}{(k+1)!} \frac{k!}{e^k} = \frac{e}{k+1}$  which goes to zero when  $k$  becomes big.

**Example.** Again, we apply the ratio test,  $\frac{(k+1)!}{(2k+2)!} \frac{(2k)!}{k!} = \frac{k+1}{(2k+2)(2k+1)}$  which clearly goes to zero as  $k$  becomes big.

**Example.** Applying the ratio test, we get  $\frac{\ln k}{\ln(k+1)}$  which tends to 1 as  $k$  becomes big (just use l'Hospital rule!) However, since  $\frac{1}{\ln(k+1)} < \frac{1}{\ln k}$  ( $\ln$  is monotone increasing!) the series is alternating and thus, it converges!