

# A Family of 4-point Dyadic High Resolution Subdivision Schemes

**DANIEL LEMIRE**

Research Officer

National Research Council of Canada (NRC)

email: [lemire@ondelette.com](mailto:lemire@ondelette.com)

# Subdivision: why care?

✓ multiscale approach

# Subdivision: why care?


- ✓ multiscale approach
- ✓ local interpolation

# Subdivision: why care?

- ✓ multiscale approach
- ✓ local interpolation
- ✓ compactly supported wavelets


# How Dubuc did it!

✓ 4 points  $\longrightarrow$  cubic polynomial




# How Dubuc did it!

- ✓ 4 points  $\longrightarrow$  cubic polynomial
- $\longrightarrow$  midpoint value (*dyadic interpolation*)



# How Dubuc did it!


- ✓ 4 points  $\longrightarrow$  cubic polynomial  
 $\longrightarrow$  midpoint value (*dyadic interpolation*)
- ✓ repeat at finer scales



## One follow-up...

Same story **but...**

✓ 4 points  $\longrightarrow$  cubic polynomial






## One follow-up...

Same story **but...**

- ✓ 4 points  $\longrightarrow$  cubic polynomial  
 $\longrightarrow$  midpoint **and quartile** values (*tetradic interpolation*)



## Question

Given a 4—point subdivision scheme, can we reproduce more than just cubic polynomials?

## One more simple trick?

*Recall Richardson's extrapolation:*

1. *an error of order  $p$*

$$a_{\Delta x} = a_{true} + \underbrace{k(\Delta x)^p + O(\Delta x^{p+1})}_{\text{some error}}$$

## One more simple trick?

**Recall Richardson's extrapolation:**

1. *an error of order  $p$*

$$a_{\Delta x} = a_{true} + \underbrace{k(\Delta x)^p + O(\Delta x^{p+1})}_{\text{some error}}$$


2.  $\Delta x \rightarrow \Delta x/2$

$$a_{\frac{\Delta x}{2}} = a_{true} + \underbrace{k\left(\frac{\Delta x}{2}\right)^p + O(\Delta x^{p+1})}_{\text{somewhat smaller error}}$$

3. Combine  $a_{\Delta x}$  and  $a_{\frac{\Delta x}{2}}$ ,

$$\frac{2^p \times a_{\frac{\Delta x}{2}} - a_{\Delta x}}{2^p - 1} = a_{true} + \underbrace{O(\Delta x^{p+1})}_{\text{we gain an order!}}$$

## Guessing early or coarsing it up



✓ Ample storage: why not use it early?

# HRS Algorithm

1. recopy stable data:  $y_{j+1,4k} = y_{j,2k}$  ;

# HRS Algorithm

1. recopy stable data:  $y_{j+1,4k} = y_{j,2k}$  ;
2. Tetradic scheme:
  - ✓ cubic  $p$  interpolates  $y_{j,2k-2}, y_{j,2k}, y_{j,2k+2}, y_{j,2k+4}$



# HRS Algorithm

1. recopy stable data:  $y_{j+1,4k} = y_{j,2k}$  ;

2. Tetradic scheme:

✓ cubic  $p$  interpolates  $y_{j,2k-2}, y_{j,2k}, y_{j,2k+2}, y_{j,2k+4}$

✓  $y_{j+1,4k+1}^{\text{coarse}} = p(x_{j+1,4k+1}), y_{j+1,4k+2}^{\text{fine}} = p(x_{j+1,4k+2}), y_{j+1,4k+3}^{\text{coarse}} = p(x_{j+1,4k+3})$

# HRS Algorithm

1. recopy stable data:  $y_{j+1,4k} = y_{j,2k}$  ;

2. Tetradic scheme:

✓ **cubic  $p$  interpolates**  $y_{j,2k-2}, y_{j,2k}, y_{j,2k+2}, y_{j,2k+4}$

✓  $y_{j+1,4k+1}^{\text{coarse}} = p(x_{j+1,4k+1}), y_{j+1,4k+2}^{\text{fine}} = p(x_{j+1,4k+2}), y_{j+1,4k+3}^{\text{coarse}} = p(x_{j+1,4k+3})$

1. Update midpoint :

$$y_{j+1,4k+2} = (1 - \alpha)y_{j+1,4k+2}^{\text{fine}} + \alpha y_{j,2k+1}^{\text{coarse}}.$$

# Smoothness

▷  $\alpha = 0 \Rightarrow$  Dubuc

# Smoothness

▷  $\alpha = 0 \Rightarrow \text{Dubuc}$

▷  $\text{Dubuc} \Rightarrow C^1$ .

▷ For  $-25/56 < \alpha < 15/32$ , the HRS schemes are  $C^1$ .

# Reproduced polynomials

- ▷ HRS always reproduce cubic polynomials.

# Reproduced polynomials

- ▷ HRS always reproduce cubic polynomials.
- ▷ Reproduce **quartic** polynomials when  $\alpha = -3/32$ .

# Reproduced polynomials

- ▷ HRS always reproduce cubic polynomials.
- ▷ Reproduce **quartic** polynomials when  $\alpha = -3/32$ .

$y_{j-1,k} = p_4(x_{j-1,k})$  where  $p_4(x) = a_4x^4 + a_3x^3 + a_2x^2 + a_1x + a_0$ , then

✓  $y_{j,2k+1}^{\text{coarse}} = p_4(x_{j,2r+1}) - \frac{105a_4}{2^{4j}}$

✓  $y_{j,2k+1}^{\text{fine}} = p_4(x_{j,2r+1}) - \frac{9a_4}{2^{4j}}$

# Initialisation

Need subdivision scheme such that  $y_{j-1,k} = p_4(x_{j-1,k}) \implies$

✓  $y_{j,2k+1} = p_4(x_{j,2k+1}) - \frac{105a_4}{16 \times 2^{4j}}$

✓  $y_{j,2k} = p_4(x_{j,2k})$

(Possible with a 5–point s.s.)



# Reproducing quartic polynomials

▷ subdivision  $\implies$  5-point scheme

# Reproducing quartic polynomials

- ▷ subdivision  $\implies$  5-point scheme
- ▷ HRS ( $\alpha = -3/32$ )  $\implies$  5-point initialization + 4-point for the rest

# How does it compare to Vector Subdivision Schemes?

✓ Vector  $\Rightarrow$  several fixed value per node

# How does it compare to Vector Subdivision Schemes?

- ✓ Vector  $\Rightarrow$  several fixed value per node
- ✓ HRS  $\Rightarrow$  one value per node but **allowed to changed over time**

## Conclusion (a comparative table)

scheme	regularity	reproduced polynomials
Dubuc	$C^1$	cubic
Deslauriers-Dubuc	$C^1$	cubic
Dyn-Gregory-Levin	up to $C^1$	up to cubic
Hassan et al.	$C^2$	quadratic
presented HRS	up to $C^1$	cubic to quartic

# Key Mathematical Argument

Fourier  $\implies$

$$P(z) = \sum_k y_{j,k} z^k$$

# Key Mathematical Argument

Fourier  $\implies$

$$P(z) = \sum_k y_{j,k} z^k$$

and

$$\Gamma(z) = 1 + \frac{9(z + z^{-1})}{16} - \frac{9(z^3 + z^{-3})}{16}$$

then

$$P^{j+1}(z) = \Gamma(z)P^j(z^2)$$

## Key Mathematical Argument

Fourier  $\implies$

$$P(z) = \sum_k y_{j,k} z^k$$

and

$$\Gamma(z) = 1 + \frac{9(z + z^{-1})}{16} - \frac{9(z^3 + z^{-3})}{16}$$

then


$$P^{j+1}(z) = \Gamma(z) P^j(z^2)$$

HRS requires 2 trig. poly.

$$P^{j+1}(z) = \Gamma_1(z) P^j(z^2) + \Gamma_2(z) P^j(-z^2)$$



## Bonus material 1 - nice pictures



Derivatives of the fundamental functions for  $\alpha = -0.2$  (continuous line),  $\alpha = 0$  (dash-dot line), and  $\alpha = 0.15$  (dashed line).

## Bonus material 2 - intermediate result

- ▷ (Dyn) Given trigonometric polynomials  $\Gamma_1(z)$  and  $\Gamma_2(z)$ , the HRS scheme defined by


$$P^{j+1}(z) = \Gamma_1(z)P^j(z^2) + \Gamma_2(z)P^j(-z^2)$$

is  $C^n$  if the symbol corresponding to finite differences of order  $n+1$

$$dH_n^j(z) = \frac{2^{jn}(1-z)^{n+1}}{z^{n+1}}P^j(z)$$

is the symbol of a HRS scheme converging uniformly to zero for all bounded initial data.

## Bonus material 3 - more on proof



For a given  $\alpha$ , an HRS scheme is differentiable if

$$\lambda_{HR}(\alpha) = \max \{ \lambda_1(\alpha), \lambda_2(\alpha) \} < 1.$$

## Bonus material 4 - proper initialization


1. recopy data at  $x_{j+1,2k} = x_{j,k}$ :  $y_{j+1,2k} = y_{j,k}$  ;
2. extrapolate  $y_{j,k+4}$  using  $y_{j,k-2}, y_{j,k-1}, y_{j,k}, y_{j,k+1}, y_{j,k+2}$  by the formula

$$\gamma_{j,k} = 5y_{j,k-2} - 24y_{j,k-1} + 45y_{j,k} - 40y_{j,k+1} + 15y_{j,k+2}; \quad (1)$$

3. interpolate midpoint :

$$y_{j+1,2k+1} = \frac{-7y_{j,k-2} + 105y_{j,k} + 35y_{j,k+2} - 5\gamma_{j,k}}{128}.$$

# Bonus material 5 - Crunch the tetradic tree into a dyadic one



## Some references

### References

- [1] I. Daubechies, Orthonormal bases of compactly supported wavelets, Comm. Pure & Appl. Math. 41, pp. 909–996, 1988.
- [2] G. Deslauriers and S. Dubuc, Symmetric iterative interpolation processes, Constr. Approx., 5, pp. 49-68, 1989.
- [3] G. Deslauriers, S. Dubuc, and D. Lemire, Une famille d'ondelettes biorthogonales sur l'intervalle obtenue par un schéma d'interpolation itérative, Ann. Sci. Math. Québec 23 no. 1, pp. 37-48, 37-48, 1999.

- [4] F. Dubeau and R. Gervais, Procédure locale et non locale d'interpolation à l'aide de fonctions splines quadratiques, Ann. Sci. Math. Québec 23 no. 1, pp. 49-61, 1999.
- [5] S. Dubuc, Interpolation through an iterative scheme, J. Math. Anal. Appl., 114, pp. 185-204, 1986.
- [6] S. Dubuc, D. Lemire, J.-L. Merrien, Fourier analysis of 2-point Hermite interpolatory subdivision schemes, J. of Fourier Anal. Appl., 7 no. 5, 2001.
- [7] N. Dyn, Subdivision schemes in computer-aided geometric design, Advances in numerical analysis (W. Light, ed.), vol. 2, Clarendon Press, pp. 36-104, 1992.
- [8] N. Dyn, J.A. Gregory, and D. Levin, A 4-point interpolatory subdivision

scheme for curve design. Comput. Aided Geom. Design, 4, pp. 257-268, 1987.

- [9] B. Han, Approximation Properties and Construction of Hermite Interpolants and Biorthogonal Multiwavelets, Journal of Approximation Theory, 110 no.1, pp. 18-53, 2001.
- [10] M.F. Hassan, I.P. Ivriissimitzis, N.A. Dodgson, and M.A. Sabin, An Interpolating 4—Point  $C^2$  Ternary Stationary Subdivision Scheme, submitted for publication to CAGD (December 18, 2001).
- [11] C. Heil and D. Colella, Matrix refinement equations and subdivision schemes, J. Fourier Anal. Appl., 2, pp. 363-377, 1996.
- [12] F. Kuijt and R. van Damme, Stability of subdivision schemes, Mem-



orandum no. 1469, Faculty of Applied Mathematics, University of Twente, the Netherlands.

- [13] J.-L. Merrien, A family of Hermite interpolants by bisection algorithms. Numer. Algorithms 2, pp. 187-200, 1992.
- [14] J.-L. Merrien, Interpolants d'Hermite  $C^2$  obtenus par subdivision. M2An Math. Model. Numer. Anal. 33, pp. 55-65, 1999.
- [15] G. Plonka, Approximation order provided by refinable function vectors, Constr. Approx. 13, pp. 221-244, 1997.