





PEKING UNIVERSITY

JUDGE ONLINE FOR ACM/ICPC



Online Judge	Problem Set	Authors	Online Contests	User
Web Board Home Page F.A.Qs Statistical Charts	Problems Submit Problem Online Status Prob.ID: <input style="width: 50px;" type="text"/> <input type="button" value="Go"/>	Register Update your info Authors ranklist <input style="width: 50px;" type="text"/> <input type="button" value="Search"/>	Current Contest Past Contests Scheduled Contests Award Contest	sidepi Log Out Mail:2(0) Login Log Archive

Sequence

Language: ▼

Time Limit: 6000MS **Memory Limit:** 65536K
Total Submissions: 9776 **Accepted:** 3252

Description

Given m sequences, each contains n non-negative integer. Now we may select one number from each sequence to form a sequence with m integers. It's clear that we may get n^m this kind of sequences. Then we can calculate the sum of numbers in each sequence, and get n^m values. What we need is the smallest n sums. Could you help us?

Input

The first line is an integer T , which shows the number of test cases, and then T test cases follow. The first line of each case contains two integers m, n ($0 < m \leq 100, 0 < n \leq 2000$). The following m lines indicate the m sequence respectively. No integer in the sequence is greater than 10000.

Output

For each test case, print a line with the smallest n sums in increasing order, which is separated by a space.

Sample Input

```

1
2 3
1 2 3
2 2 3
  
```

Sample Output

3 3 4

Source

POJ Monthly,Guang Lin

[\[Submit\]](#) [\[Go Back\]](#) [\[Status\]](#) [\[Discuss\]](#)



[Home Page](#)



[Go Back](#)



[To top](#)

All Rights Reserved 2003-2013 Ying Fuchen,Xu Pengcheng,Xie Di
Any problem, Please [Contact Administrator](#)