



Online Judge	Problem Set	Authors	Online Contests	User
Web Board Home Page F.A.Qs Statistical Charts	Problems Submit Problem Online Status Prob.ID: <input type="text"/> <input type="button" value="Go"/>	Register Update your info Authors ranklist <input type="text"/> <input type="button" value="Search"/>	Current Contest Past Contests Scheduled Contests Award Contest	lydliyudong Log Out Mail:5(0) Login Log Archive

To the Max

Language: ▾

Time Limit: 1000MS

Memory Limit: 10000K

Total Submissions: 49646

Accepted: 26308

Description

Given a two-dimensional array of positive and negative integers, a sub-rectangle is any contiguous sub-array of size 1×1 or greater located within the whole array. The sum of a rectangle is the sum of all the elements in that rectangle. In this problem the sub-rectangle with the largest sum is referred to as the maximal sub-rectangle.

As an example, the maximal sub-rectangle of the array:

0 -2 -7 0

9 2 -6 2

-4 1 -4 1

-1 8 0 -2

is in the lower left corner:

9 2

-4 1

-1 8

and has a sum of 15.

Input

The input consists of an $N \times N$ array of integers. The input begins with a single positive integer N on a line by itself, indicating the size of the square two-dimensional array. This is followed by N^2 integers separated by whitespace (spaces and newlines). These are the N^2 integers of the array, presented in row-major order. That is, all numbers in the first row, left to right, then all numbers in the second row, left to right, etc. N may be as large as 100. The numbers in the array will be in the range $[-127, 127]$.

Output

Output the sum of the maximal sub-rectangle.

Sample Input

```
4
0 -2 -7 0 9 2 -6 2
-4 1 -4 1 -1
8 0 -2
```

Sample Output

```
15
```

Source

Greater New York 2001

[\[Submit\]](#) [\[Go Back\]](#) [\[Status\]](#) [\[Discuss\]](#)



[Home Page](#)



[Go Back](#)



[To top](#)